



# Full wwPDB X-ray Structure Validation Report ⓘ

May 17, 2020 – 03:11 am BST

PDB ID : 4X4S  
Title : Crystal structure of the A.fulgidus CCA-adding enzyme in complex with a G70A arginyl-tRNA minihelix ending in CCACC and CTP  
Authors : Kuhn, C.-D.; Joshua-Tor, L.  
Deposited on : 2014-12-03  
Resolution : 3.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1251 (3.30-3.22)
Ramachandran outliers	138981	1229 (3.30-3.22)
Sidechain outliers	138945	1228 (3.30-3.22)
RSRZ outliers	127900	1154 (3.30-3.22)
RNA backbone	3102	1072 (3.62-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	457	<div> <div style="width: 93%;"></div> <div>93%</div> </div>
1	C	457	<div> <div style="width: 5%; background-color: red;"></div> <div style="width: 93%; background-color: green;"></div> <div>5%</div> <div>93%</div> </div>
2	B	37	<div> <div style="width: 16%; background-color: red;"></div> <div style="width: 73%; background-color: green;"></div> <div style="width: 14%; background-color: yellow;"></div> <div style="width: 8%; background-color: orange;"></div> <div style="width: 5%; background-color: grey;"></div> <div>16%</div> <div>73%</div> <div>14%</div> <div>8%</div> <div>5%</div> </div>
2	D	37	<div> <div style="width: 68%; background-color: green;"></div> <div style="width: 14%; background-color: yellow;"></div> <div style="width: 19%; background-color: grey;"></div> <div>68%</div> <div>14%</div> <div>19%</div> </div>

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 16877 atoms, of which 8096 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CCA-adding enzyme.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	443	Total	C	H	N	O	S	0	0	0
			7344	2354	3676	639	662	13			
1	C	437	Total	C	H	N	O	S	0	0	0
			7270	2333	3641	632	651	13			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	438	SER	-	expression tag	UNP O28126
A	439	ASN	-	expression tag	UNP O28126
A	440	SER	-	expression tag	UNP O28126
A	441	SER	-	expression tag	UNP O28126
A	442	SER	-	expression tag	UNP O28126
A	443	VAL	-	expression tag	UNP O28126
A	444	ASP	-	expression tag	UNP O28126
A	445	LYS	-	expression tag	UNP O28126
A	446	LEU	-	expression tag	UNP O28126
A	447	ALA	-	expression tag	UNP O28126
A	448	ALA	-	expression tag	UNP O28126
A	449	ALA	-	expression tag	UNP O28126
A	450	LEU	-	expression tag	UNP O28126
A	451	GLU	-	expression tag	UNP O28126
A	452	HIS	-	expression tag	UNP O28126
A	453	HIS	-	expression tag	UNP O28126
A	454	HIS	-	expression tag	UNP O28126
A	455	HIS	-	expression tag	UNP O28126
A	456	HIS	-	expression tag	UNP O28126
A	457	HIS	-	expression tag	UNP O28126
C	438	SER	-	expression tag	UNP O28126
C	439	ASN	-	expression tag	UNP O28126
C	440	SER	-	expression tag	UNP O28126
C	441	SER	-	expression tag	UNP O28126
C	442	SER	-	expression tag	UNP O28126

*Continued on next page...*


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	443	VAL	-	expression tag	UNP O28126
C	444	ASP	-	expression tag	UNP O28126
C	445	LYS	-	expression tag	UNP O28126
C	446	LEU	-	expression tag	UNP O28126
C	447	ALA	-	expression tag	UNP O28126
C	448	ALA	-	expression tag	UNP O28126
C	449	ALA	-	expression tag	UNP O28126
C	450	LEU	-	expression tag	UNP O28126
C	451	GLU	-	expression tag	UNP O28126
C	452	HIS	-	expression tag	UNP O28126
C	453	HIS	-	expression tag	UNP O28126
C	454	HIS	-	expression tag	UNP O28126
C	455	HIS	-	expression tag	UNP O28126
C	456	HIS	-	expression tag	UNP O28126
C	457	HIS	-	expression tag	UNP O28126

- Molecule 2 is a RNA chain called G70A tRNA minihelix ending in CCACC.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	35	Total	C	H	N	O	P	0	0	0
			1128	331	382	133	247	35			
2	D	30	Total	C	H	N	O	P	0	0	0
			964	283	327	112	212	30			

- Molecule 3 is CYTIDINE-5'-TRIPHOSPHATE (three-letter code: CTP) (formula:  $C_9H_{16}N_3O_{14}P_3$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	H	N	O	P	
			41	9	12	3	14	3	
3	C	1	Total	C	H	N	O	P	
			41	9	12	3	14	3	

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg		
			1	1	0	0
4	C	1	Total	Mg		
			1	1	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	H	O		
			14	3	8	3	0	0
5	A	1	Total	C	H	O		
			14	3	8	3	0	0
5	A	1	Total	C	H	O		
			14	3	8	3	0	0
5	C	1	Total	C	H	O		
			14	3	8	3	0	0

- Molecule 6 is D(-)-TARTARIC ACID (three-letter code: TAR) (formula: C<sub>4</sub>H<sub>6</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	H	O	0	0
			14	4	4	6		

- Molecule 7 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula:  $C_4H_{10}O_3$ ).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	H	O	0	0
			17	4	10	3		



- Molecule 1: CCA-adding enzyme




- Chain C:  5% 93%



- Chain B: 



- Chain D:  68% 14% 19%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	111.34Å 216.45Å 58.45Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.67 – 3.25 48.67 – 3.21	Depositor EDS
% Data completeness (in resolution range)	98.4 (48.67-3.25) 92.6 (48.67-3.21)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.64 (at 3.19Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, $R_{free}$	0.220 , 0.276 0.226 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	51.6	Xtriage
Anisotropy	0.242	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 37.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.89	EDS
Total number of atoms	16877	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.99% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, PEG, CTP, TAR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.27	0/3751	0.41	0/5040
1	C	0.26	0/3712	0.40	0/4987
2	B	0.43	1/830 (0.1%)	0.71	0/1288
2	D	0.46	1/707 (0.1%)	0.73	0/1093
All	All	0.30	2/9000 (0.0%)	0.48	0/12408

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	1	G	OP3-P	-10.53	1.48	1.61
2	B	1	G	OP3-P	-10.46	1.48	1.61

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3668	3676	3668	8	0
1	C	3629	3641	3633	7	0
2	B	746	382	383	2	0
2	D	637	327	329	1	0
3	A	29	12	12	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	29	12	12	1	0
4	A	1	0	0	0	0
4	C	1	0	0	0	0
5	A	18	24	24	0	0
5	C	6	8	8	1	0
6	A	10	4	4	1	0
7	A	7	10	9	0	0
All	All	8781	8096	8082	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:17:A:O2'	2:B:19:U:OP2	1.97	0.81
1:C:358:PHE:O	1:C:363:ARG:NH2	2.13	0.80
1:A:363:ARG:NH1	1:A:378:GLU:OE2	2.30	0.64
1:A:274:ARG:NH1	1:A:332:GLU:OE2	2.36	0.58
1:C:133:HIS:HD1	3:C:501:CTP:HO2'	1.52	0.58
1:C:252:GLY:O	1:C:257:LYS:NZ	2.37	0.55
1:A:284:ARG:NH1	1:A:325:GLU:OE2	2.42	0.51
1:C:274:ARG:NH1	1:C:332:GLU:OE2	2.44	0.50
1:C:274:ARG:NH2	1:C:433:MET:O	2.45	0.50
1:A:344:ARG:NH1	1:A:378:GLU:OE1	2.49	0.46
1:A:240:HIS:ND1	6:A:504:TAR:O11	2.46	0.46
1:A:133:HIS:ND1	3:A:501:CTP:O2'	2.44	0.44
1:A:79:GLY:HA3	1:A:113:PRO:HG3	1.99	0.43
1:C:143:GLY:N	1:C:145:GLU:OE2	2.48	0.43
2:B:7:G:H3'	2:B:8:G:H5''	2.00	0.42
2:D:17:A:H2'	2:D:19:U:C5	2.54	0.42
1:A:311:GLU:HG2	1:A:382:PHE:HE1	1.85	0.40
1:C:344:ARG:NH2	5:C:503:GOL:O3	2.54	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	441/457 (96%)	420 (95%)	20 (4%)	1 (0%)	47	77
1	C	435/457 (95%)	411 (94%)	23 (5%)	1 (0%)	47	77
All	All	876/914 (96%)	831 (95%)	43 (5%)	2 (0%)	47	77

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	119	GLU
1	C	119	GLU

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	393/404 (97%)	391 (100%)	2 (0%)	88	93
1	C	387/404 (96%)	386 (100%)	1 (0%)	92	96
All	All	780/808 (96%)	777 (100%)	3 (0%)	91	94

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	224	ARG
1	A	411	TYR
1	C	217	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	B	33/37 (89%)	5 (15%)	1 (3%)
2	D	27/37 (72%)	2 (7%)	0
All	All	60/74 (81%)	7 (11%)	1 (1%)

All (7) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	B	7	G
2	B	8	G
2	B	10	A
2	B	18	G
2	B	32	G
2	D	7	G
2	D	8	G

All (1) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
2	B	17	A

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	GOL	A	506	-	5,5,5	0.35	0	5,5,5	0.32	0
6	TAR	A	504	-	3,9,9	0.59	0	6,12,12	0.80	0
5	GOL	A	507	-	5,5,5	0.42	0	5,5,5	0.16	0
5	GOL	C	503	-	5,5,5	0.40	0	5,5,5	0.19	0
5	GOL	A	503	-	5,5,5	0.37	0	5,5,5	0.22	0
3	CTP	A	501	4	23,30,30	0.79	0	30,47,47	1.39	5 (16%)
3	CTP	C	501	4	23,30,30	0.79	0	30,47,47	1.50	6 (20%)
7	PEG	A	505	-	6,6,6	0.60	0	5,5,5	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	506	-	-	2/4/4/4	-
6	TAR	A	504	-	-	4/4/12/12	-
5	GOL	A	507	-	-	4/4/4/4	-
5	GOL	C	503	-	-	2/4/4/4	-
5	GOL	A	503	-	-	2/4/4/4	-
3	CTP	A	501	4	-	8/20/38/38	0/2/2/2
3	CTP	C	501	4	-	9/20/38/38	0/2/2/2
7	PEG	A	505	-	-	2/4/4/4	-

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	501	CTP	C3'-C2'-C1'	3.78	106.67	100.98
3	C	501	CTP	C3'-C2'-C1'	3.75	106.62	100.98
3	C	501	CTP	C2-N3-C4	3.60	119.99	116.34
3	A	501	CTP	C2-N3-C4	3.13	119.51	116.34
3	C	501	CTP	PB-O3A-PA	-3.05	122.36	132.83
3	C	501	CTP	PB-O3B-PG	-2.95	122.71	132.83
3	A	501	CTP	PB-O3B-PG	-2.68	123.64	132.83

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	501	CTP	PB-O3A-PA	-2.57	124.00	132.83
3	C	501	CTP	N4-C4-N3	2.50	120.44	116.49
3	A	501	CTP	N4-C4-N3	2.49	120.43	116.49
3	C	501	CTP	C5-C4-N3	-2.25	119.12	121.72

There are no chirality outliers.

All (33) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	A	506	GOL	O1-C1-C2-C3
6	A	504	TAR	C1-C2-C3-O3
6	A	504	TAR	C1-C2-C3-C4
6	A	504	TAR	O2-C2-C3-O3
6	A	504	TAR	O2-C2-C3-C4
5	A	507	GOL	O1-C1-C2-C3
5	C	503	GOL	O1-C1-C2-C3
3	A	501	CTP	C3'-C4'-C5'-O5'
3	A	501	CTP	O4'-C4'-C5'-O5'
3	A	501	CTP	C5'-O5'-PA-O1A
3	A	501	CTP	C5'-O5'-PA-O3A
3	C	501	CTP	C3'-C4'-C5'-O5'
3	C	501	CTP	O4'-C4'-C5'-O5'
3	C	501	CTP	C5'-O5'-PA-O2A
3	C	501	CTP	C5'-O5'-PA-O3A
5	A	507	GOL	O1-C1-C2-O2
5	A	507	GOL	C1-C2-C3-O3
7	A	505	PEG	O2-C3-C4-O4
5	C	503	GOL	O1-C1-C2-O2
3	C	501	CTP	PG-O3B-PB-O1B
5	A	506	GOL	O1-C1-C2-O2
5	A	507	GOL	O2-C2-C3-O3
5	A	503	GOL	O1-C1-C2-O2
3	C	501	CTP	PA-O3A-PB-O1B
3	C	501	CTP	C5'-O5'-PA-O1A
3	A	501	CTP	PA-O3A-PB-O2B
3	A	501	CTP	PG-O3B-PB-O1B
7	A	505	PEG	O1-C1-C2-O2
3	A	501	CTP	PG-O3B-PB-O2B
3	A	501	CTP	PA-O3A-PB-O1B
3	C	501	CTP	PA-O3A-PB-O2B
3	C	501	CTP	PG-O3B-PB-O2B
5	A	503	GOL	O1-C1-C2-C3


There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	504	TAR	1	0
5	C	503	GOL	1	0
3	A	501	CTP	1	0
3	C	501	CTP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	443/457 (96%)	0.06	2 (0%) 91 90	13, 40, 70, 105	0
1	C	437/457 (95%)	0.26	21 (4%) 30 28	13, 47, 81, 127	0
2	B	35/37 (94%)	0.91	6 (17%) 1 1	27, 94, 155, 169	0
2	D	30/37 (81%)	0.56	0 100 100	44, 92, 137, 152	0
All	All	945/988 (95%)	0.20	29 (3%) 49 47	13, 45, 94, 169	0

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	18	G	5.1
1	C	1	MET	4.7
1	C	207	GLU	3.6
1	C	212	GLU	3.2
1	C	121	LYS	3.2
1	C	211	GLY	3.2
1	A	353	ARG	3.1
1	C	222	GLU	3.1
2	B	26	G	3.0
2	B	27	C	2.9
1	C	217	VAL	2.8
2	B	6	C	2.7
1	C	2	LYS	2.6
1	C	223	LYS	2.6
1	C	258	HIS	2.5
1	C	3	VAL	2.5
1	C	210	LYS	2.5
1	C	8	GLU	2.5
1	C	122	ASN	2.4
1	C	105	LYS	2.3
2	B	8	G	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	22	ARG	2.3
2	B	31	C	2.2
1	C	139	GLY	2.2
1	C	38	LEU	2.2
1	C	14	VAL	2.1
1	C	15	ILE	2.1
1	A	121	LYS	2.1
1	C	124	LYS	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	MG	C	502	1/1	0.75	0.13	87,87,87,87	0
3	CTP	C	501	29/29	0.81	0.29	75,144,173,229	0
7	PEG	A	505	7/7	0.84	0.27	40,59,84,84	0
5	GOL	C	503	6/6	0.85	0.23	28,51,90,90	0
5	GOL	A	507	6/6	0.85	0.25	28,44,68,68	0
6	TAR	A	504	10/10	0.87	0.30	40,55,75,75	0
3	CTP	A	501	29/29	0.90	0.21	36,74,104,126	0
4	MG	A	502	1/1	0.91	0.28	49,49,49,49	0
5	GOL	A	503	6/6	0.93	0.20	29,42,54,54	0
5	GOL	A	506	6/6	0.93	0.25	19,37,46,49	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around CTP C 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers ⓘ

There are no such residues in this entry.