



Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 03:30 pm BST

PDB ID : 5X6Q
Title : Crystal structure of Pseudomonas fluorescens KMO in complex with Ro 61-8048
Authors : Kim, H.T.; Hwang, K.Y.
Deposited on : 2017-02-23
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	463	<div> <div>2%</div> <div>87%</div> <div>9%</div> <div>.</div> </div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3761 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Kynurenine 3-monooxygenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	445	3449	2165	638	629	17	0	0	0


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP Q84HF5
A	0	SER	-	expression tag	UNP Q84HF5
A	252	SER	CYS	engineered mutation	UNP Q84HF5
A	461	SER	CYS	engineered mutation	UNP Q84HF5

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).



- Molecule 3 is 3,4-dimethoxy-N-[4-(3-nitrophenyl)-1,3-thiazol-2-yl]benzenesulfonamide (three-letter code: 7ZR) (formula: $C_{17}H_{15}N_3O_6S_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			28	17	3	6	2		
3	A	1	Total	C	N	O	S	0	0
			28	17	3	6	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	203	Total	O	0	0
			203	203		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Kynurenine 3-monooxygenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	68.54Å 45.65Å 76.71Å 90.00° 105.14° 90.00°	Depositor
Resolution (Å)	33.65 – 1.90 38.86 – 1.90	Depositor EDS
% Data completeness (in resolution range)	97.6 (33.65-1.90) 97.7 (38.86-1.90)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.30 (at 1.89Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.196 , 0.234 0.197 , 0.234	Depositor DCC
R_{free} test set	1789 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	29.2	Xtriage
Anisotropy	0.237	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 46.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3761	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 7ZR, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.41	0/3523	0.57	1/4784 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	197	GLN	CA-CB-CG	6.43	127.56	113.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	197	GLN	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3449	0	3417	26	0
2	A	53	0	31	0	0
3	A	56	0	0	2	0
4	A	203	0	0	5	0
All	All	3761	0	3448	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:84:ARG:NH2	3:A:502:7ZR:S19	2.50	0.85
1:A:85:MET:HG2	1:A:93:PRO:HB2	1.65	0.78
1:A:184:GLU:HG2	1:A:297:LEU:HD22	1.79	0.65
1:A:437:ASP:OD1	1:A:438:LEU:N	2.30	0.64
1:A:84:ARG:HH22	3:A:502:7ZR:C15	2.11	0.63
1:A:94:ASN:O	4:A:601:HOH:O	2.17	0.60
1:A:437:ASP:N	4:A:614:HOH:O	2.43	0.51
1:A:84:ARG:HH11	1:A:215:ILE:HD11	1.75	0.51
1:A:340:SER:O	4:A:602:HOH:O	2.20	0.50
1:A:180:VAL:HG11	1:A:302:VAL:HG22	1.96	0.47
1:A:197:GLN:OE1	1:A:232:SER:HB2	2.15	0.46
1:A:215:ILE:HG22	1:A:217:PRO:HD3	1.97	0.46
1:A:344:ASN:O	1:A:348:LEU:HG	2.16	0.45
1:A:157:LYS:HE3	1:A:158:ARG:O	2.17	0.44
1:A:309:LEU:HD12	1:A:330:ASP:HB3	2.00	0.43
1:A:53:ILE:HD11	4:A:649:HOH:O	2.18	0.43
1:A:78:ALA:HB2	1:A:109:ILE:HG22	2.02	0.42
1:A:138:VAL:HB	1:A:175:ALA:HB1	2.02	0.42
1:A:399:ARG:NH1	1:A:454:LEU:O	2.53	0.41
1:A:193:TYR:HA	1:A:237:LEU:O	2.20	0.41
1:A:239:LEU:HB3	1:A:253:PHE:CE2	2.55	0.41
1:A:396:GLN:HE22	1:A:451:THR:HG22	1.86	0.41
1:A:113:ARG:NH1	1:A:228:ASN:OD1	2.40	0.41
1:A:66:ARG:NH1	4:A:608:HOH:O	2.33	0.41
1:A:182:LEU:CD2	1:A:299:THR:HG23	2.51	0.40
1:A:182:LEU:HA	1:A:299:THR:CG2	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	439/463 (95%)	428 (98%)	9 (2%)	2 (0%)	29	18

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	143	GLN
1	A	398	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	359/373 (96%)	356 (99%)	3 (1%)	81	82

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	91	THR
1	A	155	LEU
1	A	272	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	7ZR	A	503	-	26,30,30	3.72	11 (42%)	35,43,43	2.69	11 (31%)
3	7ZR	A	502	-	26,30,30	3.56	11 (42%)	35,43,43	3.29	12 (34%)
2	FAD	A	501	-	51,58,58	1.13	4 (7%)	60,89,89	2.21	7 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	7ZR	A	503	-	-	0/19/23/23	0/3/3/3
3	7ZR	A	502	-	-	3/19/23/23	0/3/3/3
2	FAD	A	501	-	-	2/30/50/50	0/6/6/6

All (26) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	503	7ZR	O27-N26	14.51	1.47	1.22
3	A	502	7ZR	O27-N26	13.61	1.45	1.22
3	A	503	7ZR	O05-C04	6.72	1.47	1.37
3	A	502	7ZR	O05-C04	6.37	1.47	1.37
3	A	503	7ZR	C20-C17	5.27	1.57	1.48
2	A	501	FAD	C4X-C10	4.69	1.43	1.38
3	A	503	7ZR	C18-S19	-4.30	1.63	1.70
3	A	502	7ZR	C18-S19	-4.20	1.64	1.70
3	A	502	7ZR	C20-C17	3.74	1.54	1.48
3	A	502	7ZR	C08-S11	3.49	1.81	1.76
3	A	503	7ZR	O12-S11	3.43	1.47	1.43
2	A	501	FAD	C4-N3	3.32	1.38	1.33
3	A	502	7ZR	O12-S11	3.27	1.47	1.43
3	A	502	7ZR	C21-C20	-3.25	1.34	1.39
3	A	502	7ZR	S11-N14	3.07	1.68	1.63
3	A	503	7ZR	C17-N16	3.04	1.47	1.37
3	A	502	7ZR	C17-N16	2.98	1.46	1.37
3	A	503	7ZR	C08-S11	2.81	1.80	1.76
3	A	503	7ZR	C21-C20	-2.47	1.35	1.39
3	A	502	7ZR	O13-S11	-2.47	1.40	1.43
3	A	502	7ZR	C24-C25	-2.44	1.33	1.38
3	A	503	7ZR	S11-N14	2.28	1.67	1.63
2	A	501	FAD	C9A-N10	2.20	1.41	1.38
3	A	503	7ZR	C24-C25	-2.11	1.34	1.38
2	A	501	FAD	C5X-N5	2.08	1.38	1.35
3	A	503	7ZR	C25-C20	2.01	1.43	1.39

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	7ZR	O13-S11-O12	-13.62	102.81	119.55
2	A	501	FAD	C4-N3-C2	12.77	125.92	115.14
3	A	503	7ZR	O13-S11-O12	-11.66	105.22	119.55
3	A	502	7ZR	C08-S11-N14	8.21	117.16	106.83
2	A	501	FAD	C4X-C4-N3	-6.78	114.16	123.43
2	A	501	FAD	C10-C4X-N5	4.87	124.62	121.26
3	A	502	7ZR	O02-C03-C04	4.53	121.72	115.41
2	A	501	FAD	C4-C4X-C10	-4.20	117.17	119.95
3	A	502	7ZR	O05-C04-C03	4.13	121.16	115.41
3	A	503	7ZR	C21-C22-N26	4.01	122.28	118.75
3	A	503	7ZR	O05-C04-C03	3.76	120.65	115.41
2	A	501	FAD	C4X-C10-N10	-3.69	116.51	120.30
3	A	502	7ZR	O02-C03-C10	-3.58	118.24	124.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	503	7ZR	O13-S11-C08	3.45	112.21	107.97
3	A	503	7ZR	O12-S11-C08	3.35	112.09	107.97
3	A	502	7ZR	O05-C04-C07	-3.17	118.67	124.12
3	A	503	7ZR	O27-N26-C22	3.13	123.22	118.80
3	A	503	7ZR	O02-C03-C04	3.05	119.65	115.41
3	A	502	7ZR	C15-N14-S11	-2.97	118.76	124.16
3	A	502	7ZR	O12-S11-C08	2.85	111.48	107.97
3	A	502	7ZR	C25-C20-C21	2.69	121.96	118.16
3	A	503	7ZR	O05-C04-C07	-2.62	119.61	124.12
3	A	502	7ZR	C06-O05-C04	-2.59	113.62	117.53
3	A	502	7ZR	C21-C22-N26	2.57	121.01	118.75
3	A	503	7ZR	O02-C03-C10	-2.48	120.12	124.37
2	A	501	FAD	C1'-N10-C9A	2.39	120.17	118.29
3	A	502	7ZR	C17-C18-S19	-2.24	109.05	111.79
3	A	503	7ZR	C25-C20-C21	2.14	121.20	118.16
3	A	503	7ZR	O13-S11-N14	2.09	111.96	106.73
2	A	501	FAD	C5A-C6A-N6A	2.08	123.52	120.35

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	502	7ZR	C10-C03-O02-C01
3	A	502	7ZR	C04-C03-O02-C01
2	A	501	FAD	O4B-C4B-C5B-O5B
3	A	502	7ZR	C09-C08-S11-O12
2	A	501	FAD	C1'-C2'-C3'-O3'


There are no ring outliers.


1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	502	7ZR	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	445/463 (96%)	0.04	11 (2%) 57 60	19, 33, 53, 69	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	246	ALA	6.4
1	A	377	VAL	4.7
1	A	459	HIS	3.7
1	A	245	ALA	3.6
1	A	398	THR	3.2
1	A	247	GLN	3.1
1	A	8	ARG	2.8
1	A	142	ARG	2.5
1	A	375	SER	2.4
1	A	458	SER	2.3
1	A	378	ALA	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	7ZR	A	502	28/28	0.95	0.10	30,38,51,54	0
3	7ZR	A	503	28/28	0.96	0.11	27,37,52,58	0
2	FAD	A	501	53/53	0.98	0.12	17,22,27,32	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 7ZR A 502:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 7ZR A 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD A 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.