



wwPDB X-ray Structure Validation Summary Report ⓘ

May 25, 2020 – 12:18 pm BST

PDB ID : 1XAE
Title : Crystal structure of wild type yellow fluorescent protein zFP538 from *Zoanthus*
Authors : Remington, S.J.; Wachter, R.M.; Yarbrough, D.K.; Branchaud, B.; Anderson, D.C.; Kallio, K.; Lukyanov, K.A.
Deposited on : 2004-08-25
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	229	 3% 38% 45% 14% . .
1	B	229	 3% 35% 45% 15% . .

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3616 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called fluorescent protein FP538.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	223	Total	C	N	O	S	0	0	0
			1766	1135	293	321	17			
1	B	222	Total	C	N	O	S	0	0	0
			1759	1130	291	321	17			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	66	CH7	LYS	CHROMOPHORE	UNP 6090875
A	66	CH7	TYR	CHROMOPHORE	UNP 6090875
A	66	CH7	GLY	CHROMOPHORE	UNP 6090875
B	66	CH7	LYS	CHROMOPHORE	UNP 6090875
B	66	CH7	TYR	CHROMOPHORE	UNP 6090875
B	66	CH7	GLY	CHROMOPHORE	UNP 6090875

- Molecule 2 is BETA-MERCAPTOETHANOL (three-letter code: BME) (formula: C_2H_6OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total 4	C 2	O 1	S 1	0	0
2	B	1	Total 4	C 2	O 1	S 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	43	Total 43	O 43	0	0
3	B	40	Total 40	O 40	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	121.16Å 121.16Å 111.69Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	6.00 – 2.70 39.66 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.9 (6.00-2.70) 99.9 (39.66-2.60)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.30 (at 2.61Å)	Xtriage
Refinement program	TNT 5E	Depositor
R, R_{free}	0.205 , 0.255 0.210 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	48.1	Xtriage
Anisotropy	0.057	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 104.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.034 for -h,-k,l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3616	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CSO, BME, NFA, CH7

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.45	15/1761 (0.9%)	1.08	13/2375 (0.5%)
1	B	0.81	14/1754 (0.8%)	1.09	17/2368 (0.7%)
All	All	1.18	29/3515 (0.8%)	1.08	30/4743 (0.6%)

The worst 5 of 29 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	157	LYS	CB-CG	49.81	2.87	1.52
1	B	10	GLU	CD-OE2	6.23	1.32	1.25
1	A	10	GLU	CD-OE2	6.12	1.32	1.25
1	A	117	GLU	CD-OE2	6.05	1.32	1.25
1	A	150	GLU	CD-OE2	6.03	1.32	1.25

The worst 5 of 30 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	226	PHE	C-N-CD	-7.24	104.67	120.60
1	A	101	ASP	CB-CG-OD2	-7.00	112.00	118.30
1	B	208	ASP	CB-CG-OD2	-6.87	112.12	118.30
1	B	69	ASP	CB-CG-OD2	-6.79	112.19	118.30
1	B	173	ASP	CB-CG-OD2	-6.47	112.47	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1766	0	1687	148	0
1	B	1759	0	1672	150	0
2	A	4	0	5	1	0
2	B	4	0	5	0	0
3	A	43	0	0	4	0
3	B	40	0	0	6	0
All	All	3616	0	3369	297	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 43.

The worst 5 of 297 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:76:PRO:HG2	1:B:79:ILE:HD12	1.18	1.16
1:A:28:VAL:HB	1:A:50:GLU:HB2	1.25	1.16
1:B:141:MET:HE3	1:B:172:LYS:HG3	1.33	1.09
1:A:76:PRO:HG2	1:A:79:ILE:HG13	1.42	1.00
1:A:62:SER:HB3	1:A:204:LEU:HD21	1.48	0.95

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	216/229 (94%)	204 (94%)	11 (5%)	1 (0%)	29	54
1	B	215/229 (94%)	207 (96%)	6 (3%)	2 (1%)	17	40
All	All	431/458 (94%)	411 (95%)	17 (4%)	3 (1%)	22	46

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	115	VAL
1	B	212	ALA
1	B	227	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	182/194 (94%)	154 (85%)	28 (15%)	2	7
1	B	181/194 (93%)	145 (80%)	36 (20%)	1	3
All	All	363/388 (94%)	299 (82%)	64 (18%)	2	4

5 of 64 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	10	GLU
1	B	86	SER
1	B	211	ASP
1	B	13	THR
1	B	46	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	180	GLN
1	A	202	HIS
1	B	42	GLN
1	A	158	GLN
1	B	17	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

8 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CSO	A	47	1	3,6,7	0.84	0	0,6,8	0.00	-
1	NFA	A	65	1	12,12,12	1.34	2 (16%)	15,15,15	0.90	1 (6%)
1	NFA	B	65	1	12,12,12	1.51	2 (16%)	15,15,15	0.70	0
1	CH7	A	66	1	23,25,26	2.43	11 (47%)	28,34,36	1.46	4 (14%)
1	CH7	B	66	1	23,25,26	2.41	10 (43%)	28,34,36	1.46	5 (17%)
1	CSO	B	47	1	3,6,7	0.94	0	0,6,8	0.00	-
1	CSO	A	21	1	3,6,7	0.79	0	0,6,8	0.00	-
1	CSO	B	21	1	3,6,7	0.93	0	0,6,8	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSO	A	47	1	-	0/1/5/7	-
1	NFA	A	65	1	-	5/8/8/8	0/1/1/1
1	NFA	B	65	1	-	5/8/8/8	0/1/1/1
1	CH7	A	66	1	-	1/6/35/36	0/3/3/3
1	CH7	B	66	1	-	1/6/35/36	0/3/3/3
1	CSO	B	47	1	-	0/1/5/7	-
1	CSO	A	21	1	-	0/1/5/7	-
1	CSO	B	21	1	-	0/1/5/7	-

The worst 5 of 25 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	66	CH7	OH-CZ	-4.91	1.25	1.37
1	A	66	CH7	OH-CZ	-4.83	1.25	1.37
1	B	66	CH7	CE1-CZ	4.00	1.46	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	66	CH7	CE1-CZ	3.98	1.46	1.38
1	A	66	CH7	CB2-CA2	3.88	1.38	1.35

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	66	CH7	CE-NZ-CA1	-3.40	112.41	117.16
1	B	66	CH7	CE-NZ-CA1	-3.29	112.56	117.16
1	B	66	CH7	CE1-CZ-CE2	-2.97	114.76	119.77
1	A	66	CH7	CE1-CZ-CE2	-2.97	114.76	119.77
1	B	66	CH7	O2-C2-CA2	2.51	132.37	130.96

There are no chirality outliers.

5 of 12 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	65	NFA	NXT-C-CA-N
1	A	65	NFA	O-C-CA-CB
1	A	65	NFA	NXT-C-CA-CB
1	B	65	NFA	NXT-C-CA-N
1	A	66	CH7	C3-CA3-N3-C2

There are no ring outliers.

6 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	65	NFA	1	0
1	A	66	CH7	2	0
1	B	66	CH7	5	0
1	B	47	CSO	1	0
1	A	21	CSO	1	0
1	B	21	CSO	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	BME	B	587	1	3,3,3	0.98	0	1,2,2	1.44	0
2	BME	A	587	1	3,3,3	0.62	0	1,2,2	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BME	B	587	1	-	1/1/1/1	-
2	BME	A	587	1	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	587	BME	O1-C1-C2-S2
2	A	587	BME	O1-C1-C2-S2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	587	BME	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	219/229 (95%)	0.29	7 (3%) 47 48	38, 52, 75, 91	0
1	B	218/229 (95%)	0.31	7 (3%) 47 48	33, 54, 74, 97	0
All	All	437/458 (95%)	0.30	14 (3%) 47 48	33, 53, 74, 97	0

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	212	ALA	5.2
1	A	190	VAL	5.1
1	B	212	ALA	3.6
1	B	211	ASP	3.3
1	A	211	ASP	3.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CSO	B	21	7/8	0.84	0.17	49,56,62,70	0
1	CH7	B	66	23/24	0.86	0.23	29,57,100,100	0
1	CH7	A	66	23/24	0.87	0.24	42,62,100,100	0
1	NFA	B	65	12/12	0.88	0.20	43,48,61,100	0
1	CSO	B	47	7/8	0.89	0.16	48,50,73,96	0
1	CSO	A	21	7/8	0.90	0.18	34,50,56,64	0
1	NFA	A	65	12/12	0.91	0.23	34,47,66,69	0
1	CSO	A	47	7/8	0.94	0.15	40,49,81,94	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	BME	A	587	4/4	0.93	0.19	39,82,97,100	0
2	BME	B	587	4/4	0.95	0.16	54,67,73,100	0

6.5 Other polymers [i](#)

There are no such residues in this entry.