



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 16, 2021 – 11:35 AM EST

PDB ID : 6XA9  
Title : SARS CoV-2 PLpro in complex with ISG15 C-terminal domain propargylamide  
Authors : Klemm, T.; Calleja, D.J.; Richardson, L.W.; Lechtenberg, B.C.; Komander, D.  
Deposited on : 2020-06-04  
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.16
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.16


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	318	<div> <div>4%</div> <div>91%</div> <div>8%</div> <div>.</div> </div>
1	C	318	<div> <div>4%</div> <div>92%</div> <div>6%</div> <div>.</div> </div>
1	E	318	<div> <div>16%</div> <div>90%</div> <div>7%</div> <div>.</div> </div>
2	B	80	<div> <div>78%</div> <div>23%</div> </div>
2	D	80	<div> <div>3%</div> <div>80%</div> <div>19%</div> <div>.</div> </div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	80	<div><div></div><div>15%</div><div>86%</div><div>13%</div><div></div></div>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8969 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Non-structural protein 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	313	Total	C	N	O	S	0	1	0
			2430	1549	392	470	19			
1	C	313	Total	C	N	O	S	0	0	0
			2392	1519	393	461	19			
1	E	307	Total	C	N	O	S	0	0	0
			2299	1456	376	449	18			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP P0DTD1
A	-1	PRO	-	expression tag	UNP P0DTD1
C	-2	GLY	-	expression tag	UNP P0DTD1
C	-1	PRO	-	expression tag	UNP P0DTD1
E	-2	GLY	-	expression tag	UNP P0DTD1
E	-1	PRO	-	expression tag	UNP P0DTD1

- Molecule 2 is a protein called ISG15 CTD-propargylamide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	80	Total	C	N	O	S	0	0	0
			623	396	107	118	2			
2	D	79	Total	C	N	O	S	0	0	0
			589	379	100	109	1			
2	F	79	Total	C	N	O	S	0	0	0
			555	353	92	109	1			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	78	MET	-	initiating methionine	UNP P05161
B	157	AYE	GLY	engineered mutation	UNP P05161

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	78	MET	-	initiating methionine	UNP P05161
D	157	AYE	GLY	engineered mutation	UNP P05161
F	78	MET	-	initiating methionine	UNP P05161
F	157	AYE	GLY	engineered mutation	UNP P05161

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	E	1	Total	C	O	0	0
			6	3	3		
3	E	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total 1	Zn 1	0	0
4	C	1	Total 1	Zn 1	0	0
4	E	1	Total 1	Zn 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	8	Total 8	O 8	0	0
5	B	1	Total 1	O 1	0	0
5	C	16	Total 16	O 16	0	0
5	E	5	Total 5	O 5	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Non-structural protein 3




- Molecule 1: Non-structural protein 3




- Molecule 1: Non-structural protein 3




- Molecule 2: ISG15 CTD-propargylamide







- Molecule 2: ISG15 CTD-propargylamide



- Molecule 2: ISG15 CTD-propargylamide





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	124.17Å 124.17Å 238.17Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.28 – 2.90 49.28 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.9 (49.28-2.90) 100.0 (49.28-2.90)	Depositor EDS
$R_{merge}$	0.16	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.50 (at 2.91Å)	Xtriage
Refinement program	PHENIX 1.16_3549	Depositor
R, $R_{free}$	0.200 , 0.231 0.199 , 0.231	Depositor DCC
$R_{free}$ test set	1983 reflections (4.71%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	89.8	Xtriage
Anisotropy	0.287	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 73.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	8969	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	102.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.36% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AYE, GOL, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.39	0/2490	0.62	0/3396
1	C	0.36	0/2451	0.59	0/3346
1	E	0.35	0/2356	0.57	0/3221
2	B	0.36	0/630	0.55	0/854
2	D	0.33	0/596	0.52	0/812
2	F	0.27	0/560	0.47	0/768
All	All	0.36	0/9083	0.58	0/12397

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2430	0	2285	17	0
1	C	2392	0	2209	13	0
1	E	2299	0	2040	14	0
2	B	623	0	615	10	0
2	D	589	0	572	10	0
2	F	555	0	502	8	0
3	A	18	0	24	3	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	18	0	24	0	0
3	E	12	0	16	0	0
4	A	1	0	0	0	0
4	C	1	0	0	0	0
4	E	1	0	0	0	0
5	A	8	0	0	0	0
5	B	1	0	0	0	0
5	C	16	0	0	0	0
5	E	5	0	0	1	0
All	All	8969	0	8287	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:162:LEU:HD23	2:F:155:ARG:HB3	1.66	0.77
1:C:263:GLU:OE2	1:C:296:TYR:OH	2.07	0.73
1:E:140:ARG:NH1	5:E:601:HOH:O	2.27	0.68
1:C:162:LEU:HD23	2:D:155:ARG:HB3	1.75	0.67
2:F:136:PRO:HG2	2:F:139:GLU:HG3	1.78	0.66
1:A:49:SER:HB3	3:A:402:GOL:H11	1.80	0.64
1:E:157:LYS:HE3	1:E:163:GLY:HA2	1.82	0.62
1:A:157:LYS:HE3	1:A:163:GLY:HA2	1.82	0.60
2:D:82:LEU:HD11	2:D:144:PRO:HG3	1.84	0.59
1:A:263:GLU:OE2	1:A:296:TYR:OH	2.18	0.58
1:A:121:GLN:OE1	1:A:140:ARG:NH2	2.36	0.57
2:F:123:TRP:HD1	2:F:153:ARG:HB2	1.71	0.56
2:B:105:ALA:HB2	2:B:133:ASP:HB3	1.86	0.55
2:B:123:TRP:HD1	2:B:153:ARG:HB2	1.71	0.55
1:C:157:LYS:HE3	1:C:163:GLY:HA2	1.88	0.55
2:B:123:TRP:CD1	2:B:153:ARG:HB2	2.42	0.54
1:C:13:ASN:HB2	1:C:56:TYR:OH	2.07	0.54
2:F:82:LEU:HD22	2:F:100:LEU:HD11	1.91	0.52
2:D:86:VAL:HG22	2:D:148:VAL:HB	1.93	0.51
2:D:86:VAL:HB	2:D:96:TYR:CE1	2.45	0.51
1:A:13:ASN:HB2	1:A:56:TYR:OH	2.11	0.51
1:C:89:HIS:HB2	1:C:159:VAL:HG21	1.93	0.51
2:D:103:THR:HA	2:D:136:PRO:HA	1.93	0.50
1:C:263:GLU:O	1:C:273:TYR:HA	2.11	0.50

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:13:ASN:HB2	1:E:56:TYR:OH	2.12	0.50
1:E:166:ARG:NH2	2:F:151:ASN:HB3	2.27	0.48
2:D:102:GLN:OE1	2:D:106:HIS:ND1	2.42	0.48
2:F:104:VAL:HG22	2:F:137:LEU:HD23	1.95	0.48
1:A:47:HIS:HB3	3:A:402:GOL:H32	1.97	0.46
1:C:283:TYR:HB3	1:C:290:LEU:HD11	1.97	0.46
1:A:263:GLU:O	1:A:273:TYR:HA	2.16	0.46
2:D:99:ARG:H	2:D:102:GLN:NE2	2.14	0.46
2:B:86:VAL:HG22	2:B:148:VAL:HB	1.98	0.46
1:A:96:PRO:HB3	1:E:289:LEU:HD11	1.97	0.45
1:A:121:GLN:HA	1:A:136:TYR:OH	2.16	0.45
1:A:147:PHE:CE2	1:A:151:ILE:HD11	2.51	0.45
2:F:123:TRP:CD1	2:F:153:ARG:HB2	2.51	0.45
2:D:87:ARG:HA	2:D:93:SER:HA	1.98	0.45
1:E:253:LEU:HB2	1:E:296:TYR:HB3	1.99	0.45
1:A:4:THR:HG22	1:A:22:ASP:HA	2.00	0.44
1:C:120:LEU:O	1:C:136:TYR:OH	2.34	0.44
2:D:99:ARG:O	2:D:102:GLN:HG2	2.17	0.44
2:B:85:LEU:HB2	2:B:147:THR:HG22	2.00	0.44
2:D:123:TRP:CD1	2:D:153:ARG:HB3	2.52	0.44
1:E:80:LEU:HA	1:E:80:LEU:HD23	1.74	0.44
1:A:49:SER:CB	3:A:402:GOL:H11	2.47	0.43
2:B:137:LEU:HB3	2:B:142:LEU:HD12	2.00	0.43
2:B:81:PRO:HD3	2:B:99:ARG:CZ	2.48	0.43
1:E:219:GLY:HA2	1:E:232:LYS:O	2.19	0.43
1:A:89:HIS:HB2	1:A:159:VAL:HG21	2.01	0.43
1:E:147:PHE:CE2	1:E:151:ILE:HD11	2.54	0.43
1:C:80:LEU:HA	1:C:80:LEU:HD23	1.79	0.42
2:B:106:HIS:O	2:B:110:GLN:HG3	2.20	0.42
1:C:301:THR:HG23	1:C:302:ASP:OD2	2.20	0.42
1:C:147:PHE:O	1:C:151:ILE:HG13	2.19	0.41
1:A:164:ASP:HB2	2:B:154:LEU:O	2.20	0.41
1:A:95:TYR:CD1	1:A:144:ALA:HB3	2.56	0.41
1:A:95:TYR:OH	1:A:145:ALA:HA	2.20	0.41
2:B:120:ASP:HB3	2:B:155:ARG:HH21	1.86	0.41
1:E:224:CYS:SG	1:E:225:THR:N	2.94	0.41
2:F:82:LEU:HD12	2:F:82:LEU:HA	1.90	0.41
1:C:112:TYR:CD2	1:C:163:GLY:HA3	2.55	0.41
1:A:224:CYS:SG	1:A:225:THR:N	2.94	0.41
1:E:132:LEU:HA	1:E:132:LEU:HD23	1.93	0.41
1:E:247:PRO:HB3	1:E:268:TYR:OH	2.21	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:252:GLU:HA	1:E:296:TYR:O	2.20	0.41
1:C:112:TYR:CE2	1:C:163:GLY:HA3	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	312/318 (98%)	302 (97%)	10 (3%)	0	100	100
1	C	311/318 (98%)	302 (97%)	9 (3%)	0	100	100
1	E	303/318 (95%)	295 (97%)	8 (3%)	0	100	100
2	B	77/80 (96%)	76 (99%)	1 (1%)	0	100	100
2	D	76/80 (95%)	75 (99%)	1 (1%)	0	100	100
2	F	76/80 (95%)	75 (99%)	1 (1%)	0	100	100
All	All	1155/1194 (97%)	1125 (97%)	30 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	258/277 (93%)	258 (100%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	247/277 (89%)	247 (100%)	0	100	100
1	E	226/277 (82%)	226 (100%)	0	100	100
2	B	68/71 (96%)	67 (98%)	1 (2%)	65	87
2	D	61/71 (86%)	61 (100%)	0	100	100
2	F	53/71 (75%)	53 (100%)	0	100	100
All	All	913/1044 (88%)	912 (100%)	1 (0%)	93	98

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	82	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

Of 11 ligands modelled in this entry, 3 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GOL	A	403	-	5,5,5	1.06	0	5,5,5	1.02	0
3	GOL	C	403	-	5,5,5	0.86	0	5,5,5	1.02	0
3	GOL	E	503	-	5,5,5	0.83	0	5,5,5	0.89	0
3	GOL	C	401	-	5,5,5	1.17	1 (20%)	5,5,5	1.35	0
3	GOL	A	401	-	5,5,5	1.15	0	5,5,5	0.92	0
3	GOL	E	502	-	5,5,5	0.91	0	5,5,5	1.01	0
3	GOL	A	402	-	5,5,5	0.99	0	5,5,5	1.07	1 (20%)
3	GOL	C	402	-	5,5,5	0.98	0	5,5,5	0.88	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	403	-	-	0/4/4/4	-
3	GOL	C	403	-	-	2/4/4/4	-
3	GOL	E	503	-	-	0/4/4/4	-
3	GOL	C	401	-	-	2/4/4/4	-
3	GOL	A	401	-	-	2/4/4/4	-
3	GOL	E	502	-	-	0/4/4/4	-
3	GOL	A	402	-	-	2/4/4/4	-
3	GOL	C	402	-	-	0/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	401	GOL	O2-C2	-2.17	1.36	1.43

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402	GOL	C3-C2-C1	-2.01	103.89	111.70

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	401	GOL	C1-C2-C3-O3
3	A	402	GOL	O1-C1-C2-C3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
3	A	401	GOL	O2-C2-C3-O3
3	C	403	GOL	C1-C2-C3-O3
3	C	401	GOL	O1-C1-C2-C3
3	A	402	GOL	O1-C1-C2-O2
3	C	401	GOL	O1-C1-C2-O2
3	C	403	GOL	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	402	GOL	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å²)	Q<0.9
1	A	313/318 (98%)	0.37	12 (3%)	40	36	53, 78, 142, 231	0
1	C	313/318 (98%)	0.45	14 (4%)	33	29	54, 88, 157, 218	0
1	E	307/318 (96%)	0.88	52 (16%)	1	1	60, 109, 211, 255	0
2	B	79/80 (98%)	0.12	0	100	100	62, 76, 98, 110	0
2	D	78/80 (97%)	0.22	2 (2%)	56	52	64, 103, 151, 189	0
2	F	78/80 (97%)	0.77	12 (15%)	2	1	96, 158, 186, 238	0
All	All	1168/1194 (97%)	0.53	92 (7%)	12	10	53, 93, 183, 255	0

All (92) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	50	HIS	6.5
1	E	233	TYR	6.4
1	E	229	GLN	5.4
1	E	21	VAL	4.9
1	E	55	PHE	4.8
1	E	35	TYR	4.6
1	C	233	TYR	4.6
1	E	7	VAL	4.5
1	E	194	GLN	4.5
1	E	31	PHE	4.4
1	C	185	LEU	4.4
1	E	209	GLY	4.2
1	E	210	THR	4.0
1	A	233	TYR	4.0
1	E	312	THR	4.0
1	C	234	LEU	4.0
1	E	216	PHE	3.9
1	E	187	VAL	3.8
1	A	188	VAL	3.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	314	ILE	3.5
1	E	56	TYR	3.5
1	E	232	LYS	3.5
1	E	196	GLN	3.5
1	E	27	TYR	3.4
1	C	231	THR	3.4
1	E	186	ASN	3.4
2	F	124	LEU	3.3
2	F	135	LEU	3.3
1	C	230	ALA	3.2
2	F	83	SER	3.1
1	E	22	ASP	3.1
1	E	80	LEU	3.1
1	E	48	ASN	3.1
1	E	231	THR	3.1
1	E	234	LEU	3.1
1	A	313	THR	3.1
1	E	30	GLN	3.0
1	E	34	THR	3.0
1	E	220	VAL	3.0
1	E	28	GLY	2.9
1	E	181	CYS	2.9
1	E	9	THR	2.8
1	C	194	GLN	2.8
1	E	193	GLY	2.8
1	E	185	LEU	2.8
1	E	215	GLN	2.8
2	F	144	PRO	2.7
2	F	148	VAL	2.7
1	C	211	LEU	2.7
1	E	20	VAL	2.7
1	E	178	LEU	2.7
1	E	32	GLY	2.7
1	C	253	LEU	2.7
2	F	140	TYR	2.6
2	D	85	LEU	2.6
1	A	234	LEU	2.6
1	E	223	PRO	2.6
1	E	40	ASP	2.6
1	E	24	SER	2.6
1	A	227	GLY	2.5
1	E	184	VAL	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	F	84	ILE	2.5
1	A	230	ALA	2.4
1	C	193	GLY	2.4
2	F	104	VAL	2.4
1	A	236	GLN	2.4
1	C	47	HIS	2.4
1	E	42	THR	2.3
1	E	230	ALA	2.3
1	A	312	THR	2.3
1	C	216	PHE	2.3
1	A	187	VAL	2.3
1	E	19	GLN	2.3
1	E	33	PRO	2.3
2	F	96	TYR	2.3
1	E	313	THR	2.2
2	D	79	ASP	2.2
1	C	244	MET	2.2
1	E	197	THR	2.2
1	E	311	THR	2.2
1	A	216	PHE	2.2
1	A	231	THR	2.2
1	E	258	PHE	2.2
1	E	195	GLN	2.1
2	F	150	MET	2.1
1	C	187	VAL	2.1
1	E	154	TYR	2.1
1	A	235	VAL	2.1
1	E	310	TYR	2.0
2	F	139	GLU	2.0
2	F	131	LEU	2.0
1	E	8	PHE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	GOL	A	403	6/6	0.58	0.27	102,112,116,118	0
4	ZN	A	404	1/1	0.69	0.15	212,212,212,212	0
3	GOL	C	402	6/6	0.87	0.21	103,107,117,120	0
3	GOL	C	403	6/6	0.92	0.23	97,100,107,110	0
3	GOL	E	503	6/6	0.93	0.28	90,96,101,104	0
3	GOL	A	402	6/6	0.94	0.12	108,112,113,117	0
3	GOL	C	401	6/6	0.94	0.29	68,75,92,97	0
3	GOL	A	401	6/6	0.94	0.21	72,91,94,100	0
3	GOL	E	502	6/6	0.95	0.18	96,102,105,108	0
4	ZN	E	501	1/1	0.95	0.03	275,275,275,275	0
4	ZN	C	404	1/1	0.96	0.04	224,224,224,224	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.