



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 9, 2021 – 08:03 PM EST

PDB ID : 6XCS
Title : Erythromycin esterase mutant EreC H289N in its open conformation
Authors : Zielinski, M.; Berghuis, A.M.
Deposited on : 2020-06-09
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.17
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.17


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	438	
1	B	438	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6169 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called EreC.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	408	Total	C	N	O	S	Se	0	2	0
			3052	1928	514	596	4	10			
1	B	408	Total	C	N	O	S	Se	0	3	0
			3051	1912	532	593	4	10			

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MSE	-	initiating methionine	UNP C7C425
A	-18	GLY	-	expression tag	UNP C7C425
A	-17	SER	-	expression tag	UNP C7C425
A	-16	SER	-	expression tag	UNP C7C425
A	-15	HIS	-	expression tag	UNP C7C425
A	-14	HIS	-	expression tag	UNP C7C425
A	-13	HIS	-	expression tag	UNP C7C425
A	-12	HIS	-	expression tag	UNP C7C425
A	-11	HIS	-	expression tag	UNP C7C425
A	-10	HIS	-	expression tag	UNP C7C425
A	-9	SER	-	expression tag	UNP C7C425
A	-8	SER	-	expression tag	UNP C7C425
A	-7	GLY	-	expression tag	UNP C7C425
A	-6	LEU	-	expression tag	UNP C7C425
A	-5	VAL	-	expression tag	UNP C7C425
A	-4	PRO	-	expression tag	UNP C7C425
A	-3	ARG	-	expression tag	UNP C7C425
A	-2	GLY	-	expression tag	UNP C7C425
A	-1	SER	-	expression tag	UNP C7C425
A	0	HIS	-	expression tag	UNP C7C425
A	289	ASN	HIS	engineered mutation	UNP C7C425
B	-19	MSE	-	initiating methionine	UNP C7C425
B	-18	GLY	-	expression tag	UNP C7C425
B	-17	SER	-	expression tag	UNP C7C425
B	-16	SER	-	expression tag	UNP C7C425


Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-15	HIS	-	expression tag	UNP C7C425
B	-14	HIS	-	expression tag	UNP C7C425
B	-13	HIS	-	expression tag	UNP C7C425
B	-12	HIS	-	expression tag	UNP C7C425
B	-11	HIS	-	expression tag	UNP C7C425
B	-10	HIS	-	expression tag	UNP C7C425
B	-9	SER	-	expression tag	UNP C7C425
B	-8	SER	-	expression tag	UNP C7C425
B	-7	GLY	-	expression tag	UNP C7C425
B	-6	LEU	-	expression tag	UNP C7C425
B	-5	VAL	-	expression tag	UNP C7C425
B	-4	PRO	-	expression tag	UNP C7C425
B	-3	ARG	-	expression tag	UNP C7C425
B	-2	GLY	-	expression tag	UNP C7C425
B	-1	SER	-	expression tag	UNP C7C425
B	0	HIS	-	expression tag	UNP C7C425
B	289	ASN	HIS	engineered mutation	UNP C7C425

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	38	Total O 38 38	0	0
2	B	28	Total O 28 28	0	0



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	95.37Å 74.21Å 129.29Å 90.00° 96.50° 90.00°	Depositor
Resolution (Å)	44.80 – 2.40 44.80 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.5 (44.80-2.40) 99.6 (44.80-2.40)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.58 (at 2.39Å)	Xtriage
Refinement program	PHENIX 1.15.1_3469	Depositor
R, R_{free}	0.240 , 0.290 0.240 , 0.290	Depositor DCC
R_{free} test set	1697 reflections (4.83%)	wwPDB-VP
Wilson B-factor (Å ²)	51.7	Xtriage
Anisotropy	0.729	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 66.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6169	wwPDB-VP
Average B, all atoms (Å ²)	73.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.43	0/3091	0.66	2/4187 (0.0%)
1	B	0.46	0/3092	0.82	13/4190 (0.3%)
All	All	0.45	0/6183	0.74	15/8377 (0.2%)

There are no bond length outliers.

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	226	ARG	CB-CG-CD	-12.28	79.69	111.60
1	B	226	ARG	CA-CB-CG	10.16	135.76	113.40
1	B	226	ARG	NE-CZ-NH1	-8.93	115.84	120.30
1	B	355	ARG	NE-CZ-NH1	8.59	124.59	120.30
1	B	226	ARG	CG-CD-NE	8.19	129.00	111.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3052	0	2982	90	0
1	B	3051	0	2934	101	0
2	A	38	0	0	3	0
2	B	28	0	0	7	0
All	All	6169	0	5916	185	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 185 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:359:MSE:HE2	1:A:404:ILE:HG13	1.52	0.89
1:A:184[B]:SER:HB2	1:A:245:MSE:HE3	1.63	0.81
1:A:16:THR:HG22	1:A:29:GLU:HB3	1.66	0.76
1:A:184[A]:SER:HB2	1:A:245:MSE:HE3	1.66	0.76
1:B:44:GLY:HA3	1:B:290:LEU:HD22	1.68	0.76

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	408/438 (93%)	392 (96%)	14 (3%)	2 (0%)	29	41
1	B	409/438 (93%)	393 (96%)	15 (4%)	1 (0%)	47	62
All	All	817/876 (93%)	785 (96%)	29 (4%)	3 (0%)	34	48

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	220	VAL
1	A	220	VAL
1	A	127	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	318/359 (89%)	310 (98%)	8 (2%)	47	67
1	B	312/359 (87%)	302 (97%)	10 (3%)	39	59
All	All	630/718 (88%)	612 (97%)	18 (3%)	42	62

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	272	MSE
1	B	415	LEU
1	B	392	SER
1	B	14	ARG
1	B	233	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	149	GLN
1	B	131	GLN
1	B	334	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	398/438 (90%)	0.20	13 (3%) 46 45	49, 70, 98, 109	0
1	B	398/438 (90%)	0.41	22 (5%) 25 24	53, 72, 100, 120	0
All	All	796/876 (90%)	0.31	35 (4%) 34 33	49, 71, 99, 120	0

The worst 5 of 35 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	418	LEU	5.3
1	B	380	PRO	5.2
1	B	417	ASP	5.1
1	A	342	LEU	3.6
1	A	94	THR	3.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.