



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 15, 2020 – 08:37 am BST

PDB ID : 4XDD
Title : Apo [FeFe]-Hydrogenase CpI
Authors : Esselborn, J.; Hofmann, E.; Kurisu, G.; Happe, T.
Deposited on : 2014-12-19
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	583	<div> <div>8%</div> <div>97%</div> </div>
1	B	583	<div> <div>4%</div> <div>98%</div> </div>

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 19534 atoms, of which 8986 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Iron hydrogenase 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	582	Total	C	H	N	O	S	0	1	0
			9012	2840	4482	779	871	40			
1	B	581	Total	C	H	N	O	S	0	1	0
			9000	2833	4480	778	868	41			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	575	SER	-	expression tag	UNP P29166
A	576	ALA	-	expression tag	UNP P29166
A	577	TRP	-	expression tag	UNP P29166
A	578	SER	-	expression tag	UNP P29166
A	579	HIS	-	expression tag	UNP P29166
A	580	PRO	-	expression tag	UNP P29166
A	581	GLN	-	expression tag	UNP P29166
A	582	PHE	-	expression tag	UNP P29166
A	583	LYS	-	expression tag	UNP P29166
B	575	SER	-	expression tag	UNP P29166
B	576	ALA	-	expression tag	UNP P29166
B	577	TRP	-	expression tag	UNP P29166
B	578	SER	-	expression tag	UNP P29166
B	579	HIS	-	expression tag	UNP P29166
B	580	PRO	-	expression tag	UNP P29166
B	581	GLN	-	expression tag	UNP P29166
B	582	PHE	-	expression tag	UNP P29166
B	583	LYS	-	expression tag	UNP P29166

- Molecule 2 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	Fe	S	0	0
			8	4	4		
2	A	1	Total	Fe	S	0	0
			8	4	4		
2	A	1	Total	Fe	S	0	0
			8	4	4		
2	A	1	Total	Fe	S	0	0
			8	4	4		
2	B	1	Total	Fe	S	0	0
			8	4	4		
2	B	1	Total	Fe	S	0	0
			8	4	4		
2	B	1	Total	Fe	S	0	0
			8	4	4		
2	B	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total	Mg	0	0
			2	2		
3	A	2	Total	Mg	0	0
			2	2		

- Molecule 4 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe₂S₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	Fe	S	0	0
			4	2	2		
4	B	1	Total	Fe	S	0	0
			4	2	2		

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	2	Total	Cl	0	0
			2	2		
5	A	2	Total	Cl	0	0
			2	2		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	H	O	0	0
			14	3	8	3		
6	B	1	Total	C	H	O	0	0
			14	3	8	3		
6	B	1	Total	C	H	O	0	0
			14	3	8	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	704	Total	O	0	0
			704	704		
7	B	696	Total	O	0	0
			696	696		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Iron hydrogenase 1



- Molecule 1: Iron hydrogenase 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	90.06 Å 71.81 Å 103.31 Å 90.00° 97.93° 90.00°	Depositor
Resolution (Å)	38.96 – 1.60 39.49 – 1.60	Depositor EDS
% Data completeness (in resolution range)	99.6 (38.96-1.60) 99.6 (39.49-1.60)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.57 (at 1.60 Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.144 , 0.166 0.147 , 0.169	Depositor DCC
R_{free} test set	8589 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	22.6	Xtriage
Anisotropy	0.129	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 49.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	19534	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, SF4, FES, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.31	0/4608	0.49	0/6208
1	B	0.30	0/4600	0.48	0/6196
All	All	0.30	0/9208	0.48	0/12404

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4530	4482	4479	12	0
1	B	4520	4480	4482	7	0
2	A	32	0	0	1	0
2	B	32	0	0	0	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	4	0	0	0	0
4	B	4	0	0	0	0
5	A	2	0	0	0	0
5	B	2	0	0	0	0
6	A	6	8	8	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	12	16	16	0	0
7	A	704	0	0	7	5
7	B	696	0	0	2	2
All	All	10548	8986	8985	19	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:155:ASN:ND2	7:B:788:HOH:O	2.03	0.91
1:B:550:LYS:NZ	7:B:790:HOH:O	2.26	0.69
1:A:15:GLU:OE2	7:A:837:HOH:O	2.13	0.66
1:A:42:ASP:OD2	7:A:838:HOH:O	2.15	0.65
1:A:2:LYS:NZ	1:A:15:GLU:O	2.30	0.64
1:A:288:GLU:OE1	7:A:839:HOH:O	2.16	0.61
1:B:556:PHE:HE1	1:B:567:ILE:HD11	1.74	0.53
6:A:609:GOL:H2	7:A:873:HOH:O	2.09	0.51
1:A:215:LYS:NZ	7:A:844:HOH:O	2.40	0.49
1:A:556:PHE:HE1	1:A:567:ILE:HD11	1.77	0.49
1:A:320:SER:HB3	1:A:571:LYS:HD3	1.97	0.47
1:A:569:HIS:CE1	7:A:876:HOH:O	2.69	0.44
1:A:215:LYS:CE	7:A:844:HOH:O	2.67	0.43
1:A:479:LYS:NZ	1:B:39[B]:CYS:SG	2.92	0.42
1:B:432:LYS:HE3	1:B:440:LEU:HD13	2.01	0.42
1:A:226:ILE:HD11	1:A:336:TYR:CD2	2.56	0.41
1:B:336:TYR:HB3	1:B:337:PRO:HD3	2.01	0.41
1:B:277:MET:HG3	1:B:548:LEU:HD21	2.02	0.41
1:A:301:PRO:HD2	2:A:601:SF4:S2	2.61	0.40

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:723:HOH:O	7:A:799:HOH:O[2_455]	1.50	0.70
7:A:781:HOH:O	7:A:822:HOH:O[2_545]	1.93	0.27
7:A:741:HOH:O	7:A:800:HOH:O[2_455]	2.02	0.18
7:A:820:HOH:O	7:B:777:HOH:O[2_545]	2.02	0.18
7:A:745:HOH:O	7:B:759:HOH:O[2_455]	2.07	0.13

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	581/583 (100%)	566 (97%)	15 (3%)	0	100	100
1	B	580/583 (100%)	568 (98%)	12 (2%)	0	100	100
All	All	1161/1166 (100%)	1134 (98%)	27 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	497/498 (100%)	496 (100%)	1 (0%)	93	88
1	B	497/498 (100%)	496 (100%)	1 (0%)	93	88
All	All	994/996 (100%)	992 (100%)	2 (0%)	93	88

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	B	1	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 21 ligands modelled in this entry, 8 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SF4	A	603	1	0,12,12	0.00	-	-		
2	SF4	A	604	1	0,12,12	0.00	-	-		
2	SF4	B	602	1	0,12,12	0.00	-	-		
4	FES	A	606	1	0,4,4	0.00	-	-		
2	SF4	B	603	1	0,12,12	0.00	-	-		
6	GOL	B	610	-	5,5,5	0.44	0	5,5,5	0.62	0
2	SF4	A	602	1	0,12,12	0.00	-	-		
2	SF4	B	604	1	0,12,12	0.00	-	-		
6	GOL	A	609	-	5,5,5	0.41	0	5,5,5	0.61	0
6	GOL	B	611	-	5,5,5	0.38	0	5,5,5	0.34	0
2	SF4	A	601	1	0,12,12	0.00	-	-		
2	SF4	B	601	1	0,12,12	0.00	-	-		
4	FES	B	605	1	0,4,4	0.00	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SF4	A	603	1	-	-	0/6/5/5
2	SF4	A	604	1	-	-	0/6/5/5
2	SF4	B	602	1	-	-	0/6/5/5
4	FES	A	606	1	-	-	0/1/1/1
2	SF4	B	603	1	-	-	0/6/5/5
6	GOL	B	610	-	-	2/4/4/4	-
2	SF4	A	602	1	-	-	0/6/5/5
2	SF4	B	604	1	-	-	0/6/5/5
6	GOL	A	609	-	-	2/4/4/4	-
6	GOL	B	611	-	-	2/4/4/4	-
2	SF4	A	601	1	-	-	0/6/5/5
2	SF4	B	601	1	-	-	0/6/5/5
4	FES	B	605	1	-	-	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	B	611	GOL	O1-C1-C2-C3
6	A	609	GOL	O2-C2-C3-O3
6	A	609	GOL	C1-C2-C3-O3
6	B	610	GOL	C1-C2-C3-O3
6	B	610	GOL	O2-C2-C3-O3
6	B	611	GOL	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	609	GOL	1	0
2	A	601	SF4	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	582/583 (99%)	0.19	46 (7%)	12 11	18, 25, 63, 93	0
1	B	581/583 (99%)	0.07	26 (4%)	33 30	20, 28, 51, 72	0
All	All	1163/1166 (99%)	0.13	72 (6%)	20 18	18, 27, 56, 93	0

All (72) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	5	ILE	9.7
1	A	71	ILE	8.0
1	A	8	GLY	7.8
1	A	1	MET	7.7
1	A	4	ILE	7.6
1	A	3	THR	7.0
1	A	55	GLY	6.4
1	A	10	GLN	6.1
1	A	12	ASN	5.8
1	A	11	PHE	5.7
1	A	56	THR	5.3
1	A	14	ASP	5.3
1	A	582	PHE	4.8
1	A	581	GLN	4.5
1	A	22	PHE	4.5
1	A	2	LYS	4.5
1	A	53	VAL	4.4
1	A	70	MET	4.4
1	A	9	VAL	4.4
1	A	68	ASP	4.3
1	A	16	ASP	4.3
1	A	13	THR	4.3
1	A	69	GLY	4.3
1	A	66	ILE	4.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	6	ILE	4.1
1	B	581	GLN	3.9
1	A	67	GLU	3.7
1	B	504	VAL	3.6
1	A	28	ILE	3.6
1	B	437	ASN	3.5
1	A	57	GLY	3.4
1	A	54	GLU	3.4
1	B	172	ASN	3.3
1	B	580	PRO	3.2
1	B	470	VAL	3.2
1	A	58	LEU	3.1
1	A	15	GLU	3.0
1	A	127	LYS	2.9
1	A	27	ASN	2.9
1	A	579	HIS	2.9
1	A	580	PRO	2.9
1	B	566	GLU	2.9
1	A	59	VAL	2.9
1	A	233	VAL	2.9
1	B	463	ASN	2.8
1	A	72	ILE	2.7
1	B	417	PHE	2.7
1	B	471	ILE	2.7
1	B	473	GLY	2.7
1	A	17	THR	2.7
1	B	499	CYS	2.6
1	B	418	GLY	2.6
1	A	65	LEU	2.6
1	A	51	VAL	2.6
1	A	25	ASP	2.6
1	B	576	ALA	2.6
1	B	503	CYS	2.6
1	B	538	SER	2.4
1	B	440	LEU	2.4
1	A	577	TRP	2.3
1	B	496	VAL	2.3
1	B	558	LYS	2.2
1	B	577	TRP	2.2
1	A	230	ALA	2.2
1	B	299	CYS	2.2
1	B	506	GLY	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	480	PHE	2.1
1	A	381	THR	2.1
1	B	420	THR	2.1
1	B	355	CYS	2.1
1	B	507	GLY	2.0
1	A	7	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	GOL	B	611	6/6	0.89	0.25	44,56,68,82	0
3	MG	B	609	1/1	0.93	0.04	26,26,26,26	0
6	GOL	B	610	6/6	0.94	0.09	25,35,42,44	0
3	MG	A	607	1/1	0.94	0.14	45,45,45,45	0
6	GOL	A	609	6/6	0.95	0.08	24,32,41,47	0
3	MG	A	605	1/1	0.97	0.03	26,26,26,26	0
5	CL	B	607	1/1	0.98	0.03	36,36,36,36	0
2	SF4	A	604	8/8	0.99	0.06	24,25,26,27	0
5	CL	A	610	1/1	0.99	0.03	32,32,32,32	0
2	SF4	B	604	8/8	0.99	0.06	26,27,28,28	0
3	MG	B	608	1/1	0.99	0.04	28,28,28,28	0
4	FES	A	606	4/4	0.99	0.06	25,26,28,28	0
2	SF4	B	603	8/8	1.00	0.07	21,21,22,22	0
2	SF4	A	603	8/8	1.00	0.07	19,20,20,20	0
5	CL	A	608	1/1	1.00	0.12	20,20,20,20	0
2	SF4	A	602	8/8	1.00	0.09	18,19,19,19	0
2	SF4	B	602	8/8	1.00	0.09	20,21,21,21	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	SF4	A	601	8/8	1.00	0.10	18,18,19,19	0
2	SF4	B	601	8/8	1.00	0.12	21,21,21,22	0
4	FES	B	605	4/4	1.00	0.07	21,21,22,22	0
5	CL	B	606	1/1	1.00	0.14	23,23,23,23	0

6.5 Other polymers [i](#)

There are no such residues in this entry.