Full wwPDB X-ray Structure Validation Report (i)

May 22, 2020 – 05:35 am BST

PDB ID	:	4XF5
Title	:	Crystal structure of a TRAP periplasmic solute binding protein from Chromohalobacter salexigens DSM 3043 (Csal_0678), Target EFI-501078, with bound (S)-(+)-2-Amino-1-propanol.
Authors	:	Vetting, M.W.; Al Obaidi, N.F.; Toro, R.; Morisco, L.L.; Benach, J.; Wasserman, S.R.; Attonito, J.D.; Scott Glenn, A.; Chamala, S.; Chowdhury, S.; Lafleur, J.; Love, J.; Seidel, R.D.; Whalen, K.L.; Gerlt, J.A.; Almo, S.C.; Enzyme Function Initiative (EFI)
Deposited on	:	2014-12-26
Resolution	:	1.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1156 (1.46-1.46)
Clashscore	141614	1202 (1.46-1.46)
Ramachandran outliers	138981	1178 (1.46-1.46)
Sidechain outliers	138945	1178 (1.46-1.46)
RSRZ outliers	127900	1139 (1.46-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	347	
1	B	347	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 10594 atoms, of which 4908 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Twin-arginine translocation pathway signal.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	317	Total	C	H	N	O	S	0	4	0
			5049	1659	2465	433	482	10			
1	B	317	Total	C	H	N	O	S	0	0	0
			5005	1647	2443	428	477	10			

There are 46 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	4	MET	-	initiating methionine	UNP Q1QZR9
A	5	HIS	-	expression tag	UNP Q1QZR9
A	6	HIS	-	expression tag	UNP Q1QZR9
A	7	HIS	-	expression tag	UNP Q1QZR9
A	8	HIS	-	expression tag	UNP Q1QZR9
A	9	HIS	-	expression tag	UNP Q1QZR9
A	10	HIS	-	expression tag	UNP Q1QZR9
A	11	SER	-	expression tag	UNP Q1QZR9
A	12	SER	-	expression tag	UNP Q1QZR9
A	13	GLY	-	expression tag	UNP Q1QZR9
A	14	VAL	-	expression tag	UNP Q1QZR9
A	15	ASP	-	expression tag	UNP Q1QZR9
A	16	LEU	-	expression tag	UNP Q1QZR9
A	17	GLY	-	expression tag	UNP Q1QZR9
A	18	THR	-	expression tag	UNP Q1QZR9
A	19	GLU	-	expression tag	UNP Q1QZR9
A	20	ASN	-	expression tag	UNP Q1QZR9
A	21	LEU	-	expression tag	UNP Q1QZR9
A	22	TYR	-	expression tag	UNP Q1QZR9
A	23	PHE	-	expression tag	UNP Q1QZR9
A	24	GLN	-	expression tag	UNP Q1QZR9
A	25	SER	-	expression tag	UNP Q1QZR9
A	26	MET	-	expression tag	UNP Q1QZR9
B	4	MET	-	initiating methionine	UNP Q1QZR9
B	5	HIS	-	expression tag	UNP Q1QZR9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	6	HIS	-	expression tag	UNP Q1QZR9
B	7	HIS	-	expression tag	UNP Q1QZR9
B	8	HIS	-	expression tag	UNP Q1QZR9
B	9	HIS	-	expression tag	UNP Q1QZR9
B	10	HIS	-	expression tag	UNP Q1QZR9
B	11	SER	-	expression tag	UNP Q1QZR9
B	12	SER	-	expression tag	UNP Q1QZR9
B	13	GLY	-	expression tag	UNP Q1QZR9
B	14	VAL	-	expression tag	UNP Q1QZR9
B	15	ASP	-	expression tag	UNP Q1QZR9
B	16	LEU	-	expression tag	UNP Q1QZR9
B	17	GLY	-	expression tag	UNP Q1QZR9
B	18	THR	-	expression tag	UNP Q1QZR9
B	19	GLU	-	expression tag	UNP Q1QZR9
B	20	ASN	-	expression tag	UNP Q1QZR9
B	21	LEU	-	expression tag	UNP Q1QZR9
B	22	TYR	-	expression tag	UNP Q1QZR9
B	23	PHE	-	expression tag	UNP Q1QZR9
B	24	GLN	-	expression tag	UNP Q1QZR9
B	25	SER	-	expression tag	UNP Q1QZR9
B	26	MET	-	expression tag	UNP Q1QZR9

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	2	Total Cl 2 2	0	0
2	A	2	Total Cl 2 2	0	0

- Molecule 3 is (2S)-2-aminopropan-1-ol (three-letter code: 2A1) (formula: C_3H_9NO).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			5	3	1	1		
3	B	1	Total	C	N	O	0	0
			5	3	1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	249	Total	O	0	0
			249	249		
4	B	277	Total	O	0	0
			277	277		

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Twin-arginine translocation pathway signal



- Molecule 1: Twin-arginine translocation pathway signal



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α, β, γ	122.61Å 51.15Å 111.94Å 90.00° 115.96° 90.00°	Depositor
Resolution (Å)	27.56 – 1.45 100.65 – 1.45	Depositor EDS
% Data completeness (in resolution range)	90.0 (27.56-1.45) 90.0 (100.65-1.45)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.78 (at 1.45Å)	Xtriage
Refinement program	PHENIX 1.8.1_1168	Depositor
R, R_{free}	0.164 , 0.191 0.165 , 0.192	Depositor DCC
R_{free} test set	4973 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å^2)	14.9	Xtriage
Anisotropy	0.483	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.40 , 49.6	EDS
L-test for twinning[2]	$< \|L\| > = 0.44$, $< L^2 > = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.97	EDS
Total number of atoms	10594	wwPDB-VP
Average B, all atoms (Å^2)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.47% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, 2A1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|>5$	RMSZ	#$\|Z\|>5$
1	A	0.50	0/2665	0.63	1/3627 (0.0%)
1	B	0.51	0/2637	0.65	1/3587 (0.0%)
All	All	0.50	0/5302	0.64	2/7214 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	B	119	ARG	NE-CZ-NH1	5.46	123.03	120.30
1	A	119	ARG	NE-CZ-NH1	5.08	122.84	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2584	2465	2451	20	0
1	B	2562	2443	2436	18	0
2	A	2	0	0	1	0
2	B	2	0	0	1	0
3	A	5	0	9	1	0
3	B	5	0	9	2	0
4	A	249	0	0	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	277	0	0	4	0
All	All	5686	4908	4905	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:307:GLU:OE1	4:B:776:HOH:O	1.99	0.80
1:A:103[A]:SER:O	1:A:108[A]:THR:HG21	1.84	0.78
1:B:88:ARG:NH2	4:B:688:HOH:O	2.19	0.76
1:A:134:LYS:HD3	2:A:402:CL:CL	2.32	0.66
1:B:33:THR:HG23	1:B:33:THR:O	1.98	0.64
1:A:215:TRP:CZ3	3:A:403:2A1:H3B	2.33	0.63
1:B:206:GLN:O	1:B:207:THR:HB	2.04	0.58
1:A:343:ARG:HG2	1:A:347:TYR:CE2	2.40	0.56
1:A:162:LYS:HD2	1:A:231:ASN:OD1	2.06	0.55
1:B:135:GLU:HG3	4:B:611:HOH:O	2.07	0.54
1:A:103[A]:SER:O	1:A:108[A]:THR:CG2	2.54	0.51
1:A:175:ARG:HG3	1:A:175:ARG:HH11	1.76	0.50
1:A:251:GLU:HG3	4:A:727:HOH:O	2.11	0.49
1:A:32:PRO:HG3	1:A:67:GLN:CG	2.42	0.49
1:A:344:PRO:HB2	1:B:120:TYR:CE1	2.49	0.48
1:A:43:GLY:HA3	1:A:198:TRP:CE3	2.48	0.48
1:A:97:ASP:HB3	1:A:242:TRP:CZ2	2.49	0.47
1:B:43:GLY:HA3	1:B:198:TRP:CE3	2.50	0.47
1:A:83:PHE:HB2	1:A:101:MET:HG2	1.96	0.47
1:B:97:ASP:HB3	1:B:242:TRP:CZ2	2.50	0.46
1:A:253:TRP:CZ2	1:A:261:ARG:HG3	2.51	0.46
1:A:84:ARG:HG2	4:A:534:HOH:O	2.16	0.45
1:B:215:TRP:CZ3	3:B:403:2A1:H3B	2.50	0.45
1:A:220:GLU:O	1:A:224:VAL:HG22	2.17	0.45
1:B:33:THR:CG2	1:B:33:THR:O	2.66	0.43
1:A:155:ASP:HB2	1:A:156:PRO:HD2	2.00	0.43
1:B:154:TRP:HB2	1:B:242:TRP:HB3	2.02	0.42
1:B:32:PRO:O	1:B:33:THR:HG22	2.19	0.42
1:B:155:ASP:HB2	1:B:156:PRO:HD2	2.01	0.42
1:A:116:PHE:CE1	1:A:307:GLU:HG2	2.54	0.42
1:B:242:TRP:CE2	3:B:403:2A1:H3A	2.55	0.41
1:B:60:ARG:HD3	4:B:729:HOH:O	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:164:PRO:HD3	1:B:232:TYR:CE2	2.55	0.41
1:B:253:TRP:CZ2	1:B:261:ARG:HG3	2.56	0.41
1:B:134:LYS:HD3	2:B:402:CL:CL	2.58	0.40
1:A:206:GLN:CG	1:A:229:VAL:HG11	2.51	0.40
1:A:266[B]:VAL:HG23	4:A:746:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	319/347 (92%)	311 (98%)	8 (2%)	0	100	100
1	B	315/347 (91%)	305 (97%)	9 (3%)	1 (0%)	41	18
All	All	634/694 (91%)	616 (97%)	17 (3%)	1 (0%)	47	22

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	207	THR

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	268/292 (92%)	266 (99%)	2 (1%)	84	65
1	B	264/292 (90%)	262 (99%)	2 (1%)	81	62
All	All	532/584 (91%)	528 (99%)	4 (1%)	81	62

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	154	TRP
1	A	187	SER
1	B	98	ASP
1	B	295	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\|>2$	Counts	RMSZ	$\#\|Z\|>2$
3	2A1	A	403	-	4,4,4	0.41	0	3,4,4	1.38	1 (33%)
3	2A1	B	403	-	4,4,4	0.51	0	3,4,4	1.01	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	2A1	A	403	-	-	0/2/2/2	-
3	2A1	B	403	-	-	0/2/2/2	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	A	403	2A1	C3-C2-C1	-2.37	107.79	111.33

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	403	2A1	1	0
3	B	403	2A1	2	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	317/347 (91%)	-0.36	2 (0%)	89	91	9, 21, 36, 50	0
1	B	317/347 (91%)	-0.35	7 (2%)	62	65	10, 20, 36, 56	0
All	All	634/694 (91%)	-0.35	9 (1%)	75	76	9, 21, 36, 56	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	32	PRO	3.8
1	B	206	GLN	3.7
1	A	33	THR	3.2
1	B	348	VAL	2.9
1	B	203	VAL	2.8
1	A	348	VAL	2.7
1	B	207	THR	2.5
1	B	33	THR	2.4
1	B	205	LEU	2.2

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	2A1	B	403	5/5	0.97	0.09	12,13,14,21	0
3	2A1	A	403	5/5	0.98	0.09	12,13,14,18	0
2	CL	B	402	1/1	0.98	0.07	20,20,20,20	0
2	CL	A	402	1/1	0.99	0.05	19,19,19,19	0
2	CL	A	401	1/1	1.00	0.06	18,18,18,18	0
2	CL	B	401	1/1	1.00	0.07	14,14,14,14	0

6.5 Other polymers (i)

There are no such residues in this entry.