



Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2020 – 04:40 pm BST

PDB ID : 2XHI
Title : Separation-of-function mutants unravel the dual reaction mode of human 8-oxoguanine DNA glycosylase
Authors : Dalhus, B.; Forsbring, M.; Helle, I.H.; Backe, P.H.; Forstrom, R.J.; Alseth, I.; Bjoras, M.
Deposited on : 2010-06-16
Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1483 (1.56-1.56)
Clashscore	141614	1529 (1.56-1.56)
Ramachandran outliers	138981	1498 (1.56-1.56)
Sidechain outliers	138945	1495 (1.56-1.56)
RSRZ outliers	127900	1465 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	360	<div> <div>5%</div> <div> <div></div> <div>76%</div> <div>9%</div> <div>•</div> <div>12%</div> </div> </div>
2	B	15	<div> <div>20%</div> <div> <div></div> <div>87%</div> <div>7%</div> <div>7%</div> </div> </div>
3	C	15	<div> <div>33%</div> <div> <div>27%</div> <div>47%</div> <div>20%</div> <div>7%</div> </div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3494 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called N-GLYCOSYLASE/DNA LYASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	316	Total	C	N	O	S	23	1	0
			2518	1596	457	453	12			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-14	MET	-	expression tag	UNP O15527
A	-13	GLY	-	expression tag	UNP O15527
A	-12	SER	-	expression tag	UNP O15527
A	-11	HIS	-	expression tag	UNP O15527
A	-10	HIS	-	expression tag	UNP O15527
A	-9	HIS	-	expression tag	UNP O15527
A	-8	HIS	-	expression tag	UNP O15527
A	-7	HIS	-	expression tag	UNP O15527
A	-6	HIS	-	expression tag	UNP O15527
A	-5	SER	-	expression tag	UNP O15527
A	-4	GLN	-	expression tag	UNP O15527
A	-3	ASP	-	expression tag	UNP O15527
A	-2	PRO	-	expression tag	UNP O15527
A	-1	ASN	-	expression tag	UNP O15527
A	0	SER	-	expression tag	UNP O15527
A	249	CYS	LYS	engineered mutation	UNP O15527
A	253	LYS	CYS	engineered mutation	UNP O15527
A	268	ASN	ASP	engineered mutation	UNP O15527

- Molecule 2 is a DNA chain called 5'-D(*GP*GP*TP*AP*GP*AP*CP*CP*TP*GP*GP*AP*CP*GP*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	14	Total	C	N	O	P	0	0	0
			289	136	56	83	14			

- Molecule 3 is a DNA chain called 5'-D(*GP*CP*GP*TP*CP*CP*AP*(8OG)P*GP*TP*CP*TP*AP*CP*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	14	Total	C	N	O	P	0	0	0
			286	135	51	86	14			

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Ca	0	0
			1	1		
4	C	1	Total	Ca	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	346	Total	O	0	0
			346	346		
5	B	28	Total	O	0	0
			28	28		
5	C	25	Total	O	0	0
			25	25		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: N-GLYCOSYLASE/DNA LYASE



• Molecule 2: 5'-D(*GP*GP*TP*AP*GP*AP*CP*CP*TP*GP*GP*AP*CP*GP*C)-3'



• Molecule 3: 5'-D(*GP*CP*GP*TP*CP*CP*AP*(8OG)P*GP*TP*CP*TP*AP*CP*C)-3'



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	92.67Å 92.67Å 210.99Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	24.63 – 1.55 24.63 – 1.55	Depositor EDS
% Data completeness (in resolution range)	98.6 (24.63-1.55) 98.6 (24.63-1.55)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.01 (at 1.55Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.229 , 0.265 0.226 , 0.261	Depositor DCC
R_{free} test set	3880 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	20.9	Xtriage
Anisotropy	0.038	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 47.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3494	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.84% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 8OG, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.47	5/2592 (0.2%)	1.38	21/3527 (0.6%)
2	B	1.40	2/324 (0.6%)	2.17	25/498 (5.0%)
3	C	1.70	7/292 (2.4%)	2.56	30/445 (6.7%)
All	All	1.49	14/3208 (0.4%)	1.64	76/4470 (1.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	80	GLY	C-N	26.87	1.95	1.34
1	A	83	SER	C-N	-24.00	0.78	1.34
1	A	99	TYR	CD2-CE2	8.23	1.51	1.39
3	C	25	DT	C5-C7	7.63	1.54	1.50
3	C	22	DA	N3-C4	6.62	1.38	1.34
2	B	6	DA	C6-N6	6.38	1.39	1.33
3	C	25	DT	O4'-C1'	6.29	1.49	1.42
3	C	24	DG	N3-C4	5.69	1.39	1.35
3	C	24	DG	N7-C5	5.44	1.42	1.39
3	C	24	DG	N9-C8	5.36	1.41	1.37
1	A	68	GLN	CG-CD	5.25	1.63	1.51
3	C	25	DT	C2-N3	5.12	1.41	1.37
1	A	313	TRP	CE3-CZ3	5.11	1.47	1.38
2	B	7	DC	P-O5'	5.00	1.64	1.59

All (76) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	80	GLY	O-C-N	-25.89	81.27	122.70
1	A	80	GLY	CA-C-N	-19.80	73.65	117.20
1	A	131	ARG	NE-CZ-NH2	-17.77	111.42	120.30
3	C	24	DG	C2-N3-C4	-13.14	105.33	111.90
1	A	131	ARG	NE-CZ-NH1	12.56	126.58	120.30
3	C	25	DT	C2-N3-C4	-11.00	120.60	127.20
3	C	24	DG	N1-C2-N3	10.55	130.23	123.90
3	C	25	DT	N1-C2-N3	10.32	120.79	114.60
2	B	8	DC	C6-N1-C2	9.46	124.09	120.30
2	B	6	DA	C2-N3-C4	9.24	115.22	110.60
3	C	29	DC	O4'-C1'-N1	8.95	114.26	108.00
1	A	169	ARG	NE-CZ-NH1	-8.89	115.85	120.30
3	C	25	DT	O4'-C1'-N1	-8.75	101.87	108.00
3	C	26	DC	N1-C2-O2	8.61	124.07	118.90
1	A	46	ARG	NE-CZ-NH2	-8.59	116.01	120.30
3	C	25	DT	C4-C5-C6	8.42	123.05	118.00
1	A	144	PHE	CB-CG-CD2	-8.40	114.92	120.80
3	C	22	DA	N1-C2-N3	-8.00	125.30	129.30
2	B	8	DC	O4'-C1'-C2'	7.98	112.28	105.90
3	C	24	DG	P-O3'-C3'	7.73	128.98	119.70
2	B	9	DT	C5-C4-O4	-7.53	119.63	124.90
2	B	7	DC	N1-C2-O2	7.50	123.40	118.90
3	C	26	DC	N3-C4-N4	-7.25	112.93	118.00
2	B	9	DT	O5'-P-OP2	-7.22	99.20	105.70
2	B	9	DT	N3-C2-O2	7.20	126.62	122.30
3	C	21	DC	N1-C2-O2	7.12	123.17	118.90
3	C	22	DA	C6-N1-C2	7.06	122.84	118.60
2	B	10	DG	O4'-C1'-N9	6.90	112.83	108.00
3	C	26	DC	O4'-C1'-N1	-6.87	103.19	108.00
1	A	169	ARG	CG-CD-NE	-6.78	97.55	111.80
3	C	26	DC	C5-C4-N4	6.68	124.88	120.20
2	B	9	DT	N3-C4-O4	6.66	123.90	119.90
2	B	6	DA	C5-C6-N6	-6.64	118.39	123.70
3	C	25	DT	C6-C5-C7	-6.63	118.92	122.90
3	C	24	DG	N3-C4-N9	-6.53	122.08	126.00
3	C	24	DG	N3-C2-N2	-6.45	115.38	119.90
2	B	13	DC	O4'-C1'-N1	6.42	112.49	108.00
1	A	141	LEU	CB-CG-CD2	-6.38	100.16	111.00
2	B	13	DC	C3'-C2'-C1'	-6.30	94.94	102.50
2	B	12	DA	O4'-C1'-N9	-6.29	103.60	108.00
1	A	277	ARG	NE-CZ-NH2	6.19	123.40	120.30
2	B	8	DC	N3-C4-C5	6.18	124.37	121.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	83	SER	C-N-CA	6.10	136.94	121.70
1	A	93	LEU	CB-CG-CD1	6.03	121.25	111.00
3	C	21	DC	O4'-C1'-C2'	-5.94	101.15	105.90
3	C	17	DC	O4'-C1'-N1	5.92	112.15	108.00
1	A	228	LEU	CB-CG-CD2	5.88	120.99	111.00
1	A	79	ARG	NE-CZ-NH2	5.79	123.20	120.30
1	A	169	ARG	NE-CZ-NH2	5.79	123.20	120.30
2	B	6	DA	N3-C4-N9	5.77	132.01	127.40
2	B	5	DG	P-O3'-C3'	5.74	126.59	119.70
3	C	26	DC	P-O3'-C3'	5.59	126.41	119.70
2	B	5	DG	N1-C6-O6	-5.49	116.61	119.90
2	B	11	DG	P-O3'-C3'	5.46	126.25	119.70
1	A	213	ARG	NE-CZ-NH2	5.43	123.02	120.30
1	A	279	TYR	CB-CG-CD2	5.42	124.25	121.00
2	B	7	DC	C1'-O4'-C4'	-5.39	104.71	110.10
2	B	6	DA	N1-C2-N3	-5.37	126.61	129.30
3	C	19	DT	P-O3'-C3'	5.37	126.14	119.70
2	B	8	DC	C1'-O4'-C4'	-5.36	104.74	110.10
3	C	25	DT	C5-C6-N1	-5.32	120.51	123.70
3	C	24	DG	O4'-C1'-N9	-5.30	104.29	108.00
3	C	17	DC	P-O3'-C3'	5.28	126.03	119.70
1	A	184	LEU	CA-CB-CG	5.26	127.41	115.30
3	C	24	DG	C8-N9-C4	-5.24	104.31	106.40
3	C	21	DC	C1'-O4'-C4'	-5.21	104.89	110.10
3	C	25	DT	N3-C2-O2	-5.18	119.19	122.30
3	C	24	DG	N3-C4-C5	5.18	131.19	128.60
1	A	277	ARG	NE-CZ-NH1	-5.16	117.72	120.30
1	A	45	PHE	CB-CG-CD2	-5.14	117.20	120.80
2	B	5	DG	C5-C6-N1	5.14	114.07	111.50
1	A	46	ARG	NE-CZ-NH1	5.12	122.86	120.30
2	B	8	DC	C5-C6-N1	-5.10	118.45	121.00
2	B	6	DA	N3-C4-C5	-5.07	123.25	126.80
2	B	14	DG	P-O3'-C3'	5.05	125.76	119.70
3	C	27	DT	P-O3'-C3'	5.02	125.73	119.70

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	82	LYS	Peptide
1	A	83	SER	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2518	0	2456	20	0
2	B	289	0	157	3	0
3	C	286	0	158	4	0
4	A	1	0	0	0	0
4	C	1	0	0	0	0
5	A	346	0	0	6	0
5	B	28	0	0	0	0
5	C	25	0	0	1	0
All	All	3494	0	2771	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:322:ASP:O	5:A:2345:HOH:O	1.87	0.92
1:A:263:GLN:H	1:A:263:GLN:HE21	1.15	0.88
1:A:322:ASP:C	5:A:2345:HOH:O	2.14	0.86
3:C:22:DA:OP2	5:C:2010:HOH:O	1.94	0.85
1:A:111:HIS:HE1	5:A:2155:HOH:O	1.80	0.63
1:A:277:ARG:NH1	5:A:2302:HOH:O	2.31	0.63
2:B:14:DG:H2'	2:B:15:DC:C6	2.37	0.60
1:A:98:LYS:HE2	5:A:2049:HOH:O	2.01	0.59
1:A:46:ARG:HG3	1:A:139:GLU:OE1	2.07	0.55
1:A:287:GLN:CD	1:A:287:GLN:H	2.11	0.55
1:A:171:ILE:HD12	1:A:173:LEU:HG	1.90	0.52
2:B:2:DG:H1	3:C:29:DC:H42	1.59	0.49
1:A:185:GLN:H	1:A:185:GLN:NE2	2.11	0.49
1:A:132:LEU:HD21	1:A:256:LEU:HG	1.95	0.48
1:A:237:HIS:CE1	1:A:248:THR:HG23	2.51	0.46
1:A:46:ARG:CG	1:A:139:GLU:OE1	2.64	0.46
1:A:249:CYS:SG	1:A:253:LYS:NZ	2.63	0.45
2:B:3:DT:H2''	2:B:4:DA:OP2	2.16	0.45
1:A:141:LEU:HD21	1:A:215:ILE:HD11	1.97	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:46:ARG:NE	1:A:139:GLU:OE1	2.50	0.45
3:C:16:DG:H2'	3:C:17:DC:N1	2.32	0.44
1:A:263:GLN:N	1:A:263:GLN:HE21	1.97	0.44
3:C:16:DG:H2'	3:C:17:DC:C2	2.54	0.42
1:A:243:LEU:HA	1:A:244:PRO:HD3	1.83	0.42
1:A:223:ALA:O	1:A:227:GLN:CG	2.69	0.40
1:A:103:ASP:HB2	5:A:2057:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	315/360 (88%)	305 (97%)	8 (2%)	2 (1%)	25 7

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	84	GLN
1	A	81	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	267/302 (88%)	257 (96%)	10 (4%)	34 7

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	82	LYS
1	A	93	LEU
1	A	184	LEU
1	A	185	GLN
1	A	227	GLN
1	A	228	LEU
1	A	263	GLN
1	A	287	GLN
1	A	315	GLN
1	A	325	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	125	GLN
1	A	185	GLN
1	A	227	GLN
1	A	263	GLN
1	A	276	GLN
1	A	294	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	8OG	C	23	3	18,25,26	1.42	2 (11%)	21,37,40	2.22	5 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	8OG	C	23	3	-	0/3/21/22	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	23	8OG	C2-N2	4.06	1.42	1.33
3	C	23	8OG	C6-N1	3.02	1.38	1.33

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	23	8OG	C8-N9-C1'	6.10	136.40	125.90
3	C	23	8OG	C5-C6-N1	-5.59	115.78	123.43
3	C	23	8OG	C1'-N9-C4	-3.68	120.14	126.99
3	C	23	8OG	C6-N1-C2	2.36	119.68	115.93
3	C	23	8OG	C2-N3-C4	-2.31	112.72	115.36

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	2

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	80:GLY	C	81:ASP	N	1.95
1	A	83:SER	C	84:GLN	N	0.78

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	313/360 (86%)	0.39	19 (6%) 21 25	12, 21, 37, 55	0
2	B	14/15 (93%)	1.12	3 (21%) 0 0	26, 49, 83, 89	0
3	C	13/15 (86%)	1.76	5 (38%) 0 0	20, 47, 103, 109	0
All	All	340/390 (87%)	0.47	27 (7%) 12 15	12, 21, 47, 109	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	287	GLN	7.0
1	A	325	GLN	6.5
3	C	16	DG	5.9
1	A	70	GLU	4.2
3	C	17	DC	4.0
3	C	29	DC	3.8
1	A	285	THR	3.8
1	A	284	THR	3.7
1	A	286	SER	3.6
1	A	84	GLN	3.4
3	C	18	DG	3.4
2	B	2	DG	3.4
2	B	15	DC	3.3
1	A	17	ALA	3.3
1	A	324	ARG	3.0
2	B	14	DG	3.0
1	A	66	LEU	2.9
3	C	28	DA	2.8
1	A	201	LEU	2.7
1	A	145	ILE	2.5
1	A	288	ALA	2.4
1	A	18	SER	2.3
1	A	69	THR	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	64	TRP	2.2
1	A	142	PHE	2.1
1	A	59	LEU	2.1
1	A	250	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	8OG	C	23	23/24	0.98	0.05	10,13,16,17	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	CA	A	1327	1/1	0.88	0.07	58,58,58,58	0
4	CA	C	1030	1/1	0.99	0.09	48,48,48,48	0

6.5 Other polymers [i](#)

There are no such residues in this entry.