



Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 01:41 am BST

PDB ID : 2XHL
Title : Structure of a functional derivative of Clostridium botulinum neurotoxin type B
Authors : Masuyer, G.; Beard, M.; Cadd, V.A.; Chaddock, J.A.; Acharya, K.R.
Deposited on : 2010-06-18
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	453	<div> <div>2%</div> <div> <div></div> <div>85%</div> <div>9%</div> <div>5%</div> </div> </div>
2	B	433	<div> <div>2%</div> <div> <div></div> <div>88%</div> <div>6%</div> <div>5%</div> </div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6860 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BOTULINUM NEUROTOXIN B LIGHT CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	430	Total	C	N	O	S	0	0	0
			3499	2253	570	660	16			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	438	VAL	-	expression tag	UNP P10844
A	439	ASP	-	expression tag	UNP P10844
A	440	GLY	-	expression tag	UNP P10844
A	441	ILE	-	expression tag	UNP P10844
A	442	ILE	-	expression tag	UNP P10844
A	443	THR	-	expression tag	UNP P10844
A	444	SER	-	expression tag	UNP P10844
A	445	LYS	-	expression tag	UNP P10844
A	446	THR	-	expression tag	UNP P10844
A	447	LYS	-	expression tag	UNP P10844
A	448	SER	-	expression tag	UNP P10844
A	449	LEU	-	expression tag	UNP P10844
A	450	ILE	-	expression tag	UNP P10844
A	451	GLU	-	expression tag	UNP P10844
A	452	GLY	-	expression tag	UNP P10844
A	453	ARG	-	expression tag	UNP P10844

- Molecule 2 is a protein called BOTULINUM NEUROTOXIN B HEAVY CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	410	Total	C	N	O	S	0	0	0
			3329	2148	515	657	9			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	454	ASN	-	expression tag	UNP P10844
B	455	LYS	-	expression tag	UNP P10844
B	456	ALA	-	expression tag	UNP P10844
B	457	LEU	-	expression tag	UNP P10844
B	458	ASN	-	expression tag	UNP P10844
B	459	LEU	-	expression tag	UNP P10844
B	460	GLN	-	expression tag	UNP P10844
B	874	LEU	-	expression tag	UNP P10844
B	875	GLU	-	expression tag	UNP P10844
B	876	ALA	-	expression tag	UNP P10844
B	877	LEU	-	expression tag	UNP P10844
B	878	ALA	-	expression tag	UNP P10844
B	879	SER	-	expression tag	UNP P10844
B	880	GLY	-	expression tag	UNP P10844
B	881	HIS	-	expression tag	UNP P10844
B	882	HIS	-	expression tag	UNP P10844
B	883	HIS	-	expression tag	UNP P10844
B	884	HIS	-	expression tag	UNP P10844
B	885	HIS	-	expression tag	UNP P10844
B	886	HIS	-	expression tag	UNP P10844

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Zn 1 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	22	Total O 22 22	0	0
4	B	9	Total O 9 9	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: BOTULINUM NEUROTOXIN B LIGHT CHAIN



• Molecule 2: BOTULINUM NEUROTOXIN B HEAVY CHAIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	66.89Å 149.10Å 113.49Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	113.49 – 2.80 37.64 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.7 (113.49-2.80) 99.7 (37.64-2.80)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.98 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.240 , 0.282 0.231 , 0.245	Depositor DCC
R_{free} test set	1448 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	48.3	Xtriage
Anisotropy	0.455	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 36.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	6860	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.36	0/3577	0.47	0/4828
2	B	0.34	0/3390	0.46	0/4592
All	All	0.35	0/6967	0.46	0/9420

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3499	0	3471	29	0
2	B	3329	0	3294	23	0
3	A	1	0	0	0	0
4	A	22	0	0	1	0
4	B	9	0	0	0	0
All	All	6860	0	6765	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:718:GLN:O	2:B:722:THR:HB	1.73	0.87
1:A:247:LEU:HD12	1:A:248:PRO:HD2	1.59	0.84
1:A:368:LYS:H	1:A:412:GLN:HE22	1.25	0.82
2:B:862:THR:HG22	2:B:864:LEU:H	1.47	0.80
1:A:107:LYS:O	1:A:111:MET:HG2	1.85	0.76
2:B:640:GLY:CA	2:B:649:THR:HG22	2.18	0.72
1:A:1:MET:HB2	1:A:2:PRO:HD2	1.70	0.71
1:A:436:MET:HG2	2:B:545:ILE:HD11	1.74	0.70
2:B:640:GLY:HA3	2:B:649:THR:HG22	1.76	0.68
1:A:438:VAL:O	1:A:439:ASP:HB3	1.97	0.64
2:B:552:ILE:HG22	2:B:643:LEU:HB3	1.79	0.64
1:A:184:ARG:HG2	1:A:243:LYS:O	2.00	0.61
1:A:410:ARG:HG2	1:A:416:ILE:HD13	1.83	0.61
2:B:640:GLY:HA2	2:B:649:THR:HG22	1.83	0.59
2:B:640:GLY:HA2	2:B:649:THR:CG2	2.35	0.56
2:B:671:PRO:HG2	2:B:722:THR:CG2	2.36	0.55
1:A:8:PHE:CZ	1:A:37:PHE:HB3	2.42	0.54
4:A:2013:HOH:O	2:B:499:ASN:ND2	2.35	0.51
1:A:438:VAL:O	1:A:439:ASP:CB	2.59	0.50
1:A:222:SER:HB3	1:A:227:ILE:HD11	1.94	0.50
1:A:230:HIS:ND1	1:A:358:GLU:OE1	2.44	0.49
2:B:862:THR:HG22	2:B:863:ILE:N	2.27	0.49
1:A:102:LYS:HE3	1:A:365:TYR:HA	1.95	0.49
1:A:246:ASP:OD2	1:A:288:ILE:HG13	2.13	0.49
2:B:671:PRO:CG	2:B:722:THR:CG2	2.92	0.48
1:A:439:ASP:O	1:A:439:ASP:OD1	2.30	0.48
2:B:822:LEU:HD22	2:B:840:LYS:HG3	1.96	0.48
2:B:683:LEU:HD11	2:B:821:LEU:HD23	1.95	0.47
1:A:75:ASP:HB3	1:A:167:LEU:HD11	1.95	0.47
2:B:551:THR:OG1	2:B:552:ILE:N	2.47	0.47
1:A:16:ASN:HA	1:A:20:ILE:HG22	1.95	0.47
1:A:8:PHE:HE1	1:A:89:LEU:HD13	1.79	0.47
1:A:8:PHE:CE1	1:A:89:LEU:HD13	2.50	0.47
2:B:768:PHE:HA	2:B:771:ILE:HD12	1.96	0.46
2:B:833:ILE:HG22	2:B:834:GLY:H	1.81	0.45
2:B:673:LEU:HD11	2:B:802:MET:HG2	1.99	0.45
1:A:434:ILE:HG13	2:B:469:LEU:HD11	2.00	0.43
1:A:399:ASN:HB3	1:A:406:GLU:HA	2.00	0.42
1:A:256:PHE:HB2	2:B:545:ILE:HD12	2.01	0.42
1:A:196:CYS:HA	1:A:197:PRO:HD3	1.88	0.42
2:B:863:ILE:O	2:B:863:ILE:HG22	2.18	0.42
1:A:5:ILE:HD12	1:A:96:PHE:HB3	2.02	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:246:ASP:OD1	1:A:247:LEU:N	2.47	0.41
1:A:8:PHE:HE1	1:A:89:LEU:HD22	1.86	0.41
1:A:378:LEU:HB3	1:A:379:PRO:HD2	2.02	0.41
1:A:107:LYS:HA	1:A:110:GLU:HG2	2.02	0.41
2:B:646:GLY:HA2	2:B:668:GLU:HA	2.03	0.41
2:B:818:LYS:O	2:B:822:LEU:HG	2.20	0.41
1:A:147:GLU:HG2	1:A:148:VAL:H	1.84	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	426/453 (94%)	416 (98%)	10 (2%)	0	100	100
2	B	406/433 (94%)	394 (97%)	11 (3%)	1 (0%)	47	78
All	All	832/886 (94%)	810 (97%)	21 (2%)	1 (0%)	51	81

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	833	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	391/411 (95%)	391 (100%)	0	100	100
2	B	377/395 (95%)	376 (100%)	1 (0%)	92	98
All	All	768/806 (95%)	767 (100%)	1 (0%)	93	98

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	570	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	86	ASN
1	A	207	GLN
1	A	412	GLN
2	B	653	ASN
2	B	656	ASN
2	B	699	ASN
2	B	828	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	430/453 (94%)	0.04	11 (2%) 56 46	14, 32, 47, 52	0
2	B	410/433 (94%)	0.06	8 (1%) 65 56	13, 40, 67, 69	0
All	All	840/886 (94%)	0.05	19 (2%) 60 51	13, 35, 62, 69	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	861	ASP	6.9
1	A	334	SER	4.7
1	A	440	GLY	4.5
1	A	145	PRO	3.4
2	B	769	ASN	3.2
1	A	207	GLN	3.0
2	B	606	ALA	2.9
1	A	404	ASP	2.9
1	A	149	GLU	2.9
1	A	402	ASP	2.8
2	B	458	ASN	2.8
2	B	688	ASP	2.8
1	A	395	GLU	2.4
1	A	427	GLU	2.4
2	B	566	ARG	2.3
1	A	146	GLY	2.3
1	A	335	GLU	2.2
2	B	563	LEU	2.1
2	B	764	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	ZN	A	1442	1/1	0.96	0.06	33,33,33,33	0

6.5 Other polymers [i](#)

There are no such residues in this entry.