



# Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2020 – 03:12 pm BST

PDB ID : 6XIA  
Title : REFINEMENT OF GLUCOSE ISOMERASE FROM STREPTOMYCES ALBUS AT 1.65 ANGSTROMS WITH DATA FROM AN IMAGING PLATE  
Authors : Dauter, Z.; Terry, H.; Wilson, K.S.  
Deposited on : 1990-09-13  
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1931 (1.66-1.66)
Ramachandran outliers	138981	1891 (1.66-1.66)
Sidechain outliers	138945	1891 (1.66-1.66)
RSRZ outliers	127900	1791 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	387	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%; text-align: center;">%</div> <div style="position: absolute; top: 10px; left: 0; width: 100%; text-align: center;">79%18%..</div> </div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3499 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called D-XYLOSE ISOMERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	387	Total	C	N	O	S	25	0	0
			3045	1915	546	576	8			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	454	Total	O	0	0
			454	454		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: D-XYLOSE ISOMERASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	93.90 Å 99.70 Å 102.90 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 1.65 19.71 – 1.66	Depositor EDS
% Data completeness (in resolution range)	(Not available) (8.00-1.65) 94.9 (19.71-1.66)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	8.45 (at 1.66 Å)	Xtriage
Refinement program	PROLSQ	Depositor
R, $R_{free}$	0.141 , (Not available) 0.136 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	13.5	Xtriage
Anisotropy	0.230	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 78.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.013 for -h,-l,-k	Xtriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	3499	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.59% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.06	22/3118 (0.7%)	1.67	59/4221 (1.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	206	GLU	CD-OE2	7.17	1.33	1.25
1	A	69	GLU	CD-OE2	6.89	1.33	1.25
1	A	44	GLU	CD-OE2	6.88	1.33	1.25
1	A	37	GLU	CD-OE2	6.87	1.33	1.25
1	A	327	GLU	CD-OE2	6.85	1.33	1.25
1	A	324	GLU	CD-OE2	6.24	1.32	1.25
1	A	293	GLU	CD-OE2	6.18	1.32	1.25
1	A	180	GLU	CD-OE2	6.07	1.32	1.25
1	A	143	GLU	CD-OE2	6.05	1.32	1.25
1	A	209	GLU	CD-OE2	6.03	1.32	1.25
1	A	66	GLU	CD-OE2	5.83	1.32	1.25
1	A	159	GLU	CD-OE2	5.76	1.31	1.25
1	A	203	GLU	CD-OE2	5.70	1.31	1.25
1	A	166	GLU	CD-OE2	5.65	1.31	1.25
1	A	275	GLU	CD-OE2	5.62	1.31	1.25
1	A	127	GLU	CD-OE2	5.58	1.31	1.25
1	A	140	GLU	CD-OE2	5.54	1.31	1.25
1	A	357	GLU	CD-OE2	5.44	1.31	1.25
1	A	372	GLU	CD-OE2	5.43	1.31	1.25
1	A	358	GLU	CD-OE2	5.10	1.31	1.25

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	336	GLU	CD-OE2	5.08	1.31	1.25
1	A	314	GLU	CD-OE2	5.01	1.31	1.25

All (59) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	204	ARG	NE-CZ-NH2	-19.59	110.50	120.30
1	A	204	ARG	NE-CZ-NH1	17.07	128.84	120.30
1	A	339	ARG	NE-CZ-NH2	-16.94	111.83	120.30
1	A	265	ARG	NE-CZ-NH2	-15.82	112.39	120.30
1	A	73	ARG	NE-CZ-NH2	-14.80	112.90	120.30
1	A	330	ARG	NE-CZ-NH2	-12.44	114.08	120.30
1	A	67	ARG	NE-CZ-NH1	-12.04	114.28	120.30
1	A	367	ARG	NE-CZ-NH1	11.43	126.02	120.30
1	A	207	ARG	NE-CZ-NH2	-11.25	114.67	120.30
1	A	40	ARG	NE-CZ-NH1	10.60	125.60	120.30
1	A	258	ARG	NE-CZ-NH1	10.18	125.39	120.30
1	A	75	ARG	NE-CZ-NH2	-10.12	115.24	120.30
1	A	73	ARG	NE-CZ-NH1	10.03	125.31	120.30
1	A	386	ARG	NE-CZ-NH1	9.87	125.23	120.30
1	A	187	ARG	NE-CZ-NH1	9.74	125.17	120.30
1	A	367	ARG	CD-NE-CZ	9.45	136.83	123.60
1	A	380	ASP	CB-CG-OD1	9.06	126.45	118.30
1	A	339	ARG	NH1-CZ-NH2	8.73	129.00	119.40
1	A	244	ASP	CB-CG-OD1	8.56	126.00	118.30
1	A	9	ARG	NE-CZ-NH2	8.05	124.33	120.30
1	A	151	ARG	NE-CZ-NH2	-8.01	116.30	120.30
1	A	149	ASP	CB-CG-OD1	7.88	125.39	118.30
1	A	23	ASP	CB-CG-OD2	-7.87	111.21	118.30
1	A	307	ARG	NE-CZ-NH2	-7.85	116.38	120.30
1	A	34	ASP	CB-CG-OD1	7.75	125.27	118.30
1	A	339	ARG	CD-NE-CZ	-7.59	112.97	123.60
1	A	41	ARG	CD-NE-CZ	7.46	134.04	123.60
1	A	211	TYR	CB-CG-CD2	-7.14	116.72	121.00
1	A	272	ASP	CB-CG-OD1	6.99	124.59	118.30
1	A	22	ARG	NE-CZ-NH1	6.94	123.77	120.30
1	A	149	ASP	CB-CG-OD2	-6.94	112.06	118.30
1	A	100	ASP	CB-CG-OD2	-6.86	112.12	118.30
1	A	34	ASP	CB-CG-OD2	-6.77	112.21	118.30
1	A	68	TYR	CB-CA-C	-6.61	97.17	110.40
1	A	116	ARG	NE-CZ-NH2	6.61	123.60	120.30
1	A	322	ASP	CB-CG-OD1	6.52	124.17	118.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	204	ARG	CD-NE-CZ	6.42	132.59	123.60
1	A	1	ASN	CB-CA-C	6.37	123.14	110.40
1	A	367	ARG	NE-CZ-NH2	-6.31	117.15	120.30
1	A	283	ARG	NE-CZ-NH2	-6.30	117.15	120.30
1	A	330	ARG	NE-CZ-NH1	6.29	123.45	120.30
1	A	67	ARG	NE-CZ-NH2	6.25	123.42	120.30
1	A	283	ARG	NE-CZ-NH1	6.02	123.31	120.30
1	A	163	LEU	CA-CB-CG	5.92	128.92	115.30
1	A	265	ARG	NE-CZ-NH1	5.92	123.26	120.30
1	A	22	ARG	CG-CD-NE	-5.83	99.55	111.80
1	A	375	ASP	CB-CG-OD1	5.67	123.41	118.30
1	A	336	GLU	CG-CD-OE1	5.62	129.53	118.30
1	A	314	GLU	CG-CD-OE1	5.60	129.50	118.30
1	A	327	GLU	CG-CD-OE2	-5.44	107.41	118.30
1	A	258	ARG	NE-CZ-NH2	-5.36	117.62	120.30
1	A	320	ARG	NE-CZ-NH1	5.32	122.96	120.30
1	A	75	ARG	NE-CZ-NH1	5.28	122.94	120.30
1	A	344	ASP	CB-CG-OD1	5.24	123.02	118.30
1	A	23	ASP	CB-CA-C	5.23	120.86	110.40
1	A	112	ARG	NE-CZ-NH1	-5.18	117.71	120.30
1	A	56	ASP	CB-CG-OD1	5.14	122.93	118.30
1	A	67	ARG	CD-NE-CZ	-5.14	116.41	123.60
1	A	336	GLU	CG-CD-OE2	-5.06	108.18	118.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	204	ARG	Sidechain
1	A	207	ARG	Sidechain
1	A	75	ARG	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3045	0	2912	34	10

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	454	0	0	12	3
All	All	3499	0	2912	34	11

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:385:ALA:O	1:A:386:ARG:HB2	1.56	1.03
1:A:95:HIS:HD2	1:A:97:VAL:H	1.07	1.02
1:A:386:ARG:HG3	2:A:762:HOH:O	1.68	0.91
1:A:265:ARG:HD2	2:A:535:HOH:O	1.78	0.83
1:A:174:ASP:HB2	2:A:718:HOH:O	1.81	0.79
1:A:91:ASN:HD21	1:A:94:THR:H	1.29	0.78
1:A:95:HIS:CD2	1:A:97:VAL:H	1.99	0.75
1:A:91:ASN:HD22	1:A:91:ASN:C	1.95	0.69
1:A:265:ARG:CD	2:A:535:HOH:O	2.40	0.68
1:A:65:SER:O	1:A:68:TYR:HB2	1.94	0.67
1:A:216:GLU:OE1	2:A:794:HOH:O	2.15	0.64
1:A:23:ASP:HB2	1:A:24:PRO:HD2	1.80	0.64
1:A:385:ALA:O	1:A:386:ARG:CB	2.42	0.62
1:A:91:ASN:ND2	1:A:93:PHE:H	2.02	0.58
1:A:41:ARG:HA	1:A:41:ARG:HE	1.70	0.56
1:A:23:ASP:HB2	1:A:24:PRO:CD	2.36	0.56
1:A:381:HIS:HD2	2:A:756:HOH:O	1.88	0.55
1:A:68:TYR:HD2	2:A:504:HOH:O	1.90	0.54
1:A:151:ARG:NH1	2:A:697:HOH:O	2.30	0.54
1:A:322:ASP:OD2	1:A:381:HIS:HE1	1.91	0.53
1:A:91:ASN:ND2	1:A:94:THR:H	2.05	0.51
1:A:75:ARG:HD2	2:A:629:HOH:O	2.10	0.49
1:A:185:GLU:OE1	1:A:254:ASP:HB3	2.14	0.47
1:A:91:ASN:HD22	1:A:93:PHE:H	1.61	0.47
1:A:95:HIS:CD2	1:A:96:PRO:HD2	2.50	0.46
1:A:131:GLU:CG	2:A:727:HOH:O	2.65	0.44
1:A:75:ARG:CD	2:A:629:HOH:O	2.65	0.44
1:A:226:ASN:OD1	1:A:228:PRO:HD2	2.18	0.44
1:A:91:ASN:ND2	1:A:91:ASN:C	2.68	0.43
1:A:131:GLU:CB	2:A:727:HOH:O	2.67	0.43
1:A:53:HIS:CD2	1:A:89:THR:HG23	2.56	0.41
1:A:381:HIS:CE1	1:A:387:GLY:HA3	2.55	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:23:ASP:CB	1:A:24:PRO:CD	2.97	0.41
1:A:23:ASP:CB	1:A:24:PRO:HD2	2.49	0.41

All (11) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1:ASN:OD1	1:A:386:ARG:NH2[2_555]	0.86	1.34
1:A:1:ASN:OD1	1:A:386:ARG:CZ[2_555]	1.09	1.11
1:A:1:ASN:CG	1:A:386:ARG:NH2[2_555]	1.14	1.06
1:A:1:ASN:N	1:A:386:ARG:NH1[2_555]	1.51	0.69
1:A:1:ASN:OD1	1:A:386:ARG:NE[2_555]	1.56	0.64
1:A:386:ARG:CD	2:A:818:HOH:O[2_555]	1.73	0.47
1:A:1:ASN:CB	1:A:386:ARG:NH2[2_555]	1.88	0.32
1:A:1:ASN:N	1:A:386:ARG:CZ[2_555]	2.07	0.13
1:A:1:ASN:CG	1:A:386:ARG:CZ[2_555]	2.10	0.10
1:A:27:ASP:OD1	2:A:524:HOH:O[3_555]	2.12	0.08
2:A:492:HOH:O	2:A:793:HOH:O[3_555]	2.12	0.08

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	385/387 (100%)	372 (97%)	11 (3%)	2 (0%)	29 11

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	386	ARG
1	A	185	GLU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	303/303 (100%)	296 (98%)	7 (2%)	50 25

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	41	ARG
1	A	64	ASP
1	A	90	THR
1	A	91	ASN
1	A	131	GLU
1	A	163	LEU
1	A	184	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	91	ASN
1	A	95	HIS
1	A	184	ASN
1	A	221	GLN
1	A	308	ASN
1	A	381	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	387/387 (100%)	-0.70	3 (0%) 86 88	7, 13, 34, 70	7 (1%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	68	TYR	5.8
1	A	1	ASN	5.4
1	A	386	ARG	3.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.