



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 8, 2020 – 05:34 AM BST

PDB ID : 5XLD  
Title : The structure of hemagglutinin Q226L-G228S mutant from an avian-origin H4N6 influenza virus in complex with human receptor analog LSTc  
Authors : Song, H.; Qi, J.; Gao, F.G.  
Deposited on : 2017-05-10  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	327	<div> <div>6%</div> <div> <div></div> <div>85%</div> <div>10%</div> <div>..</div> </div> </div>
1	B	327	<div> <div>15%</div> <div> <div></div> <div>80%</div> <div>15%</div> <div>..</div> </div> </div>
2	C	176	<div> <div>2%</div> <div> <div></div> <div>91%</div> <div>6%</div> <div>..</div> </div> </div>
2	D	176	<div> <div>%</div> <div> <div></div> <div>91%</div> <div>5%</div> <div>..</div> </div> </div>
3	E	3	<div> <div></div> <div>100%</div> </div>
3	F	3	<div> <div>33%</div> <div>67%</div> </div>

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 8217 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hemagglutinin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	319	Total	C	N	O	S	0	0	0
			2460	1542	436	470	12			
1	B	319	Total	C	N	O	S	0	0	0
			2460	1542	436	470	12			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	223	LEU	GLN	engineered mutation	UNP A3KF09
A	225	SER	GLY	engineered mutation	UNP A3KF09
B	223	LEU	GLN	engineered mutation	UNP A3KF09
B	225	SER	GLY	engineered mutation	UNP A3KF09

- Molecule 2 is a protein called Hemagglutinin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	172	Total	C	N	O	S	0	0	0
			1404	871	250	279	4			
2	D	172	Total	C	N	O	S	0	0	0
			1404	871	250	279	4			

- Molecule 3 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-6)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	E	3	Total	C	N	O	0	0	0
			46	25	2	19			

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	F	3	Total	C	N	O	0	0	0
			46	25	2	19			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	97	Total	O	0	0
			97	97		
5	C	100	Total	O	0	0
			100	100		
5	B	63	Total	O	0	0
			63	63		
5	D	81	Total	O	0	0
			81	81		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Hemagglutinin



#### • Molecule 1: Hemagglutinin




#### • Molecule 2: Hemagglutinin



#### • Molecule 2: Hemagglutinin





● Molecule 3: N-acetyl-alpha-neuraminic acid-(2-6)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



● Molecule 3: N-acetyl-alpha-neuraminic acid-(2-6)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	100.41Å 100.41Å 686.91Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	42.73 – 2.30 42.73 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.7 (42.73-2.30) 90.7 (42.73-2.30)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.53 (at 2.29Å)	Xtrriage
Refinement program	PHENIX 1.7.3 _928	Depositor
R, $R_{free}$	0.214 , 0.251 0.213 , 0.247	Depositor DCC
$R_{free}$ test set	3043 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	31.8	Xtrriage
Anisotropy	0.587	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 45.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	8217	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	50.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 50.88 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.0777e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SIA, GAL, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.34	0/2514	0.51	0/3423
1	B	0.32	0/2514	0.50	0/3423
2	C	0.39	0/1428	0.52	0/1922
2	D	0.38	0/1428	0.49	0/1922
All	All	0.35	0/7884	0.51	0/10690

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2460	0	2413	24	0
1	B	2460	0	2413	41	0
2	C	1404	0	1320	8	0
2	D	1404	0	1320	10	0
3	E	46	0	39	0	0
3	F	46	0	40	0	0
4	A	28	0	26	0	0
4	B	28	0	26	3	0
5	A	97	0	0	4	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	63	0	0	7	0
5	C	100	0	0	5	0
5	D	81	0	0	5	0
All	All	8217	0	7597	81	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (81) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:124:ASN:HB2	1:A:161:GLN:HE22	1.39	0.87
2:D:346:ASP:OD1	5:D:701:HOH:O	1.92	0.86
1:A:28:ASP:O	5:A:701:HOH:O	1.93	0.86
1:B:5:GLY:HA2	2:D:468:PHE:HA	1.62	0.82
1:B:152:LYS:NZ	1:B:189:THR:O	2.13	0.82
1:A:124:ASN:CB	1:A:161:GLN:HE22	1.94	0.80
1:B:213:ASN:O	1:B:217:ARG:NH2	2.15	0.79
1:B:244:THR:HG22	1:B:246:GLY:H	1.49	0.75
1:B:187:GLU:O	5:B:701:HOH:O	2.05	0.74
1:A:244:THR:HG22	1:A:246:GLY:H	1.55	0.71
1:B:5:GLY:N	5:B:707:HOH:O	2.25	0.69
2:C:346:ASP:OD2	5:C:601:HOH:O	2.09	0.68
1:A:134:ALA:O	1:A:221:ARG:NH1	2.28	0.67
2:C:369:GLN:OE1	5:C:602:HOH:O	2.11	0.67
1:B:102:GLN:OE1	5:B:703:HOH:O	2.12	0.67
1:B:167:ASN:OD1	5:B:702:HOH:O	2.12	0.67
1:A:302:ASP:OD1	5:A:702:HOH:O	2.14	0.66
2:D:413:ASP:OD1	5:D:702:HOH:O	2.14	0.65
2:D:450:ARG:NH2	5:D:703:HOH:O	2.29	0.64
1:A:166:ILE:HA	1:A:239:LEU:HB3	1.83	0.61
2:D:382:LEU:HD11	2:D:426:LEU:HD21	1.83	0.60
1:B:185:ASP:OD2	1:B:185:ASP:N	2.33	0.60
1:A:124:ASN:HB2	1:A:161:GLN:NE2	2.13	0.59
1:B:319:ARG:NE	5:B:704:HOH:O	2.14	0.57
1:B:199:VAL:HG22	1:B:244:THR:HG23	1.86	0.57
1:A:183:SER:N	1:A:187:GLU:OE2	2.27	0.56
1:A:152:LYS:NZ	1:A:189:THR:O	2.31	0.56
2:C:451:ARG:HD2	5:C:606:HOH:O	2.06	0.54
1:B:204:LYS:HE2	1:B:238:ASP:HA	1.88	0.54
2:C:378:LYS:NZ	2:C:430:GLU:OE1	2.37	0.54

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:137:ARG:NE	1:B:142:ASP:OD2	2.33	0.54
2:C:450:ARG:NH2	5:C:605:HOH:O	2.42	0.52
1:B:43:ASN:HB3	1:B:295:ILE:HD13	1.90	0.52
1:B:86:ARG:O	5:B:705:HOH:O	2.19	0.51
1:B:182:PRO:HB3	1:B:187:GLU:HB3	1.92	0.51
1:B:181:HIS:HE2	1:B:228:SER:HB2	1.76	0.51
1:B:123:TRP:CH2	1:B:163:LEU:HD11	2.46	0.51
1:B:164:THR:CB	4:B:601:NAG:H81	2.41	0.51
2:C:357:GLU:OE2	2:C:472:ASP:HB2	2.10	0.51
1:A:185:ASP:OD2	1:A:185:ASP:N	2.42	0.50
1:B:124:ASN:O	1:B:126:VAL:N	2.41	0.50
2:D:451:ARG:HD2	5:D:704:HOH:O	2.11	0.50
1:B:289:THR:O	2:D:383:ILE:HG23	2.12	0.50
1:B:63:ILE:HG13	1:B:101:TYR:CE2	2.47	0.49
1:B:184:THR:HG23	1:B:187:GLU:HG3	1.94	0.49
1:B:100:GLU:HB2	1:B:231:TRP:HH2	1.78	0.49
1:B:164:THR:HB	4:B:601:NAG:H81	1.96	0.48
1:B:162:ASN:O	4:B:601:NAG:O7	2.31	0.48
2:D:383:ILE:O	2:D:384:GLU:HB2	2.12	0.48
1:B:100:GLU:HB2	1:B:231:TRP:CH2	2.49	0.48
1:A:173:ARG:NE	5:A:703:HOH:O	2.19	0.48
1:A:110:ASN:HA	1:A:263:SER:O	2.14	0.48
1:B:203:THR:OG1	1:B:206:SER:OG	2.25	0.48
1:A:123:TRP:CH2	1:A:163:LEU:HD11	2.48	0.48
1:B:110:ASN:HA	1:B:263:SER:O	2.15	0.47
2:D:328:GLY:N	5:D:709:HOH:O	2.48	0.46
1:B:125:THR:HG23	1:B:159:PRO:HG3	1.97	0.46
2:C:328:GLY:N	5:C:609:HOH:O	2.49	0.46
1:A:199:VAL:HG22	1:A:244:THR:HG23	1.98	0.46
1:B:175:TYR:HA	5:B:714:HOH:O	2.15	0.46
1:A:6:ASN:HB2	5:A:725:HOH:O	2.16	0.45
2:C:331:GLY:O	2:C:335:GLY:HA3	2.16	0.45
1:B:124:ASN:OD1	1:B:125:THR:HG22	2.17	0.45
1:B:93:CYS:N	1:B:221:ARG:HH12	2.15	0.44
1:A:162:ASN:HA	1:A:242:PHE:O	2.18	0.44
1:B:203:THR:HB	1:B:238:ASP:OD1	2.17	0.43
1:A:144:PHE:HB2	1:A:147:LEU:HB2	2.00	0.43
1:B:166:ILE:HA	1:B:239:LEU:HB3	2.00	0.43
1:A:6:ASN:H	1:A:7:PRO:CD	2.31	0.43
1:B:104:LEU:HB2	1:B:231:TRP:CZ3	2.54	0.42
1:A:152:LYS:HE2	1:A:193:LYS:HB2	2.01	0.42

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:43:ASN:HB3	1:A:295:ILE:HD13	2.00	0.42
1:B:219:LEU:HA	1:B:219:LEU:HD13	1.91	0.41
1:B:175:TYR:CE2	1:B:254:HIS:HB3	2.55	0.41
1:A:53:ARG:HB2	1:A:53:ARG:HE	1.53	0.41
1:A:177:TRP:CE2	1:A:201:VAL:HG21	2.55	0.41
1:B:54:LEU:HD22	1:B:82:VAL:HB	2.01	0.41
1:A:63:ILE:HG13	1:A:101:TYR:CE2	2.56	0.41
1:B:148:ASN:HB2	1:B:252:ARG:HD2	2.02	0.41
1:B:189:THR:HG22	1:B:193:LYS:O	2.20	0.40
2:D:379:LEU:HA	2:D:379:LEU:HD23	1.93	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	317/327 (97%)	301 (95%)	14 (4%)	2 (1%)	25	31
1	B	317/327 (97%)	297 (94%)	17 (5%)	3 (1%)	17	20
2	C	170/176 (97%)	160 (94%)	10 (6%)	0	100	100
2	D	170/176 (97%)	157 (92%)	11 (6%)	2 (1%)	13	14
All	All	974/1006 (97%)	915 (94%)	52 (5%)	7 (1%)	22	26

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	6	ASN
1	A	124	ASN
1	B	125	THR
2	D	384	GLU
1	B	6	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	156	ASN
2	D	383	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	276/283 (98%)	265 (96%)	11 (4%)	31	44
1	B	276/283 (98%)	266 (96%)	10 (4%)	35	49
2	C	147/150 (98%)	145 (99%)	2 (1%)	67	81
2	D	147/150 (98%)	145 (99%)	2 (1%)	67	81
All	All	846/866 (98%)	821 (97%)	25 (3%)	41	57

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	HIS
1	A	28	ASP
1	A	124	ASN
1	A	127	LYS
1	A	163	LEU
1	A	185	ASP
1	A	205	THR
1	A	219	LEU
1	A	225	SER
1	A	239	LEU
1	A	244	THR
2	C	386	THR
2	C	450	ARG
1	B	14	HIS
1	B	125	THR
1	B	163	LEU
1	B	185	ASP
1	B	187	GLU
1	B	205	THR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	219	LEU
1	B	239	LEU
1	B	244	THR
1	B	252	ARG
2	D	386	THR
2	D	450	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	124	ASN
1	A	161	GLN
1	B	14	HIS
2	D	499	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	NAG	E	1	3	15,15,15	0.52	0	21,21,21	1.54	3 (14%)
3	GAL	E	2	3	11,11,12	1.59	1 (9%)	15,15,17	1.70	4 (26%)
3	SIA	E	3	3	17,20,21	0.52	0	21,28,31	3.11	5 (23%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NAG	F	1	3	15,15,15	0.48	0	21,21,21	1.43	3 (14%)
3	GAL	F	2	3	11,11,12	0.26	0	15,15,17	0.63	0
3	SIA	F	3	3	17,20,21	0.62	0	21,28,31	3.08	4 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	E	1	3	-	2/6/26/26	0/1/1/1
3	GAL	E	2	3	-	2/2/19/22	0/1/1/1
3	SIA	E	3	3	-	2/14/34/38	0/1/1/1
3	NAG	F	1	3	-	0/6/26/26	0/1/1/1
3	GAL	F	2	3	-	0/2/19/22	0/1/1/1
3	SIA	F	3	3	-	2/14/34/38	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	2	GAL	O4-C4	-4.78	1.31	1.43

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	3	SIA	O6-C2-C3	-10.20	91.99	109.87
3	E	3	SIA	O6-C2-C3	-10.19	92.00	109.87
3	E	3	SIA	C3-C2-C1	-8.12	94.21	111.93
3	F	3	SIA	C3-C2-C1	-7.93	94.63	111.93
3	E	1	NAG	C4-C3-C2	4.24	116.55	110.34
3	F	1	NAG	C4-C3-C2	4.20	116.49	110.34
3	F	3	SIA	C4-C3-C2	4.04	117.05	109.81
3	E	3	SIA	C4-C3-C2	3.97	116.93	109.81
3	E	2	GAL	O5-C5-C6	3.56	112.78	107.20
3	E	1	NAG	C1-C2-C3	3.52	115.35	110.54
3	E	2	GAL	C1-C2-C3	3.03	113.39	109.67
3	F	1	NAG	C1-C2-C3	2.91	114.51	110.54
3	E	2	GAL	C3-C4-C5	-2.88	105.11	110.24
3	E	2	GAL	C1-O5-C5	-2.76	108.45	112.19
3	E	3	SIA	C6-O6-C2	2.60	116.89	111.34
3	F	3	SIA	C6-O6-C2	2.31	116.29	111.34

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	1	NAG	C3-C4-C5	2.27	114.28	110.24
3	E	1	NAG	C1-C2-N2	-2.19	108.19	110.73
3	E	3	SIA	C4-C5-C6	-2.09	103.80	109.10

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	E	1	NAG	C8-C7-N2-C2
3	E	1	NAG	O7-C7-N2-C2
3	F	3	SIA	C11-C10-N5-C5
3	F	3	SIA	O10-C10-N5-C5
3	E	2	GAL	O5-C5-C6-O6
3	E	2	GAL	C4-C5-C6-O6
3	E	3	SIA	C11-C10-N5-C5
3	E	3	SIA	O10-C10-N5-C5

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

## Oligosaccharide Chain E



## Oligosaccharide Chain F





## 5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
4	NAG	B	602	1	14,14,15	0.45	0	17,19,21	1.08	1 (5%)
4	NAG	A	601	1	14,14,15	0.51	0	17,19,21	1.06	1 (5%)
4	NAG	A	602	1	14,14,15	0.45	0	17,19,21	1.02	1 (5%)
4	NAG	B	601	1	14,14,15	0.30	0	17,19,21	0.61	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	B	602	1	-	0/6/23/26	0/1/1/1
4	NAG	A	601	1	-	1/6/23/26	0/1/1/1
4	NAG	A	602	1	-	0/6/23/26	0/1/1/1
4	NAG	B	601	1	-	4/6/23/26	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	602	NAG	C1-O5-C5	3.60	117.06	112.19
4	A	602	NAG	C1-O5-C5	3.11	116.40	112.19
4	A	601	NAG	O5-C1-C2	2.69	115.54	111.29

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	601	NAG	C8-C7-N2-C2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
4	B	601	NAG	O7-C7-N2-C2
4	B	601	NAG	C4-C5-C6-O6
4	B	601	NAG	O5-C5-C6-O6
4	A	601	NAG	O5-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	601	NAG	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	319/327 (97%)	0.14	18 (5%) 24 30	21, 45, 82, 120	0
1	B	319/327 (97%)	0.70	49 (15%) 2 3	24, 64, 115, 139	0
2	C	172/176 (97%)	0.07	3 (1%) 70 76	19, 33, 52, 114	0
2	D	172/176 (97%)	0.04	2 (1%) 79 83	24, 35, 62, 123	0
All	All	982/1006 (97%)	0.29	72 (7%) 15 20	19, 43, 97, 139	0

All (72) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	192	TYR	7.2
1	B	191	LEU	6.0
1	B	158	TYR	5.8
1	B	219	LEU	5.2
1	B	197	GLY	5.0
1	B	184	THR	4.9
1	B	99	PRO	4.6
1	B	123	TRP	4.5
1	B	157	ALA	4.4
1	B	193	LYS	4.3
2	C	383	ILE	4.2
1	B	220	VAL	4.2
2	D	383	ILE	4.2
1	B	183	SER	4.2
1	B	151	VAL	4.0
1	B	153	SER	3.9
1	B	160	LEU	3.9
1	A	151	VAL	3.8
1	B	94	TYR	3.8
1	B	187	GLU	3.6
1	A	157	ALA	3.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	152	LYS	3.5
1	B	134	ALA	3.5
1	B	225	SER	3.4
1	B	222	GLY	3.4
1	A	191	LEU	3.3
1	B	188	GLN	3.3
1	B	155	GLY	3.2
1	A	155	GLY	3.2
1	B	223	LEU	3.2
1	A	219	LEU	3.1
1	B	207	GLN	2.9
1	B	182	PRO	2.9
1	B	224	SER	2.9
1	B	147	LEU	2.9
1	A	190	ASN	2.9
1	A	222	GLY	2.8
1	B	125	THR	2.8
1	B	181	HIS	2.8
1	B	156	ASN	2.6
1	B	221	ARG	2.5
1	A	216	SER	2.5
1	B	216	SER	2.5
1	A	158	TYR	2.4
1	B	126	VAL	2.4
2	C	386	THR	2.4
1	A	127	LYS	2.4
1	B	186	THR	2.4
1	B	127	LYS	2.4
1	B	150	LEU	2.4
1	A	5	GLY	2.3
1	B	190	ASN	2.3
1	A	184	THR	2.3
1	A	124	ASN	2.2
1	B	211	VAL	2.2
2	D	384	GLU	2.2
1	A	186	THR	2.2
1	A	123	TRP	2.2
1	B	161	GLN	2.2
1	B	180	HIS	2.1
1	B	244	THR	2.1
1	A	187	GLU	2.1
1	A	152	LYS	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	192	TYR	2.1
1	B	189	THR	2.1
1	B	139	ASN	2.1
1	B	249	ILE	2.1
1	B	218	PRO	2.1
2	C	384	GLU	2.1
1	B	208	THR	2.0
1	B	209	SER	2.0
1	B	166	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	NAG	F	1	15/15	0.47	0.39	149,164,169,171	0
3	NAG	E	1	15/15	0.62	0.33	126,135,142,144	0
3	GAL	E	2	11/12	0.76	0.30	98,112,122,123	0
3	SIA	F	3	20/21	0.78	0.28	82,88,93,95	0
3	GAL	F	2	11/12	0.89	0.48	116,134,143,147	0
3	SIA	E	3	20/21	0.90	0.19	63,69,76,76	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around Chain E:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	NAG	A	601	14/15	0.78	0.18	90,96,100,101	0
4	NAG	B	601	14/15	0.83	0.31	101,106,113,113	0
4	NAG	A	602	14/15	0.94	0.13	29,36,42,44	0
4	NAG	B	602	14/15	0.94	0.12	39,43,48,49	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.