



Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 03:35 pm BST

PDB ID : 1XNH
Title : Crystal Structure of NH₃-dependent NAD⁺ synthetase from *Helicobacter pylori*
Authors : Kang, G.B.; Kim, Y.S.; Im, Y.J.; Rho, S.H.; Lee, J.H.; Eom, S.H.
Deposited on : 2004-10-05
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	268	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2001 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NH(3)-dependent NAD(+) synthetase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	251	Total	C	N	O	S	0	0	0
			1964	1270	318	368	8			


There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	261	LEU	-	CLONING ARTIFACT	UNP O25096
A	262	GLU	-	CLONING ARTIFACT	UNP O25096
A	263	HIS	-	CLONING ARTIFACT	UNP O25096
A	264	HIS	-	CLONING ARTIFACT	UNP O25096
A	265	HIS	-	CLONING ARTIFACT	UNP O25096
A	266	HIS	-	CLONING ARTIFACT	UNP O25096
A	267	HIS	-	CLONING ARTIFACT	UNP O25096
A	268	HIS	-	CLONING ARTIFACT	UNP O25096

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	37	Total	O	0	0
			37	37		

- Molecule 1: NH(3)-dependent NAD(+) synthetase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	93.83Å 48.32Å 64.20Å 90.00° 109.97° 90.00°	Depositor
Resolution (Å)	26.74 – 2.30 26.74 – 2.30	Depositor EDS
% Data completeness (in resolution range)	(Not available) (26.74-2.30) 99.9 (26.74-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.68 (at 2.31Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.255 , 0.295 0.271 , 0.278	Depositor DCC
R_{free} test set	1277 reflections (10.25%)	wwPDB-VP
Wilson B-factor (Å ²)	43.7	Xtriage
Anisotropy	0.181	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 38.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	2001	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.47	0/2003	0.60	0/2708

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1964	0	1992	71	0
2	A	37	0	0	2	0
All	All	2001	0	1992	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:169:ARG:HB3	1:A:169:ARG:HH11	1.29	0.96
1:A:110:CYS:HA	1:A:113:LEU:HD12	1.54	0.86
1:A:206:LEU:HD23	1:A:237:ILE:HD12	1.62	0.80
1:A:36:LEU:HD21	1:A:165:TYR:HA	1.66	0.78

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:56:LEU:HD12	1:A:118:LEU:HD23	1.67	0.77
1:A:169:ARG:CB	1:A:169:ARG:HH11	2.02	0.71
1:A:45:CYS:SG	1:A:130:ILE:HD12	2.32	0.69
1:A:169:ARG:NH1	1:A:169:ARG:HB3	2.06	0.66
1:A:221:THR:HG22	1:A:234:VAL:HG12	1.77	0.66
1:A:103:LEU:HB3	1:A:187:PHE:CE2	2.32	0.64
1:A:220:ASP:HB3	1:A:223:THR:HG22	1.78	0.63
1:A:206:LEU:O	1:A:210:ILE:HG13	1.98	0.63
1:A:199:PRO:HG2	1:A:202:VAL:HG23	1.80	0.62
1:A:154:ASN:OD1	1:A:156:ILE:HD13	2.00	0.62
1:A:36:LEU:O	1:A:40:VAL:HG23	2.01	0.61
1:A:196:LEU:HD23	1:A:240:ARG:HH21	1.66	0.60
1:A:43:VAL:O	1:A:47:LYS:HG3	2.01	0.59
1:A:6:GLN:OE1	1:A:170:ARG:HD2	2.01	0.59
1:A:32:LEU:HD13	1:A:71:ALA:HB1	1.86	0.56
1:A:213:LEU:HD11	1:A:227:LEU:HD11	1.87	0.56
1:A:165:TYR:HE1	1:A:178:LEU:HB3	1.71	0.56
1:A:32:LEU:HD11	1:A:74:LEU:HD23	1.88	0.55
1:A:28:VAL:HG11	1:A:130:ILE:HD11	1.89	0.55
1:A:36:LEU:HD21	1:A:165:TYR:CA	2.36	0.54
1:A:135:LYS:HD3	1:A:160:PHE:CE2	2.42	0.54
1:A:135:LYS:HE2	1:A:158:GLU:OE1	2.09	0.53
1:A:68:LYS:HD2	1:A:84:GLU:OE2	2.09	0.53
1:A:220:ASP:OD1	1:A:222:GLU:HB3	2.08	0.53
1:A:156:ILE:H	1:A:156:ILE:HD13	1.75	0.52
1:A:98:PHE:HB3	1:A:101:ALA:HB2	1.93	0.51
1:A:204:ASP:N	1:A:205:PRO:HD2	2.26	0.51
1:A:29:VAL:HA	1:A:54:HIS:O	2.11	0.50
1:A:11:TYR:CD2	1:A:12:LEU:HD23	2.46	0.50
1:A:191:SER:HB3	1:A:194:LYS:HD3	1.93	0.50
1:A:134:ASN:ND2	1:A:161:LYS:HD2	2.27	0.50
1:A:103:LEU:HB3	1:A:187:PHE:CD2	2.47	0.49
1:A:135:LYS:HD3	1:A:160:PHE:CZ	2.48	0.49
1:A:45:CYS:SG	1:A:130:ILE:CD1	3.01	0.49
1:A:49:PHE:HB2	1:A:53:ALA:HB2	1.95	0.48
1:A:196:LEU:CD2	1:A:240:ARG:HH21	2.26	0.48
1:A:122:SER:HB2	1:A:127:SER:O	2.15	0.47
1:A:160:PHE:O	1:A:164:VAL:HG23	2.15	0.47
1:A:140:LEU:HD11	1:A:203:ILE:HG21	1.97	0.47
1:A:60:SER:HB3	1:A:87:ILE:HD11	1.97	0.46
1:A:14:ASP:HB3	2:A:275:HOH:O	2.14	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:233:LEU:O	1:A:233:LEU:HD13	2.16	0.46
1:A:19:GLU:HB3	1:A:153:ILE:HD13	1.98	0.45
1:A:191:SER:HB3	1:A:194:LYS:CD	2.45	0.45
1:A:138:ARG:CZ	1:A:248:LEU:HD22	2.46	0.45
1:A:72:LEU:HD23	1:A:72:LEU:HA	1.76	0.45
1:A:11:TYR:HD2	1:A:12:LEU:HD23	1.82	0.45
1:A:156:ILE:HB	1:A:159:LEU:HD12	1.98	0.45
1:A:152:ALA:HB2	2:A:269:HOH:O	2.16	0.44
1:A:161:LYS:HG2	1:A:200:TYR:CD2	2.52	0.44
1:A:220:ASP:HB3	1:A:223:THR:CG2	2.45	0.44
1:A:15:PHE:O	1:A:19:GLU:HG2	2.18	0.44
1:A:23:ARG:HG3	1:A:23:ARG:HH11	1.83	0.44
1:A:138:ARG:NH2	1:A:248:LEU:HD22	2.33	0.44
1:A:49:PHE:HB3	1:A:52:ASN:HB2	2.00	0.44
1:A:146:PHE:HA	1:A:150:ALA:HB2	1.99	0.43
1:A:155:PRO:HG2	1:A:156:ILE:HG23	2.00	0.43
1:A:54:HIS:HE1	1:A:83:THR:OG1	2.03	0.42
1:A:203:ILE:O	1:A:207:LEU:HG	2.19	0.42
1:A:191:SER:HB3	1:A:194:LYS:CG	2.50	0.41
1:A:19:GLU:HB3	1:A:153:ILE:CD1	2.51	0.41
1:A:88:ALA:HB3	1:A:89:PRO:CD	2.51	0.41
1:A:56:LEU:HD12	1:A:118:LEU:CD2	2.46	0.40
1:A:58:MET:HA	1:A:85:TYR:O	2.20	0.40
1:A:20:VAL:HG21	1:A:28:VAL:HG11	2.03	0.40
1:A:191:SER:HB3	1:A:194:LYS:HG3	2.03	0.40
1:A:214:PHE:HZ	1:A:223:THR:HG23	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	249/268 (93%)	225 (90%)	22 (9%)	2 (1%)	19 23

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	185	ASP
1	A	187	PHE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	215/236 (91%)	201 (94%)	14 (6%)	17 23

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	ASP
1	A	32	LEU
1	A	69	THR
1	A	73	ASN
1	A	86	SER
1	A	133	SER
1	A	156	ILE
1	A	166	GLU
1	A	169	ARG
1	A	185	ASP
1	A	192	ASP
1	A	198	TYR
1	A	201	SER
1	A	240	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	46	GLN
1	A	54	HIS
1	A	67	ASN
1	A	190	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	251/268 (93%)	1.20	36 (14%) 2 3	1, 20, 58, 68	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	185	ASP	13.7
1	A	187	PHE	13.6
1	A	184	ALA	13.4
1	A	191	SER	8.9
1	A	188	VAL	8.5
1	A	194	LYS	6.3
1	A	189	GLY	6.0
1	A	186	LEU	6.0
1	A	195	ASP	5.6
1	A	192	ASP	5.3
1	A	190	GLN	4.9
1	A	193	GLU	4.7
1	A	221	THR	3.8
1	A	216	THR	3.7
1	A	182	PRO	3.6
1	A	224	LEU	3.6
1	A	5	TYR	3.5
1	A	227	LEU	3.4
1	A	222	GLU	3.4
1	A	142	TYR	3.3
1	A	231	GLU	3.2
1	A	213	LEU	2.9
1	A	165	TYR	2.7
1	A	130	ILE	2.7
1	A	31	GLY	2.6
1	A	55	ALA	2.5
1	A	214	PHE	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	36	LEU	2.4
1	A	170	ARG	2.4
1	A	233	LEU	2.4
1	A	181	PRO	2.3
1	A	56	LEU	2.2
1	A	132	THR	2.1
1	A	239	SER	2.1
1	A	226	GLN	2.0
1	A	215	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.