



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 25, 2020 – 05:36 am BST

PDB ID : 5XOZ  
Title : Crystal structure of a Kunitz type trypsin inhibitor from Cicer arietinumL  
Authors : Bendre, A.D.; Suresh, C.G.; Ramasamy, S.  
Deposited on : 2017-05-31  
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	223	
1	B	223	

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2717 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Trypsin protein inhibitor 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	178	Total	C	N	O	S	0	0	0
			1355	871	223	256	5			
1	B	179	Total	C	N	O	S	0	0	0
			1362	874	224	259	5			

There are 58 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLU	-	expression tag	UNP Q9M3Z7
A	2	ALA	-	expression tag	UNP Q9M3Z7
A	3	GLU	-	expression tag	UNP Q9M3Z7
A	4	ALA	-	expression tag	UNP Q9M3Z7
A	5	GLU	-	expression tag	UNP Q9M3Z7
A	6	PHE	-	expression tag	UNP Q9M3Z7
A	201	SER	-	expression tag	UNP Q9M3Z7
A	202	LEU	-	expression tag	UNP Q9M3Z7
A	203	GLU	-	expression tag	UNP Q9M3Z7
A	204	GLN	-	expression tag	UNP Q9M3Z7
A	205	LYS	-	expression tag	UNP Q9M3Z7
A	206	LEU	-	expression tag	UNP Q9M3Z7
A	207	ILE	-	expression tag	UNP Q9M3Z7
A	208	SER	-	expression tag	UNP Q9M3Z7
A	209	GLU	-	expression tag	UNP Q9M3Z7
A	210	GLU	-	expression tag	UNP Q9M3Z7
A	211	ASP	-	expression tag	UNP Q9M3Z7
A	212	LEU	-	expression tag	UNP Q9M3Z7
A	213	ASN	-	expression tag	UNP Q9M3Z7
A	214	SER	-	expression tag	UNP Q9M3Z7
A	215	ALA	-	expression tag	UNP Q9M3Z7
A	216	VAL	-	expression tag	UNP Q9M3Z7
A	217	ASP	-	expression tag	UNP Q9M3Z7
A	218	HIS	-	expression tag	UNP Q9M3Z7
A	219	HIS	-	expression tag	UNP Q9M3Z7

*Continued on next page...*


*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
A	220	HIS	-	expression tag	UNP Q9M3Z7
A	221	HIS	-	expression tag	UNP Q9M3Z7
A	222	HIS	-	expression tag	UNP Q9M3Z7
A	223	HIS	-	expression tag	UNP Q9M3Z7
B	1	GLU	-	expression tag	UNP Q9M3Z7
B	2	ALA	-	expression tag	UNP Q9M3Z7
B	3	GLU	-	expression tag	UNP Q9M3Z7
B	4	ALA	-	expression tag	UNP Q9M3Z7
B	5	GLU	-	expression tag	UNP Q9M3Z7
B	6	PHE	-	expression tag	UNP Q9M3Z7
B	201	SER	-	expression tag	UNP Q9M3Z7
B	202	LEU	-	expression tag	UNP Q9M3Z7
B	203	GLU	-	expression tag	UNP Q9M3Z7
B	204	GLN	-	expression tag	UNP Q9M3Z7
B	205	LYS	-	expression tag	UNP Q9M3Z7
B	206	LEU	-	expression tag	UNP Q9M3Z7
B	207	ILE	-	expression tag	UNP Q9M3Z7
B	208	SER	-	expression tag	UNP Q9M3Z7
B	209	GLU	-	expression tag	UNP Q9M3Z7
B	210	GLU	-	expression tag	UNP Q9M3Z7
B	211	ASP	-	expression tag	UNP Q9M3Z7
B	212	LEU	-	expression tag	UNP Q9M3Z7
B	213	ASN	-	expression tag	UNP Q9M3Z7
B	214	SER	-	expression tag	UNP Q9M3Z7
B	215	ALA	-	expression tag	UNP Q9M3Z7
B	216	VAL	-	expression tag	UNP Q9M3Z7
B	217	ASP	-	expression tag	UNP Q9M3Z7
B	218	HIS	-	expression tag	UNP Q9M3Z7
B	219	HIS	-	expression tag	UNP Q9M3Z7
B	220	HIS	-	expression tag	UNP Q9M3Z7
B	221	HIS	-	expression tag	UNP Q9M3Z7
B	222	HIS	-	expression tag	UNP Q9M3Z7
B	223	HIS	-	expression tag	UNP Q9M3Z7

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Trypsin protein inhibitor 2

Chain A:  77% 20%



- Molecule 1: Trypsin protein inhibitor 2

Chain B:  79% 20%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	38.73 Å 88.22 Å 59.34 Å 90.00° 108.32° 90.00°	Depositor
Resolution (Å)	56.33 – 2.80 36.77 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.3 (56.33-2.80) 99.4 (36.77-2.80)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	7.81 (at 2.81 Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, $R_{free}$	0.196 , 0.221 0.201 , 0.224	Depositor DCC
$R_{free}$ test set	477 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	33.0	Xtriage
Anisotropy	0.146	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 25.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.097 for h,-k,-h-l	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	2717	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.43% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.76	0/1387	0.78	0/1879
1	B	0.83	0/1394	0.82	0/1889
All	All	0.80	0/2781	0.80	0/3768

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1355	0	1343	1	0
1	B	1362	0	1348	0	0
All	All	2717	0	2691	1	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All (1) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:28:LYS:HG2	1:A:93:THR:HG21	1.87	0.56

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	176/223 (79%)	167 (95%)	9 (5%)	0	100	100
1	B	177/223 (79%)	167 (94%)	10 (6%)	0	100	100
All	All	353/446 (79%)	334 (95%)	19 (5%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	148/189 (78%)	144 (97%)	4 (3%)	44	78
1	B	149/189 (79%)	147 (99%)	2 (1%)	69	91
All	All	297/378 (79%)	291 (98%)	6 (2%)	55	84

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	50	ASP
1	B	50	ASP
1	A	103	LYS
1	A	36	ARG
1	B	13	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:



Mol	Chain	Res	Type
1	A	19	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	178/223 (79%)	-0.15	1 (0%) 89 86	16, 29, 45, 55	1 (0%)
1	B	179/223 (80%)	-0.21	1 (0%) 89 86	15, 30, 48, 55	0
All	All	357/446 (80%)	-0.18	2 (0%) 89 86	15, 29, 46, 55	1 (0%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	126	SER	4.4
1	B	190	SER	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.