



Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 08:53 am BST

PDB ID : 5XP8
Title : Crystal structure of T. thermophilus Argonaute protein complexed with a bulge 4A5 on the guide strand
Authors : Sheng, G.; Gogakos, T.; Wang, J.; Zhao, H.; Serganov, A.; Juranek, S.; Tuschl, T.; Patel, J.D.; Wang, Y.
Deposited on : 2017-06-01
Resolution : 3.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1094 (3.10-3.10)
Clashscore	141614	1184 (3.10-3.10)
Ramachandran outliers	138981	1141 (3.10-3.10)
Sidechain outliers	138945	1141 (3.10-3.10)
RSRZ outliers	127900	1067 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	685	<div> <div>2%</div> <div> <div></div> <div>76%</div> <div>21%</div> <div>..</div> </div> </div>
2	E	22	<div> <div>23%</div> <div>41%</div> <div>14%</div> <div>23%</div> </div>
3	F	19	<div> <div>21%</div> <div>42%</div> <div>16%</div> <div>21%</div> </div>
4	C	3	<div> <div>100%</div> </div>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5808 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TtAgo.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	678	Total	C	N	O	S	0	0	0
			5091	3266	938	881	6			

- Molecule 2 is a DNA chain called DNA (5'-D(P*TP*GP*AP*GP*AP*GP*TP*AP*GP*TP*AP*GP*GP*TP*TP*GP*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	17	Total	C	N	O	P	0	0	0
			359	170	67	105	17			

- Molecule 3 is a DNA chain called DNA (5'-D(*CP*AP*AP*CP*C*AP*TP*AP*CP*TP*A P*CP*CP*TP*CP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	F	15	Total	C	N	O	P	0	0	0
			296	143	52	87	14			

- Molecule 4 is a DNA chain called DNA (5'-D(*AP*GP*T)-3').


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	C	3	Total	C	N	O	P	0	0	0
			59	30	12	15	2			

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	3	Total	Mg	1	0
			3	3		

i

- Molecule 1: TtAgo



DT	DA	DT	DA	C4	A5	A6	C7	C8	T9	A10	C11	T12	A13	C14	C15	T16	C17	G18
----	----	----	----	----	----	----	----	----	----	-----	-----	-----	-----	-----	-----	-----	-----	-----

- Molecule 4: DNA (5'-D(*AP*GP*T)-3')

Chain C:  100%

There are no outlier residues recorded for this chain.

4 Data and refinement statistics

Property	Value	Source
Space group	P 41 3 2	Depositor
Cell constants a, b, c, α , β , γ	202.14Å 202.14Å 202.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.20 – 3.10 45.20 – 3.10	Depositor EDS
% Data completeness (in resolution range)	98.9 (45.20-3.10) 98.9 (45.20-3.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.18 (at 3.12Å)	Xtriage
Refinement program	PHENIX 1.8.2_1309	Depositor
R, R_{free}	0.214 , 0.276 0.217 , 0.280	Depositor DCC
R_{free} test set	1313 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	91.1	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 40.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5808	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.46	0/5213	0.77	4/7110 (0.1%)
2	E	0.99	1/403 (0.2%)	1.63	8/621 (1.3%)
3	F	1.15	2/330 (0.6%)	1.62	13/505 (2.6%)
4	C	0.72	0/66	1.42	0/101
All	All	0.57	3/6012 (0.0%)	0.94	25/8337 (0.3%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	9	DT	O3'-P	-10.40	1.48	1.61
2	E	1	DT	OP3-P	-9.44	1.49	1.61
3	F	7	DC	O3'-P	-9.23	1.50	1.61

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	9	DG	O4'-C1'-N9	9.93	114.95	108.00
2	E	11	DA	O4'-C1'-N9	9.10	114.37	108.00
2	E	10	DT	O4'-C4'-C3'	-8.74	100.76	106.00
3	F	16	DT	N3-C4-O4	8.59	125.05	119.90
3	F	7	DC	C1'-O4'-C4'	-8.47	101.63	110.10
3	F	12	DT	O4'-C1'-N1	7.68	113.38	108.00
3	F	14	DC	O4'-C4'-C3'	-7.53	101.48	106.00
3	F	7	DC	O4'-C1'-N1	7.32	113.12	108.00
1	A	365	LEU	CA-CB-CG	7.12	131.67	115.30
3	F	9	DT	O5'-P-OP2	-6.80	99.58	105.70
3	F	16	DT	C5-C4-O4	-6.70	120.21	124.90
1	A	358	PRO	N-CA-CB	6.38	110.96	103.30
3	F	7	DC	C3'-C2'-C1'	-6.23	95.03	102.50
3	F	11	DC	O4'-C1'-N1	6.21	112.35	108.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	143	PRO	N-CA-CB	5.93	110.42	103.30
2	E	9	DG	O4'-C1'-C2'	5.92	110.64	105.90
3	F	11	DC	C1'-O4'-C4'	-5.87	104.23	110.10
2	E	1	DT	N3-C4-O4	5.78	123.37	119.90
2	E	14	DT	N3-C4-O4	5.75	123.35	119.90
3	F	7	DC	P-O3'-C3'	5.58	126.39	119.70
3	F	4	DC	O4'-C1'-N1	5.56	111.89	108.00
1	A	175	CYS	N-CA-CB	-5.41	100.86	110.60
2	E	9	DG	N3-C4-N9	-5.32	122.81	126.00
3	F	7	DC	O4'-C1'-C2'	-5.17	101.76	105.90
2	E	4	DG	O4'-C4'-C3'	-5.00	102.50	104.50

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5091	0	4990	93	0
2	E	359	0	194	12	0
3	F	296	0	170	9	0
4	C	59	0	33	0	0
5	A	3	0	0	0	0
All	All	5808	0	5387	103	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (103) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:12:DG:H1	3:F:8:DC:H42	1.14	0.87
2:E:12:DG:H1	3:F:8:DC:N4	1.78	0.81
2:E:16:DG:H1	3:F:4:DC:H42	1.33	0.74
1:A:315:ARG:HH11	1:A:589:ALA:HB3	1.53	0.73

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:410:THR:O	1:A:436:ASN:HA	1.91	0.70
1:A:295:SER:O	1:A:299:ARG:HG3	1.94	0.67
1:A:365:LEU:HD13	1:A:376:LEU:HD21	1.79	0.65
1:A:513:ARG:NH2	1:A:551:GLN:O	2.32	0.63
2:E:12:DG:N2	3:F:8:DC:N3	2.44	0.62
1:A:350:ARG:NH2	1:A:354:ALA:HB3	2.16	0.61
1:A:28:ARG:HB3	1:A:93:LYS:HE3	1.81	0.60
1:A:365:LEU:HD11	1:A:378:LEU:HD11	1.83	0.60
3:F:8:DC:H2''	3:F:9:DT:H5'	1.84	0.60
1:A:295:SER:HA	1:A:306:PRO:HG3	1.83	0.59
1:A:340:ARG:NH1	1:A:499:GLY:O	2.34	0.59
1:A:473:LEU:HB3	1:A:541:VAL:HG12	1.85	0.59
1:A:350:ARG:NH2	1:A:354:ALA:CB	2.66	0.58
1:A:17:ARG:HG3	1:A:303:LEU:O	2.03	0.57
1:A:439:LEU:HD11	1:A:447:TRP:HB3	1.85	0.57
1:A:319:PRO:HB3	1:A:637:HIS:CG	2.40	0.56
1:A:315:ARG:NH1	1:A:589:ALA:HB3	2.20	0.56
1:A:109:LEU:HB2	1:A:157:VAL:HG21	1.87	0.56
1:A:128:TRP:CZ3	1:A:172:ARG:HG2	2.41	0.55
2:E:15:DT:H3	3:F:5:DA:H61	1.54	0.55
1:A:350:ARG:HG3	1:A:355:GLN:O	2.07	0.55
1:A:494:VAL:HG13	1:A:530:PHE:HD1	1.71	0.55
1:A:615:ARG:O	1:A:651:ARG:NH1	2.39	0.54
2:E:11:DA:H2''	2:E:12:DG:H5''	1.90	0.54
1:A:590:ASP:OD2	3:F:18:DG:N1	2.41	0.54
1:A:17:ARG:HD3	1:A:303:LEU:HA	1.90	0.53
1:A:114:ARG:HD3	1:A:132:LEU:HD11	1.92	0.52
1:A:416:GLU:HG2	1:A:417:ASP:N	2.24	0.52
1:A:350:ARG:CZ	1:A:354:ALA:HB3	2.40	0.51
1:A:172:ARG:NH1	2:E:9:DG:OP1	2.39	0.51
1:A:352:ASP:HB3	1:A:411:PRO:HD3	1.91	0.51
1:A:43:TYR:CZ	2:E:17:DT:H2''	2.46	0.51
1:A:606:VAL:HG13	1:A:611:ARG:HH12	1.75	0.51
1:A:408:VAL:HG13	1:A:434:ILE:HA	1.93	0.49
3:F:13:DA:H2'	3:F:14:DC:O4'	2.12	0.49
1:A:195:ASN:ND2	1:A:200:ARG:HB2	2.28	0.49
1:A:439:LEU:HA	1:A:446:ARG:NH2	2.27	0.49
1:A:439:LEU:O	1:A:440:ARG:HG2	2.13	0.49
1:A:76:GLU:HG2	1:A:89:ARG:NE	2.28	0.48
1:A:639:THR:HA	1:A:653:PRO:HA	1.94	0.48
1:A:431:PRO:HB2	1:A:457:LYS:HB3	1.96	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:444:ARG:HA	1:A:447:TRP:NE1	2.28	0.47
1:A:10:PHE:CE2	1:A:584:VAL:HG12	2.50	0.47
1:A:79:LEU:O	1:A:79:LEU:HD12	2.16	0.46
1:A:142:GLY:O	1:A:144:GLY:N	2.49	0.46
1:A:128:TRP:HZ3	1:A:172:ARG:HG2	1.81	0.46
1:A:409:LEU:HD21	1:A:435:LEU:HD23	1.98	0.46
1:A:8:GLU:OE1	1:A:310:ARG:NH1	2.49	0.46
1:A:464:ALA:HB1	1:A:497:ASP:O	2.16	0.46
1:A:300:ARG:HA	1:A:300:ARG:HD3	1.76	0.45
1:A:446:ARG:HG3	2:E:2:DG:C8	2.51	0.45
1:A:286:ARG:HD2	1:A:613:THR:HG21	1.99	0.45
1:A:124:LEU:HD23	1:A:124:LEU:HA	1.81	0.45
1:A:422:LYS:NZ	1:A:432:SER:O	2.50	0.45
1:A:181:ALA:HA	1:A:184:ALA:HB3	1.98	0.44
1:A:523:LEU:HD21	1:A:561:LEU:HD11	1.99	0.44
1:A:246:ASP:HB3	1:A:249:ASP:CB	2.48	0.44
1:A:479:ALA:HA	1:A:487:PHE:O	2.18	0.44
1:A:543:LEU:HB3	1:A:570:LEU:HD12	1.99	0.44
1:A:189:LEU:HD12	1:A:189:LEU:H	1.83	0.44
1:A:425:LEU:HD12	1:A:432:SER:HB3	2.00	0.44
1:A:607:HIS:O	1:A:608:ARG:HG2	2.18	0.43
2:E:13:DG:H2'	2:E:14:DT:H6	1.83	0.43
1:A:179:LEU:HD23	1:A:258:THR:HG23	2.00	0.43
1:A:439:LEU:HA	1:A:446:ARG:HH21	1.82	0.43
1:A:671:ILE:HA	1:A:674:LEU:HD12	2.00	0.43
1:A:548:ARG:HA	1:A:572:SER:OG	2.19	0.43
1:A:174:LEU:HD12	1:A:174:LEU:C	2.38	0.43
1:A:43:TYR:CE1	2:E:17:DT:H2''	2.54	0.43
1:A:98:LEU:HA	1:A:98:LEU:HD23	1.86	0.43
1:A:41:GLU:O	1:A:45:LEU:HB2	2.19	0.42
1:A:246:ASP:HB3	1:A:249:ASP:HB3	2.01	0.42
1:A:324:ARG:N	1:A:337:GLY:O	2.41	0.42
1:A:361:LEU:O	1:A:365:LEU:HG	2.20	0.42
1:A:58:VAL:HG22	1:A:112:LEU:HD13	2.00	0.42
2:E:15:DT:H3	3:F:5:DA:N6	2.18	0.42
1:A:180:GLU:OE2	1:A:240:ARG:HD2	2.19	0.42
1:A:350:ARG:HD2	1:A:409:LEU:HB2	2.02	0.42
1:A:348:LEU:HD21	1:A:350:ARG:HD3	2.01	0.41
1:A:4:LEU:HD13	1:A:316:LEU:HD23	2.01	0.41
1:A:426:LEU:HA	1:A:426:LEU:HD12	1.71	0.41
1:A:121:LEU:HD11	1:A:153:LEU:HD12	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:224:LEU:HD12	1:A:224:LEU:HA	1.77	0.41
1:A:362:ARG:HG2	1:A:366:LEU:HD12	2.03	0.41
1:A:134:VAL:O	1:A:150:GLY:HA3	2.21	0.41
1:A:348:LEU:HD22	1:A:357:TRP:HA	2.03	0.41
1:A:106:ARG:HG3	1:A:157:VAL:HG23	2.03	0.41
1:A:217:LEU:HA	1:A:217:LEU:HD23	1.93	0.41
1:A:45:LEU:HD12	1:A:45:LEU:HA	1.75	0.41
1:A:665:GLU:HG3	1:A:684:PHE:HA	2.02	0.41
1:A:125:GLU:HA	1:A:126:GLY:HA2	1.68	0.41
1:A:548:ARG:HG2	1:A:574:ARG:NE	2.36	0.41
1:A:303:LEU:HA	1:A:303:LEU:HD23	1.88	0.40
1:A:588:LEU:HD21	1:A:614:PRO:HG3	2.03	0.40
1:A:89:ARG:HG2	1:A:91:TYR:CZ	2.56	0.40
1:A:174:LEU:HD13	1:A:176:GLU:OE1	2.22	0.40
1:A:17:ARG:NH2	1:A:23:GLU:OE1	2.54	0.40
1:A:628:LEU:HD12	1:A:628:LEU:HA	1.71	0.40
1:A:76:GLU:HG2	1:A:89:ARG:CD	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	674/685 (98%)	634 (94%)	25 (4%)	15 (2%)	6 29

All (15) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	436	ASN
1	A	439	LEU
1	A	214	GLU
1	A	216	PRO

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	250	PRO
1	A	440	ARG
1	A	143	PRO
1	A	184	ALA
1	A	438	PRO
1	A	205	LEU
1	A	253	PRO
1	A	256	HIS
1	A	281	LEU
1	A	484	SER
1	A	231	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	475/549 (86%)	458 (96%)	17 (4%)	35 67

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	60	MET
1	A	112	LEU
1	A	177	MET
1	A	193	VAL
1	A	236	ARG
1	A	264	VAL
1	A	287	ARG
1	A	289	ARG
1	A	426	LEU
1	A	462	VAL
1	A	505	LEU
1	A	513	ARG
1	A	573	VAL
1	A	628	LEU
1	A	629	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	640	ARG
1	A	665	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	461	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	678/685 (98%)	-0.13	14 (2%) 63 43	14, 55, 97, 134	0
2	E	17/22 (77%)	-0.57	0 100 100	30, 44, 91, 113	0
3	F	15/19 (78%)	-0.34	0 100 100	42, 62, 137, 142	0
4	C	3/3 (100%)	0.02	0 100 100	76, 76, 98, 122	0
All	All	713/729 (97%)	-0.14	14 (1%) 65 44	14, 56, 98, 142	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	85	THR	3.7
1	A	33	LEU	3.6
1	A	204	LEU	3.2
1	A	203	GLU	3.0
1	A	64	LEU	2.9
1	A	63	GLY	2.9
1	A	245	ALA	2.5
1	A	346	LEU	2.5
1	A	37	PRO	2.4
1	A	279	LEU	2.4
1	A	62	ASP	2.3
1	A	79	LEU	2.3
1	A	190	PRO	2.1
1	A	31	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	MG	A	701	1/1	0.91	0.40	12,12,12,12	0
5	MG	A	703	1/1	0.99	0.11	28,28,28,28	0
5	MG	A	702	1/1	-	-	29,29,29,29	1

6.5 Other polymers [i](#)

There are no such residues in this entry.