



# Full wwPDB X-ray Structure Validation Report ⓘ

May 18, 2020 – 03:45 am BST

PDB ID : 4XQA  
Title : CRYSTAL STRUCTURE OF AD37 FIBER KNOB IN COMPLEX WITH  
TRIVALENT SIALIC ACID INHIBITOR ME0462  
Authors : Stehle, T.; Liaci, A.M.  
Deposited on : 2015-01-19  
Resolution : 1.41 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.41 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1714 (1.40-1.40)
Clashscore	141614	1812 (1.40-1.40)
Ramachandran outliers	138981	1763 (1.40-1.40)
Sidechain outliers	138945	1762 (1.40-1.40)
RSRZ outliers	127900	1674 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	194	<div> <div>10%</div> <div> <div></div> <div>85%</div> <div>10%</div> <div>5%</div> </div> </div>
1	B	194	<div> <div>8%</div> <div> <div></div> <div>88%</div> <div>6%</div> <div>5%</div> </div> </div>
1	C	194	<div> <div>4%</div> <div> <div></div> <div>90%</div> <div>5%</div> <div>6%</div> </div> </div>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5010 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fiber.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	184	Total	C	N	O	S	0	2	0
			1454	935	236	279	4			
1	B	184	Total	C	N	O	S	0	5	0
			1463	941	235	283	4			
1	C	183	Total	C	N	O	S	0	1	0
			1446	930	235	277	4			

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	172	GLY	-	expression tag	UNP Q64823
A	173	ALA	-	expression tag	UNP Q64823
A	174	MET	-	expression tag	UNP Q64823
A	175	GLY	-	expression tag	UNP Q64823
A	176	SER	-	expression tag	UNP Q64823
B	172	GLY	-	expression tag	UNP Q64823
B	173	ALA	-	expression tag	UNP Q64823
B	174	MET	-	expression tag	UNP Q64823
B	175	GLY	-	expression tag	UNP Q64823
B	176	SER	-	expression tag	UNP Q64823
C	172	GLY	-	expression tag	UNP Q64823
C	173	ALA	-	expression tag	UNP Q64823
C	174	MET	-	expression tag	UNP Q64823
C	175	GLY	-	expression tag	UNP Q64823
C	176	SER	-	expression tag	UNP Q64823

- Molecule 2 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	C	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Zn	0	0
			1	1		
3	A	1	Total	Zn	0	0
			1	1		
3	C	1	Total	Zn	0	0
			1	1		

- Molecule 4 is (1-{2-[bis(2-{4-[({(6R)-5-(acetylamino)-3,5-dideoxy-6-[(1R,2R)-1,2,3-trihydroxypropyl]-beta-L-threo-hex-2-ulopyranonosyl}oxy)methyl]-1H-1,2,3-triazol-1-yl}ethyl)amino]ethyl}-1H-1,2,3-triazol-4-yl)methyl (6R)-5-(acetylamino)-3,5-dideoxy-6-[(1R,2R)-1,2,3-trihydroxypropyl]-beta-L-threo-hex-2-ulopyranosidonic acid (three-letter code: 423) (formula: C<sub>48</sub>H<sub>75</sub>N<sub>13</sub>O<sub>27</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	0
			88	48	13	27		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	160	Total O 160 160	0	0
5	B	202	Total O 202 202	0	0
5	C	178	Total O 178 178	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Fiber



#### • Molecule 1: Fiber



#### • Molecule 1: Fiber



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	68.23Å 75.62Å 131.11Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.27 – 1.41 47.27 – 1.41	Depositor EDS
% Data completeness (in resolution range)	99.6 (47.27-1.41) 99.6 (47.27-1.41)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.08 (at 1.41Å)	Xtriage
Refinement program	PHENIX 1.8.1069	Depositor
R, $R_{free}$	0.143 , 0.172 0.145 , 0.174	Depositor DCC
$R_{free}$ test set	6567 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.7	Xtriage
Anisotropy	0.345	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 55.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	5010	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.90% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, 423, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.55	0/1493	0.66	0/2028
1	B	0.52	0/1512	0.67	0/2058
1	C	0.58	0/1483	0.68	0/2016
All	All	0.55	0/4488	0.67	0/6102

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1454	0	1453	21	1
1	B	1463	0	1452	12	0
1	C	1446	0	1435	7	0
2	A	8	0	6	0	0
2	B	4	0	3	0	0
2	C	4	0	3	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
4	B	88	0	72	2	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	160	0	0	4	1
5	B	202	0	0	2	0
5	C	178	0	0	4	0
All	All	5010	0	4424	35	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:299:SER:O	5:A:501:HOH:O	1.85	0.93
1:A:300:LYS:HE2	5:C:501:HOH:O	1.67	0.93
1:A:186[B]:LEU:HD21	1:A:267:TRP:CE2	2.08	0.88
1:A:186[B]:LEU:CD2	1:A:267:TRP:CZ2	2.57	0.88
1:C:351:GLU:OE2	5:C:501:HOH:O	1.92	0.86
1:A:186[B]:LEU:HD21	1:A:267:TRP:CZ2	2.11	0.85
4:B:401:423:NCZ	5:B:501:HOH:O	2.09	0.84
1:C:318:ASP:OD2	5:C:502:HOH:O	2.05	0.75
1:A:270:ARG:HD3	1:B:215[B]:SER:OG	2.00	0.60
1:A:318:ASP:OD2	5:A:502:HOH:O	2.17	0.59
1:A:300:LYS:CE	5:C:501:HOH:O	2.34	0.58
4:B:401:423:H53	4:B:401:423:NCX	2.19	0.58
1:A:186[B]:LEU:CD2	1:A:267:TRP:CH2	2.88	0.56
1:B:188:THR:O	1:B:189[B]:THR:HB	2.07	0.55
1:A:213:CYS:O	1:C:211:THR:HG21	2.06	0.55
1:B:189[B]:THR:HG22	1:B:191:ASP:H	1.73	0.53
1:B:270[A]:ARG:NE	1:B:272:GLY:O	2.42	0.53
1:B:189[B]:THR:HG23	1:B:190:PRO:HD2	1.92	0.51
1:B:183:THR:HB	5:B:662:HOH:O	2.12	0.50
1:A:330:GLU:OE2	1:A:338:THR:HG21	2.12	0.49
1:A:295:LYS:HD2	1:A:328:ASN:HB3	1.96	0.48
1:A:270:ARG:NE	1:A:272:GLY:O	2.41	0.48
1:B:187:TRP:CH2	1:B:190:PRO:HD3	2.49	0.48
1:A:270:ARG:NH1	1:B:215[A]:SER:OG	2.40	0.47
1:A:211:THR:HG21	1:B:213:CYS:O	2.16	0.46
1:B:211:THR:HG21	1:C:213:CYS:O	2.16	0.46
1:A:185:THR:OG1	1:B:215[B]:SER:HB2	2.16	0.46
1:A:186[B]:LEU:HD23	1:A:267:TRP:CH2	2.53	0.44
1:A:215:SER:HB2	1:C:270:ARG:HH11	1.83	0.43
1:A:187:TRP:CH2	1:A:190:PRO:HD3	2.54	0.42

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:193[B]:SER:HA	1:B:194:PRO:HD3	1.94	0.41
1:C:270:ARG:NE	1:C:272:GLY:O	2.40	0.41
1:C:251:ASN:HB3	1:C:257:LEU:HD21	2.02	0.41
1:A:242:LYS:HE2	5:A:584:HOH:O	2.21	0.41
1:A:316:LYS:CE	5:A:515:HOH:O	2.70	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:302:TYR:OH	5:A:649:HOH:O[4_455]	2.18	0.02

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	184/194 (95%)	178 (97%)	6 (3%)	0	100	100
1	B	187/194 (96%)	178 (95%)	9 (5%)	0	100	100
1	C	182/194 (94%)	176 (97%)	6 (3%)	0	100	100
All	All	553/582 (95%)	532 (96%)	21 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	164/171 (96%)	163 (99%)	1 (1%)	86	70
1	B	165/171 (96%)	164 (99%)	1 (1%)	86	70
1	C	162/171 (95%)	161 (99%)	1 (1%)	86	70
All	All	491/513 (96%)	488 (99%)	3 (1%)	86	70

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	187	TRP
1	B	187	TRP
1	C	187	TRP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 3 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	ACT	B	402	-	1,3,3	1.11	0	0,3,3	0.00	-
4	423	B	401	-	81,93,93	2.29	14 (17%)	102,135,135	2.18	22 (21%)
2	ACT	C	401	-	1,3,3	1.12	0	0,3,3	0.00	-
2	ACT	A	401	-	1,3,3	1.20	0	0,3,3	0.00	-
2	ACT	A	403	3	1,3,3	2.01	1 (100%)	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	423	B	401	-	-	4/69/147/147	0/6/6/6

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	401	423	NCZ-NCY	-11.41	1.13	1.34
4	B	401	423	NDD-NDC	-7.89	1.19	1.34
4	B	401	423	CCV-NCU	6.83	1.43	1.34
4	B	401	423	NCY-NCX	-6.01	1.22	1.34
4	B	401	423	NCU-NCT	-3.96	1.27	1.34
4	B	401	423	NDE-NDD	-3.86	1.27	1.34
4	B	401	423	CDH-NCS	3.52	1.53	1.47
4	B	401	423	CCR-CCV	2.93	1.40	1.36
4	B	401	423	NCT-NCS	-2.71	1.29	1.34
4	B	401	423	CBT-CBS	2.68	1.55	1.53
4	B	401	423	CAZ-CAY	2.48	1.55	1.53
4	B	401	423	CDG-NDC	2.27	1.51	1.47
4	B	401	423	CDB-CDF	-2.24	1.32	1.36
4	B	401	423	CAY-NAX	2.02	1.49	1.45
2	A	403	ACT	CH3-C	2.01	1.51	1.48

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	401	423	NCU-NCT-NCS	7.65	113.08	107.31
4	B	401	423	NCZ-NCY-NCX	7.25	112.78	107.31
4	B	401	423	NDE-NDD-NDC	6.71	112.37	107.31
4	B	401	423	CDJ-NDK-CDL	5.75	125.11	111.44
4	B	401	423	CDG-NDC-CDB	-5.60	116.64	129.82

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	401	423	CDM-CDI-NCX	5.00	119.25	110.83
4	B	401	423	CBK-CBI-CBC	-4.95	103.64	113.03
4	B	401	423	CCR-CCV-NCU	-4.54	104.59	111.34
4	B	401	423	CAQ-CAO-CAI	-4.27	104.94	113.03
4	B	401	423	CDJ-CDG-NDC	4.05	117.65	110.83
4	B	401	423	CCE-CCC-CBW	-3.85	105.74	113.03
4	B	401	423	CCM-CCV-CCR	-3.56	121.73	128.45
4	B	401	423	CBB-CAZ-CAY	-3.53	104.55	109.98
4	B	401	423	CDB-CDF-NDE	-3.34	106.38	111.34
4	B	401	423	CDI-NCX-CCW	3.02	136.92	129.82
4	B	401	423	CDJ-NDK-CDM	2.87	118.26	111.44
4	B	401	423	CAH-CAF-CAE	-2.85	105.59	109.98
4	B	401	423	CCW-CDA-NCZ	-2.62	107.45	111.34
4	B	401	423	CBV-CBT-CBS	-2.36	106.35	109.98
4	B	401	423	OCL-CCM-CCV	-2.13	105.19	110.64
4	B	401	423	OAA-CAB-NAD	2.07	125.76	121.95
4	B	401	423	OAU-CAV-NAX	2.01	125.64	121.95

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	401	423	CDI-CDM-NDK-CDJ
4	B	401	423	CDM-CDI-NCX-CCW
4	B	401	423	CDH-CDL-NDK-CDM
4	B	401	423	NDC-CDG-CDJ-NDK


There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	401	423	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	184/194 (94%)	0.41	20 (10%) 5 5	23, 33, 48, 56	1 (0%)
1	B	184/194 (94%)	0.22	15 (8%) 11 10	20, 32, 48, 58	3 (1%)
1	C	183/194 (94%)	0.08	7 (3%) 40 40	20, 27, 43, 53	3 (1%)
All	All	551/582 (94%)	0.24	42 (7%) 13 13	20, 30, 47, 58	7 (1%)

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	183	THR	7.7
1	A	276	VAL	6.9
1	A	266	TYR	5.9
1	A	278	THR	5.4
1	A	277	SER	5.0
1	A	258	ASP	4.7
1	B	276	VAL	4.4
1	A	183	THR	4.3
1	B	266	TYR	4.2
1	C	276	VAL	4.1
1	B	277	SER	3.9
1	B	298	ASN	3.8
1	B	182	ASP	3.8
1	A	274	SER	3.4
1	A	279	ALA	3.4
1	A	271	SER	3.3
1	B	278	THR	3.3
1	C	359	PHE	3.2
1	A	275	ASN	3.0
1	A	201	ASP	3.0
1	A	282	LYS	2.8
1	A	281	GLU	2.7
1	B	359	PHE	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	277	SER	2.6
1	C	271	SER	2.6
1	B	259	ASN	2.5
1	A	182	ASP	2.5
1	A	199	ALA	2.5
1	C	298	ASN	2.5
1	C	183	THR	2.4
1	A	272	GLY	2.3
1	B	236	LYS	2.3
1	B	218	LEU	2.2
1	B	219	ALA	2.2
1	B	237	THR	2.2
1	B	272	GLY	2.1
1	A	200	GLN	2.1
1	A	193	SER	2.1
1	A	197	THR	2.1
1	A	192	THR	2.1
1	B	357	PHE	2.1
1	C	218	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ACT	A	401	4/4	0.83	0.31	48,49,50,50	4
2	ACT	B	402	4/4	0.84	0.14	44,45,45,46	4
4	423	B	401	88/88	0.89	0.12	23,31,50,52	0
2	ACT	A	403	4/4	0.94	0.10	40,44,44,46	0

*Continued on next page...*






*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ACT	C	401	4/4	0.95	0.07	36,37,37,37	4
3	ZN	A	402	1/1	1.00	0.06	34,34,34,34	1
3	ZN	C	402	1/1	1.00	0.04	32,32,32,32	0
3	ZN	B	403	1/1	1.00	0.03	32,32,32,32	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around 423 B 401:**

2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.