



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 9, 2020 – 01:09 PM BST

PDB ID : 2XR9
Title : Crystal structure of Autotaxin (ENPP2)
Authors : Kamtekar, S.; Hausmann, J.; Day, J.E.; Christodoulou, E.; Perrakis, A.
Deposited on : 2010-09-13
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	827	<div> <div>5%</div> <div> <div></div> <div>85%</div> <div>9%</div> <div>• 5%</div> </div> </div>
2	B	3	<div> <div></div> <div>100%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	IOD	A	1873	-	-	X	-
7	IOD	A	1878	-	-	X	X
7	IOD	A	1879	-	-	X	-
7	IOD	A	1883	-	-	X	-

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 6753 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ECTONUCLEOTIDE PYROPHOSPHATASE/PHOSPHODIESTERASE FAMILY MEMBER 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	783	Total	C	N	O	S	0	1	0
			6334	4019	1094	1172	49			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	410	ALA	ASN	engineered mutation	UNP Q64610
A	591	THR	ARG	conflict	UNP Q64610

- Molecule 2 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 3 is THIOCYANATE ION (three-letter code: SCN) (formula: CNS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	S	0	0
			3	1	1	1		
3	A	1	Total	C	N	S	0	0
			3	1	1	1		
3	A	1	Total	C	N	S	0	0
			3	1	1	1		
3	A	1	Total	C	N	S	0	0
			3	1	1	1		


- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Zn	0	0
			2	2		

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Ca	0	0
			1	1		

- Molecule 6 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 7 is IODIDE ION (three-letter code: IOD) (formula: I).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	28	Total	I	0	0
			28	28		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	332	Total	O	0	0
			332	332		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ECTONUCLEOTIDE PYROPHOSPHATASE/PHOSPHODIESTERASE FAMILY MEMBER 2



- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	53.81Å 63.29Å 70.47Å 98.78° 106.22° 99.77°	Depositor
Resolution (Å)	19.98 – 2.05 19.98 – 2.05	Depositor EDS
% Data completeness (in resolution range)	96.8 (19.98-2.05) 88.6 (19.98-2.05)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.19 (at 2.06Å)	Xtriage
Refinement program	REFMAC 5.5.0110	Depositor
R, R_{free}	0.173 , 0.221 0.196 , 0.232	Depositor DCC
R_{free} test set	2386 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	31.6	Xtriage
Anisotropy	0.218	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 50.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6753	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.90% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, BMA, NAG, PO4, SCN, IOD, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.77	2/6515 (0.0%)	0.82	14/8832 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	129	CYS	CB-SG	-7.08	1.70	1.82
1	A	854	TYR	CD2-CE2	5.47	1.47	1.39

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	606	ARG	NE-CZ-NH2	7.85	124.22	120.30
1	A	299	ARG	NE-CZ-NH2	-7.51	116.54	120.30
1	A	798	ARG	NE-CZ-NH1	7.20	123.90	120.30
1	A	299	ARG	NE-CZ-NH1	6.98	123.79	120.30
1	A	798	ARG	NE-CZ-NH2	-6.89	116.85	120.30
1	A	393	ARG	NE-CZ-NH2	-6.72	116.94	120.30
1	A	746	ARG	NE-CZ-NH1	6.51	123.56	120.30
1	A	606	ARG	NE-CZ-NH1	-5.69	117.45	120.30
1	A	596	LEU	CB-CG-CD1	-5.64	101.42	111.00
1	A	393	ARG	NE-CZ-NH1	5.33	122.97	120.30
1	A	827	ARG	NE-CZ-NH2	-5.32	117.64	120.30
1	A	746	ARG	NE-CZ-NH2	-5.27	117.66	120.30
1	A	843	ARG	NE-CZ-NH2	-5.25	117.67	120.30
1	A	827	ARG	NE-CZ-NH1	5.19	122.90	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6334	0	6096	68	0
2	B	39	0	34	0	0
3	A	12	0	0	0	0
4	A	2	0	0	0	0
5	A	1	0	0	0	0
6	A	5	0	0	0	0
7	A	28	0	0	20	0
8	A	332	0	0	3	0
All	All	6753	0	6130	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:124:LEU:HD21	7:A:1878:IOD:I	1.99	1.32
1:A:124:LEU:HD11	7:A:1878:IOD:I	2.12	1.19
7:A:1873:IOD:I	7:A:1883:IOD:I	3.10	1.10
1:A:124:LEU:CD2	7:A:1878:IOD:I	2.82	0.97
1:A:278:SER:OG	7:A:1876:IOD:I	2.53	0.96
1:A:331:ILE:HD13	7:A:1879:IOD:I	2.35	0.96
1:A:269[B]:ARG:CG	1:A:269[B]:ARG:HH21	1.79	0.95
1:A:124:LEU:CD1	7:A:1878:IOD:I	2.91	0.88
1:A:716:GLN:OE1	7:A:1873:IOD:I	2.70	0.79
1:A:709:LYS:NZ	7:A:1883:IOD:I	2.86	0.79
1:A:269[B]:ARG:HG3	1:A:269[B]:ARG:HH21	1.48	0.78
1:A:313:SER:HB3	1:A:324:MET:HE1	1.67	0.76
1:A:557:TYR:O	1:A:656:VAL:HG13	1.86	0.75
1:A:209:THR:OG1	1:A:358:ASP:OD2	2.04	0.74
1:A:124:LEU:CG	7:A:1878:IOD:I	3.06	0.73
7:A:1898:IOD:I	8:A:2253:HOH:O	2.75	0.73
7:A:1888:IOD:I	7:A:1889:IOD:I	3.47	0.72
1:A:313:SER:C	1:A:324:MET:HE3	2.12	0.70
1:A:269[B]:ARG:HG2	1:A:269[B]:ARG:HH21	1.55	0.69

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:633:TYR:CE1	1:A:635:ILE:HD11	2.28	0.68
1:A:633:TYR:HE1	1:A:635:ILE:HD11	1.61	0.66
1:A:281:HIS:ND1	7:A:1879:IOD:I	2.99	0.66
1:A:546:GLU:HG2	1:A:602:ALA:HB1	1.77	0.65
1:A:313:SER:O	1:A:324:MET:HE3	1.97	0.64
1:A:635:ILE:HD13	1:A:720:VAL:CG1	2.28	0.64
1:A:201:MET:CE	1:A:356:VAL:HG23	2.31	0.60
1:A:176:SER:HA	1:A:179:LYS:HD3	1.84	0.60
1:A:313:SER:CB	1:A:324:MET:HE1	2.33	0.58
1:A:546:GLU:HG2	1:A:602:ALA:CB	2.34	0.57
1:A:201:MET:HE2	1:A:356:VAL:HG23	1.86	0.57
1:A:313:SER:HB3	1:A:324:MET:CE	2.32	0.57
1:A:269[A]:ARG:NH2	8:A:2068:HOH:O	2.38	0.56
1:A:296:ASP:OD1	1:A:299:ARG:NH2	2.38	0.56
1:A:222:PRO:HA	1:A:225:HIS:CE1	2.41	0.55
1:A:201:MET:HE1	1:A:356:VAL:CG2	2.36	0.54
1:A:153:VAL:HG22	1:A:154:PRO:HD2	1.89	0.54
1:A:134:GLN:HG3	7:A:1878:IOD:I	2.79	0.53
1:A:61:ARG:HH11	1:A:68:VAL:HG21	1.75	0.52
1:A:269[B]:ARG:NH2	1:A:269[B]:ARG:HG2	2.20	0.52
1:A:606:ARG:HG2	1:A:606:ARG:HH21	1.74	0.51
1:A:682:PRO:HB3	1:A:716:GLN:HB3	1.92	0.51
1:A:526:THR:OG1	1:A:827:ARG:HD3	2.10	0.51
1:A:606:ARG:CG	1:A:606:ARG:HH21	2.23	0.51
1:A:555:ILE:HG23	1:A:656:VAL:HG11	1.93	0.50
1:A:793:PHE:CD2	1:A:795:LEU:HG	2.46	0.50
1:A:61:ARG:NH1	1:A:68:VAL:HG21	2.29	0.48
1:A:331:ILE:CD1	7:A:1879:IOD:I	3.22	0.48
1:A:534:LEU:HD12	1:A:537:ASN:OD1	2.14	0.47
1:A:455:ARG:NH2	8:A:2142:HOH:O	2.45	0.47
1:A:665:ASN:OD1	1:A:667:LEU:HB3	2.16	0.46
1:A:555:ILE:HG23	1:A:656:VAL:CG1	2.46	0.45
1:A:681:PHE:CD1	1:A:682:PRO:HD2	2.51	0.45
1:A:555:ILE:O	1:A:555:ILE:HG22	2.16	0.45
1:A:713:ALA:HB2	7:A:1883:IOD:I	2.86	0.45
1:A:68:VAL:HG12	1:A:69:GLY:H	1.82	0.45
1:A:201:MET:CE	1:A:356:VAL:CG2	2.93	0.44
1:A:344:GLN:HB3	7:A:1878:IOD:I	2.87	0.44
1:A:176:SER:HG	1:A:320:PHE:HZ	1.65	0.44
1:A:495:LYS:HG2	7:A:1882:IOD:I	2.88	0.44
1:A:479:LYS:O	1:A:859:GLU:HG3	2.18	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:600:ARG:HD2	1:A:611:ILE:HD11	1.99	0.43
1:A:92:LEU:HA	1:A:95:LYS:HE2	2.00	0.43
1:A:376:LEU:HB2	1:A:379:VAL:HG23	2.01	0.43
1:A:372:LEU:HD12	1:A:384:LEU:HD22	2.01	0.42
1:A:134:GLN:CG	7:A:1878:IOD:I	3.37	0.42
1:A:704:MET:HA	1:A:797:HIS:NE2	2.34	0.42
1:A:795:LEU:HD21	1:A:819:MET:HG2	2.01	0.41
1:A:635:ILE:HD13	1:A:720:VAL:HG11	2.02	0.41
1:A:843:ARG:NH1	7:A:1872:IOD:I	3.24	0.41
1:A:63:PHE:CE2	1:A:283:ARG:HD2	2.56	0.40
1:A:656:VAL:CG1	1:A:656:VAL:O	2.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	776/827 (94%)	748 (96%)	27 (4%)	1 (0%)	51 45

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	555	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	709/750 (94%)	698 (98%)	11 (2%)	62 59

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	192	ARG
1	A	278	SER
1	A	389	LEU
1	A	536	THR
1	A	545	ASP
1	A	569	ASP
1	A	606	ARG
1	A	640	GLU
1	A	665	ASN
1	A	674	GLN
1	A	843	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	674	GLN
1	A	716	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

3 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	B	1	1,2	14,14,15	0.77	0	17,19,21	1.46	3 (17%)
2	NAG	B	2	2	14,14,15	0.82	1 (7%)	17,19,21	1.61	3 (17%)
2	BMA	B	3	2	11,11,12	0.85	0	15,15,17	2.28	6 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	0/6/23/26	0/1/1/1
2	BMA	B	3	2	-	0/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2	NAG	O5-C1	-2.01	1.40	1.43

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	3	BMA	O5-C1-C2	-4.75	103.44	110.77
2	B	3	BMA	C1-O5-C5	3.68	117.18	112.19
2	B	2	NAG	C8-C7-N2	3.55	122.11	116.10
2	B	2	NAG	C4-C3-C2	3.23	115.75	111.02
2	B	3	BMA	O5-C5-C4	3.17	118.55	110.83
2	B	3	BMA	C3-C4-C5	3.05	115.68	110.24
2	B	2	NAG	O7-C7-C8	-2.99	116.50	122.06
2	B	1	NAG	C1-O5-C5	2.85	116.06	112.19
2	B	1	NAG	O4-C4-C5	-2.62	102.80	109.30
2	B	3	BMA	O2-C2-C3	2.44	115.03	110.14
2	B	1	NAG	O5-C1-C2	-2.30	107.65	111.29
2	B	3	BMA	O5-C5-C6	2.01	110.35	107.20


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 36 ligands modelled in this entry, 31 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	PO4	A	1870	4	4,4,4	0.98	0	6,6,6	2.29	1 (16%)
3	SCN	A	1866	-	1,2,2	0.96	0	0,1,1	0.00	-
3	SCN	A	1864	-	1,2,2	0.72	0	0,1,1	0.00	-
3	SCN	A	1865	-	1,2,2	0.59	0	0,1,1	0.00	-
3	SCN	A	1863	-	1,2,2	0.72	0	0,1,1	0.00	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
6	A	1870	PO4	O4-P-O2	4.75	123.21	107.97

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	783/827 (94%)	0.16	39 (4%) 28 31	15, 33, 62, 87	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	397	ILE	8.5
1	A	70	PRO	6.5
1	A	569	ASP	6.2
1	A	161	VAL	6.1
1	A	69	GLY	5.1
1	A	68	VAL	4.7
1	A	563	ASP	4.6
1	A	578	LEU	4.5
1	A	67	GLU	4.5
1	A	396	SER	4.3
1	A	538	THR	4.1
1	A	590	SER	3.6
1	A	468	CYS	3.6
1	A	667	LEU	3.0
1	A	662	PHE	2.9
1	A	366	CYS	2.9
1	A	567	THR	2.8
1	A	579	GLU	2.7
1	A	673	LYS	2.7
1	A	646	GLU	2.6
1	A	320	PHE	2.6
1	A	670	LYS	2.6
1	A	593	GLU	2.6
1	A	72	ASP	2.5
1	A	556	MET	2.5
1	A	580	GLU	2.4
1	A	698	VAL	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	59	LYS	2.4
1	A	60	GLY	2.3
1	A	182	SER	2.3
1	A	559	GLN	2.3
1	A	568	CYS	2.2
1	A	66	GLN	2.2
1	A	183	LYS	2.2
1	A	246	ARG	2.1
1	A	82	TYR	2.1
1	A	656	VAL	2.1
1	A	87	HIS	2.1
1	A	577	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	BMA	B	3	11/12	0.67	0.38	58,61,63,65	0
2	NAG	B	2	14/15	0.92	0.18	34,40,48,49	0
2	NAG	B	1	14/15	0.93	0.09	20,23,29,30	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	SCN	A	1865	3/3	0.60	0.18	62,62,62,64	0
7	IOD	A	1878	1/1	0.65	2.05	149,149,149,149	1
3	SCN	A	1866	3/3	0.74	0.21	73,73,73,74	0
3	SCN	A	1863	3/3	0.78	0.23	50,50,50,51	0
3	SCN	A	1864	3/3	0.84	0.19	69,69,69,70	0
7	IOD	A	1892	1/1	0.86	0.14	59,59,59,59	1
7	IOD	A	1891	1/1	0.92	0.08	52,52,52,52	1
7	IOD	A	1890	1/1	0.92	0.13	74,74,74,74	1
7	IOD	A	1883	1/1	0.92	0.13	42,42,42,42	1
7	IOD	A	1897	1/1	0.93	0.11	55,55,55,55	1
7	IOD	A	1896	1/1	0.93	0.08	63,63,63,63	1
7	IOD	A	1885	1/1	0.94	0.19	61,61,61,61	1
7	IOD	A	1882	1/1	0.94	0.11	47,47,47,47	1
7	IOD	A	1889	1/1	0.96	0.15	53,53,53,53	1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
7	IOD	A	1886	1/1	0.97	0.08	66,66,66,66	1
7	IOD	A	1895	1/1	0.97	0.20	63,63,63,63	1
7	IOD	A	1884	1/1	0.97	0.03	57,57,57,57	1
7	IOD	A	1880	1/1	0.97	0.04	45,45,45,45	1
4	ZN	A	1868	1/1	0.97	0.08	44,44,44,44	0
7	IOD	A	1898	1/1	0.97	0.04	55,55,55,55	1
7	IOD	A	1876	1/1	0.98	0.05	40,40,40,40	1
7	IOD	A	1879	1/1	0.98	0.04	42,42,42,42	1
7	IOD	A	1888	1/1	0.98	0.07	44,44,44,44	1
7	IOD	A	1877	1/1	0.98	0.03	45,45,45,45	1
7	IOD	A	1887	1/1	0.98	0.05	43,43,43,43	1
7	IOD	A	1893	1/1	0.98	0.02	59,59,59,59	1
6	PO4	A	1870	5/5	0.98	0.07	30,30,37,37	0
7	IOD	A	1874	1/1	0.98	0.05	43,43,43,43	1
4	ZN	A	1867	1/1	0.99	0.06	33,33,33,33	0
7	IOD	A	1881	1/1	0.99	0.03	38,38,38,38	1
7	IOD	A	1894	1/1	0.99	0.09	58,58,58,58	1
7	IOD	A	1875	1/1	0.99	0.03	42,42,42,42	1
5	CA	A	1869	1/1	1.00	0.09	25,25,25,25	0
7	IOD	A	1872	1/1	1.00	0.04	33,33,33,33	1
7	IOD	A	1871	1/1	1.00	0.04	30,30,30,30	0
7	IOD	A	1873	1/1	1.00	0.03	43,43,43,43	1

6.5 Other polymers ⓘ

There are no such residues in this entry.