



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 10, 2020 – 01:11 AM BST

PDB ID : 4XUQ
Title : Structure of the CBM22-2 xylan-binding domain from *Paenibacillus barcinonensis* Xyn10C in complex with xylotriase
Authors : Sainz-Polo, M.A.; Sanz-Aparicio, J.
Deposited on : 2015-01-26
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	181	<div> <div>2%</div> <div> <div></div> <div>82%</div> <div>7%</div> <div>11%</div> </div> </div>
1	B	181	<div> <div></div> <div> <div>87%</div> <div>11%</div> </div> </div>
1	C	181	<div> <div>%</div> <div> <div></div> <div>87%</div> <div>11%</div> </div> </div>
2	D	2	<div> <div></div> <div> <div>50%</div> <div>50%</div> </div> </div>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3942 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Endo-1,4-beta-xylanase C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	161	Total	C	N	O	S	0	0	0
			1263	796	211	253	3			
1	B	161	Total	C	N	O	S	0	0	0
			1263	796	211	253	3			
1	C	161	Total	C	N	O	S	0	0	0
			1263	796	211	253	3			

There are 3 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	177	ASP	TYR	conflict	UNP O69230
B	177	ASP	TYR	conflict	UNP O69230
C	177	ASP	TYR	conflict	UNP O69230

- Molecule 2 is an oligosaccharide called beta-D-xylopyranose-(1-4)-beta-D-xylopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	D	2	Total	C	O	0	0	0
			19	10	9			

- Molecule 3 is beta-D-xylopyranose (three-letter code: XYP) (formula: C₅H₁₀O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			9	5	4		
3	B	1	Total	C	O	0	0
			9	5	4		
3	B	1	Total	C	O	0	0
			10	5	5		
3	C	1	Total	C	O	0	0
			9	5	4		
3	C	1	Total	C	O	0	0
			10	5	5		

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Ca	0	0
			1	1		
4	A	1	Total	Ca	0	0
			1	1		
4	C	1	Total	Ca	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	35	Total	O	0	0
			35	35		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	24	Total	O	0	0
			24	24		
5	C	25	Total	O	0	0
			25	25		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Endo-1,4-beta-xylanase C



- Molecule 1: Endo-1,4-beta-xylanase C



- Molecule 1: Endo-1,4-beta-xylanase C



- Molecule 2: beta-D-xylopyranose-(1-4)-beta-D-xylopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, α , β , γ	92.45Å 92.45Å 48.42Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.42 – 1.95 46.23 – 1.95	Depositor EDS
% Data completeness (in resolution range)	100.0 (48.42-1.95) 99.9 (46.23-1.95)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.03	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.05 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.8.0103	Depositor
R, R_{free}	0.178 , 0.213 0.186 , 0.220	Depositor DCC
R_{free} test set	1715 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	18.2	Xtriage
Anisotropy	0.107	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 37.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.028 for -h,-k,l 0.039 for h,-h-k,-l 0.056 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3942	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: XYP, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.85	0/1291	0.83	0/1753
1	B	0.84	0/1291	0.78	0/1753
1	C	0.83	0/1291	0.78	0/1753
All	All	0.84	0/3873	0.80	0/5259

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1263	0	1216	13	0
1	B	1263	0	1216	3	0
1	C	1263	0	1216	2	0
2	D	19	0	0	4	0
3	A	9	0	0	4	0
3	B	19	0	0	3	0
3	C	19	0	0	3	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	35	0	0	0	0
5	B	24	0	0	0	0
5	C	25	0	0	0	0
All	All	3942	0	3648	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:1001:XYP:C1	2:D:2:XYP:O4	1.63	1.45
3:C:401:XYP:C1	3:C:402:XYP:O4	1.66	1.41
3:B:401:XYP:C1	3:B:402:XYP:O4	1.69	1.39
3:A:1001:XYP:C1	2:D:2:XYP:C4	2.51	0.87
3:C:401:XYP:C1	3:C:402:XYP:C4	2.57	0.83
3:B:401:XYP:C1	3:B:402:XYP:C4	2.59	0.81
3:A:1001:XYP:C2	2:D:2:XYP:O4	2.44	0.66
3:C:401:XYP:C2	3:C:402:XYP:O4	2.45	0.59
1:A:308:TRP:O	1:A:309:LYS:HG2	2.04	0.57
1:A:310:GLN:O	1:A:314:VAL:HG23	2.04	0.57
3:B:401:XYP:C2	3:B:402:XYP:O4	2.50	0.57
1:A:251:ILE:HG22	1:A:276:LEU:CD1	2.39	0.52
1:A:257:LYS:HG2	1:A:300:LEU:CD1	2.40	0.51
1:A:251:ILE:CG2	1:A:284:ILE:HD11	2.42	0.48
1:B:236:ILE:CG2	1:B:302:MET:HE2	2.43	0.48
1:C:202:HIS:HD2	1:C:326:ASP:OD2	1.97	0.47
3:A:1001:XYP:O5	2:D:2:XYP:O4	2.25	0.47
1:A:251:ILE:HG22	1:A:276:LEU:HD13	1.97	0.47
1:A:309:LYS:HD3	1:A:309:LYS:HA	1.67	0.47
1:B:236:ILE:HG21	1:B:302:MET:HE2	1.96	0.47
1:A:202:HIS:HD2	1:A:326:ASP:OD2	1.98	0.46
1:B:302:MET:HG2	1:B:303:TYR:N	2.33	0.44
1:A:316:GLU:HG3	1:C:274:THR:HB	2.02	0.42
1:A:276:LEU:HD12	1:A:276:LEU:N	2.35	0.41
1:A:251:ILE:HG21	1:A:284:ILE:HD11	2.03	0.41
1:A:257:LYS:HG2	1:A:300:LEU:HD13	2.02	0.41
1:A:257:LYS:HG2	1:A:300:LEU:HD12	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	159/181 (88%)	157 (99%)	2 (1%)	0	100	100
1	B	159/181 (88%)	158 (99%)	1 (1%)	0	100	100
1	C	159/181 (88%)	157 (99%)	2 (1%)	0	100	100
All	All	477/543 (88%)	472 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	137/150 (91%)	136 (99%)	1 (1%)	84	82
1	B	137/150 (91%)	136 (99%)	1 (1%)	84	82
1	C	137/150 (91%)	136 (99%)	1 (1%)	84	82
All	All	411/450 (91%)	408 (99%)	3 (1%)	84	82

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	301	ARG
1	B	301	ARG
1	C	301	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	202	HIS
1	A	203	ASN
1	A	229	GLN
1	A	277	ASN
1	A	280	GLN
1	B	203	ASN
1	B	265	ASN
1	B	277	ASN
1	B	280	GLN
1	C	202	HIS
1	C	203	ASN
1	C	229	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	XYP	D	1	2	10,10,10	1.51	3 (30%)	14,14,14	1.03	0
2	XYP	D	2	2	9,9,10	1.52	1 (11%)	10,12,14	1.75	2 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	XYP	D	1	2	-	-	0/1/1/1
2	XYP	D	2	2	-	-	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	2	XYP	O5-C1	3.44	1.49	1.42
2	D	1	XYP	O5-C1	2.51	1.46	1.43
2	D	1	XYP	O5-C5	-2.29	1.40	1.43
2	D	1	XYP	C5-C4	-2.28	1.47	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	2	XYP	O4-C4-C3	-3.71	102.71	110.14
2	D	2	XYP	C5-C4-C3	3.52	114.00	109.67

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	2	XYP	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 3 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	XYP	B	402	-	10,10,10	0.95	1 (10%)	14,14,14	1.09	2 (14%)
3	XYP	B	401	-	9,9,10	0.72	0	10,12,14	1.53	3 (30%)
3	XYP	C	401	-	9,9,10	0.63	0	10,12,14	2.07	5 (50%)
3	XYP	A	1001	-	9,9,10	0.65	0	10,12,14	2.42	4 (40%)
3	XYP	C	402	-	10,10,10	1.24	1 (10%)	14,14,14	1.09	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	XYP	B	402	-	-	-	0/1/1/1
3	XYP	B	401	-	-	-	0/1/1/1
3	XYP	C	401	-	-	-	0/1/1/1
3	XYP	A	1001	-	-	-	0/1/1/1
3	XYP	C	402	-	-	-	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	402	XYP	C5-C4	-2.21	1.47	1.52
3	B	402	XYP	O5-C5	-2.17	1.40	1.43

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1001	XYP	C1-C2-C3	-5.39	103.04	109.67
3	A	1001	XYP	C5-O5-C1	-3.36	106.36	111.52
3	C	401	XYP	C1-C2-C3	-3.34	105.56	109.67
3	B	401	XYP	C1-C2-C3	-3.01	105.96	109.67
3	A	1001	XYP	O2-C2-C3	2.92	115.98	110.14
3	C	401	XYP	O3-C3-C2	-2.91	104.42	109.99
3	C	401	XYP	C5-C4-C3	2.80	113.11	109.67
3	B	402	XYP	O4-C4-C5	-2.57	103.90	109.15
3	C	401	XYP	O3-C3-C4	2.48	114.74	109.99
3	C	401	XYP	O2-C2-C3	2.44	115.03	110.14
3	B	401	XYP	O2-C2-C3	2.30	114.74	110.14
3	A	1001	XYP	O4-C4-C3	-2.06	106.00	110.14
3	B	401	XYP	C5-C4-C3	2.04	112.17	109.67
3	B	402	XYP	C4-C3-C2	-2.03	107.38	110.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	402	XYP	3	0
3	B	401	XYP	3	0
3	C	401	XYP	3	0
3	A	1001	XYP	4	0
3	C	402	XYP	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	161/181 (88%)	-0.14	3 (1%) 66 74	10, 17, 33, 81	0
1	B	161/181 (88%)	-0.27	0 100 100	12, 20, 36, 52	0
1	C	161/181 (88%)	-0.27	1 (0%) 89 93	12, 19, 37, 56	0
All	All	483/543 (88%)	-0.23	4 (0%) 86 90	10, 19, 37, 81	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	311	SER	8.5
1	A	310	GLN	6.8
1	A	312	SER	5.3
1	C	310	GLN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	XYP	D	1	10/10	0.92	0.10	21,25,27,27	0
2	XYP	D	2	9/10	0.94	0.09	15,17,19,20	0

The following is a graphical depiction of the model fit to experimental electron density for oligosac-

charide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	XYP	B	402	10/10	0.86	0.17	36,40,44,46	0
3	XYP	B	401	9/10	0.88	0.13	33,36,41,42	0
3	XYP	A	1001	9/10	0.88	0.14	23,33,41,41	0
3	XYP	C	401	9/10	0.91	0.13	24,30,37,44	0
3	XYP	C	402	10/10	0.92	0.14	30,33,39,40	0
4	CA	C	403	1/1	0.97	0.05	25,25,25,25	0
4	CA	A	1004	1/1	0.99	0.08	18,18,18,18	0
4	CA	B	403	1/1	0.99	0.08	17,17,17,17	0

6.5 Other polymers ⓘ

There are no such residues in this entry.