



wwPDB X-ray Structure Validation Summary Report ⓘ

May 21, 2020 – 09:17 pm BST

PDB ID : 5XV4
Title : Crystal structure of ATG101-ATG13HORMA
Authors : Kim, B.-W.; Song, H.K.
Deposited on : 2017-06-26
Resolution : 2.95 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3104 (3.00-2.92)
Clashscore	141614	3462 (3.00-2.92)
Ramachandran outliers	138981	3340 (3.00-2.92)
Sidechain outliers	138945	3343 (3.00-2.92)
RSRZ outliers	127900	2986 (3.00-2.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	190	<div> <div>5%</div> <div> <div></div> <div>83%</div> <div>11%</div> <div>• •</div> </div> </div>
1	C	190	<div> <div>6%</div> <div> <div></div> <div>78%</div> <div>17%</div> <div>• •</div> </div> </div>
1	E	190	<div> <div>4%</div> <div> <div></div> <div>76%</div> <div>19%</div> <div>• •</div> </div> </div>
1	G	190	<div> <div>3%</div> <div> <div></div> <div>77%</div> <div>17%</div> <div>• •</div> </div> </div>
1	I	190	<div> <div>9%</div> <div> <div></div> <div>76%</div> <div>18%</div> <div>• •</div> </div> </div>
1	K	190	<div> <div>6%</div> <div> <div></div> <div>78%</div> <div>17%</div> <div>• •</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	M	190	
1	O	190	
2	B	218	
2	D	218	
2	F	218	
2	H	218	
2	J	218	
2	L	218	
2	N	218	
2	P	218	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 23827 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Autophagy-related protein 13.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	182	Total	C	N	O	S	0	0	0
			1356	865	231	252	8			
1	C	183	Total	C	N	O	S	0	0	0
			1358	865	233	253	7			
1	E	182	Total	C	N	O	S	0	0	0
			1357	868	228	253	8			
1	G	182	Total	C	N	O	S	0	0	0
			1357	866	232	252	7			
1	I	182	Total	C	N	O	S	0	0	0
			1363	871	231	253	8			
1	K	182	Total	C	N	O	S	0	0	0
			1365	872	233	252	8			
1	M	183	Total	C	N	O	S	0	0	0
			1375	878	233	256	8			
1	O	182	Total	C	N	O	S	0	0	0
			1354	863	231	252	8			

- Molecule 2 is a protein called Autophagy-related protein 101.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	207	Total	C	N	O	S	0	0	0
			1613	1020	275	309	9			
2	D	207	Total	C	N	O	S	0	0	0
			1610	1018	275	309	8			
2	F	207	Total	C	N	O	S	0	0	0
			1607	1017	272	309	9			
2	H	210	Total	C	N	O	S	0	0	0
			1627	1028	277	312	10			
2	J	207	Total	C	N	O	S	0	0	0
			1613	1020	275	309	9			
2	L	209	Total	C	N	O	S	0	0	0
			1626	1028	277	311	10			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	N	212	Total	C	N	O	S	0	0	0
			1636	1036	279	312	9			
2	P	207	Total	C	N	O	S	0	0	0
			1610	1018	275	309	8			


There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	40	ALA	LYS	engineered mutation	UNP Q9BSB4
B	41	ALA	LYS	engineered mutation	UNP Q9BSB4
B	42	ALA	GLU	engineered mutation	UNP Q9BSB4
D	40	ALA	LYS	engineered mutation	UNP Q9BSB4
D	41	ALA	LYS	engineered mutation	UNP Q9BSB4
D	42	ALA	GLU	engineered mutation	UNP Q9BSB4
F	40	ALA	LYS	engineered mutation	UNP Q9BSB4
F	41	ALA	LYS	engineered mutation	UNP Q9BSB4
F	42	ALA	GLU	engineered mutation	UNP Q9BSB4
H	40	ALA	LYS	engineered mutation	UNP Q9BSB4
H	41	ALA	LYS	engineered mutation	UNP Q9BSB4
H	42	ALA	GLU	engineered mutation	UNP Q9BSB4
J	40	ALA	LYS	engineered mutation	UNP Q9BSB4
J	41	ALA	LYS	engineered mutation	UNP Q9BSB4
J	42	ALA	GLU	engineered mutation	UNP Q9BSB4
L	40	ALA	LYS	engineered mutation	UNP Q9BSB4
L	41	ALA	LYS	engineered mutation	UNP Q9BSB4
L	42	ALA	GLU	engineered mutation	UNP Q9BSB4
N	40	ALA	LYS	engineered mutation	UNP Q9BSB4
N	41	ALA	LYS	engineered mutation	UNP Q9BSB4
N	42	ALA	GLU	engineered mutation	UNP Q9BSB4
P	40	ALA	LYS	engineered mutation	UNP Q9BSB4
P	41	ALA	LYS	engineered mutation	UNP Q9BSB4
P	42	ALA	GLU	engineered mutation	UNP Q9BSB4


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Autophagy-related protein 13




- Molecule 1: Autophagy-related protein 13




- Molecule 1: Autophagy-related protein 13





- Molecule 1: Autophagy-related protein 13









• Molecule 1: Autophagy-related protein 13





• Molecule 1: Autophagy-related protein 13




• Molecule 1: Autophagy-related protein 13




• Molecule 1: Autophagy-related protein 13




- Molecule 2: Autophagy-related protein 101




- Molecule 2: Autophagy-related protein 101




- Molecule 2: Autophagy-related protein 101



- Molecule 2: Autophagy-related protein 101






- Molecule 2: Autophagy-related protein 101









• Molecule 2: Autophagy-related protein 101



• Molecule 2: Autophagy-related protein 101



• Molecule 2: Autophagy-related protein 101



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	101.29Å 125.36Å 174.62Å 90.00° 95.69° 90.00°	Depositor
Resolution (Å)	35.78 – 2.95 35.78 – 2.95	Depositor EDS
% Data completeness (in resolution range)	95.3 (35.78-2.95) 95.3 (35.78-2.95)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.17 (at 2.95Å)	Xtriage
Refinement program	PHENIX 1.10.1_2155	Depositor
R, R_{free}	0.286 , 0.313 0.286 , 0.313	Depositor DCC
R_{free} test set	1959 reflections (2.24%)	wwPDB-VP
Wilson B-factor (Å ²)	66.1	Xtriage
Anisotropy	0.554	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 37.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	23827	wwPDB-VP
Average B, all atoms (Å ²)	76.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 66.85 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.6577e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.27	0/1376	0.48	0/1867
1	C	0.26	0/1378	0.46	0/1870
1	E	0.26	0/1378	0.46	0/1871
1	G	0.26	0/1377	0.47	0/1868
1	I	0.27	0/1384	0.48	0/1878
1	K	0.27	0/1386	0.46	0/1880
1	M	0.27	0/1396	0.47	0/1893
1	O	0.27	0/1374	0.47	0/1863
2	B	0.28	0/1643	0.46	0/2228
2	D	0.27	0/1640	0.46	0/2225
2	F	0.28	0/1637	0.46	0/2221
2	H	0.26	0/1657	0.46	0/2248
2	J	0.27	0/1643	0.45	0/2228
2	L	0.27	0/1656	0.46	0/2245
2	N	0.27	0/1666	0.46	0/2262
2	P	0.27	0/1640	0.47	0/2225
All	All	0.27	0/24231	0.46	0/32872

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1356	0	1336	13	0
1	C	1358	0	1331	17	0
1	E	1357	0	1332	21	0
1	G	1357	0	1340	19	0
1	I	1363	0	1343	19	0
1	K	1365	0	1348	16	0
1	M	1375	0	1358	17	0
1	O	1354	0	1335	15	0
2	B	1613	0	1563	24	0
2	D	1610	0	1556	16	0
2	F	1607	0	1552	23	0
2	H	1627	0	1565	18	0
2	J	1613	0	1563	21	0
2	L	1626	0	1574	15	0
2	N	1636	0	1576	17	0
2	P	1610	0	1556	25	0
All	All	23827	0	23228	278	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 278 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:J:131:LEU:HD11	2:J:136:GLU:HB3	1.60	0.84
1:O:104:ASN:ND2	1:O:182:THR:OG1	2.16	0.78
2:P:106:LYS:HB3	2:P:115:GLU:HG3	1.68	0.75
2:D:106:LYS:HB3	2:D:115:GLU:HG3	1.69	0.74
2:N:4:ARG:NH2	2:N:120:GLU:OE1	2.21	0.74

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	178/190 (94%)	173 (97%)	5 (3%)	0	100	100
1	C	179/190 (94%)	173 (97%)	6 (3%)	0	100	100
1	E	178/190 (94%)	172 (97%)	6 (3%)	0	100	100
1	G	178/190 (94%)	172 (97%)	6 (3%)	0	100	100
1	I	178/190 (94%)	171 (96%)	7 (4%)	0	100	100
1	K	178/190 (94%)	173 (97%)	5 (3%)	0	100	100
1	M	179/190 (94%)	174 (97%)	5 (3%)	0	100	100
1	O	178/190 (94%)	173 (97%)	5 (3%)	0	100	100
2	B	205/218 (94%)	197 (96%)	8 (4%)	0	100	100
2	D	205/218 (94%)	198 (97%)	7 (3%)	0	100	100
2	F	205/218 (94%)	198 (97%)	7 (3%)	0	100	100
2	H	208/218 (95%)	202 (97%)	6 (3%)	0	100	100
2	J	205/218 (94%)	197 (96%)	8 (4%)	0	100	100
2	L	207/218 (95%)	199 (96%)	8 (4%)	0	100	100
2	N	210/218 (96%)	203 (97%)	7 (3%)	0	100	100
2	P	205/218 (94%)	199 (97%)	6 (3%)	0	100	100
All	All	3076/3264 (94%)	2974 (97%)	102 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	140/167 (84%)	129 (92%)	11 (8%)	12	37
1	C	139/167 (83%)	131 (94%)	8 (6%)	20	51
1	E	140/167 (84%)	134 (96%)	6 (4%)	29	62
1	G	140/167 (84%)	132 (94%)	8 (6%)	20	52
1	I	141/167 (84%)	131 (93%)	10 (7%)	14	43

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	K	141/167 (84%)	135 (96%)	6 (4%)	29	62
1	M	143/167 (86%)	136 (95%)	7 (5%)	25	58
1	O	140/167 (84%)	135 (96%)	5 (4%)	35	67
2	B	173/194 (89%)	164 (95%)	9 (5%)	23	56
2	D	172/194 (89%)	168 (98%)	4 (2%)	50	78
2	F	172/194 (89%)	168 (98%)	4 (2%)	50	78
2	H	173/194 (89%)	165 (95%)	8 (5%)	27	60
2	J	173/194 (89%)	165 (95%)	8 (5%)	27	60
2	L	174/194 (90%)	163 (94%)	11 (6%)	18	48
2	N	173/194 (89%)	167 (96%)	6 (4%)	36	68
2	P	172/194 (89%)	167 (97%)	5 (3%)	42	73
All	All	2506/2888 (87%)	2390 (95%)	116 (5%)	27	60

5 of 116 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	H	25	LEU
1	I	153	ARG
1	O	9	ASP
2	H	124	VAL
1	I	10	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	E	26	GLN
1	M	26	GLN
1	M	144	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å²)	Q<0.9
1	A	182/190 (95%)	0.32	9 (4%)	29	18	49, 75, 105, 141	0
1	C	183/190 (96%)	0.31	11 (6%)	21	13	48, 85, 119, 145	0
1	E	182/190 (95%)	0.26	8 (4%)	34	21	43, 71, 101, 129	0
1	G	182/190 (95%)	0.32	6 (3%)	46	30	56, 86, 121, 147	0
1	I	182/190 (95%)	0.52	18 (9%)	7	4	50, 77, 109, 146	0
1	K	182/190 (95%)	0.51	12 (6%)	18	10	47, 72, 103, 115	0
1	M	183/190 (96%)	0.35	12 (6%)	18	10	42, 69, 102, 123	0
1	O	182/190 (95%)	0.52	19 (10%)	6	4	48, 79, 109, 139	0
2	B	207/218 (94%)	0.33	11 (5%)	26	16	53, 78, 116, 136	0
2	D	207/218 (94%)	0.13	5 (2%)	59	42	51, 76, 107, 117	0
2	F	207/218 (94%)	0.33	12 (5%)	23	14	47, 74, 108, 124	0
2	H	210/218 (96%)	0.17	10 (4%)	30	19	52, 76, 109, 137	0
2	J	207/218 (94%)	0.44	13 (6%)	20	11	50, 73, 109, 123	0
2	L	209/218 (95%)	0.40	13 (6%)	20	12	45, 71, 99, 135	0
2	N	212/218 (97%)	0.44	16 (7%)	14	8	45, 70, 106, 135	0
2	P	207/218 (94%)	0.41	16 (7%)	13	7	48, 74, 112, 123	0
All	All	3124/3264 (95%)	0.36	191 (6%)	21	12	42, 75, 110, 147	0

The worst 5 of 191 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	207	THR	7.8
2	J	207	THR	6.8
2	J	162	HIS	6.7
1	A	90	GLU	6.6
2	N	210	ARG	6.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.