



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 16, 2020 – 10:41 pm BST

PDB ID : 1XXV  
Title : Yersinia YopH (residues 163-468) binds phosphonodifluoromethyl-Phe containing hexapeptide at two sites  
Authors : Ivanov, M.I.; Stuckey, J.A.; Schubert, H.L.; Saper, M.A.; Bliska, J.B.  
Deposited on : 2004-11-08  
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	306	
1	B	306	
2	C	8	
2	D	8	
2	E	8	
2	F	8	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4831 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein-tyrosine phosphatase yopH.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	282	Total	C	N	O	S	0	0	0
			2166	1323	402	425	16			
1	B	282	Total	C	N	O	S	0	0	0
			2166	1323	402	425	16			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	235	ARG	CYS	ENGINEERED MUTATION	UNP P15273
B	235	ARG	CYS	ENGINEERED MUTATION	UNP P15273

- Molecule 2 is a protein called Epidermal growth factor receptor derived peptide.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	C	5	Total 44	C 25	F 2	N 5	O 11	P 1	0	0	1
2	D	7	Total 57	C 32	F 2	N 7	O 15	P 1	0	0	1
2	E	5	Total 44	C 25	F 2	N 5	O 11	P 1	4	0	1
2	F	7	Total 57	C 32	F 2	N 7	O 15	P 1	0	0	1

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	154	Total	O	0	0
			154	154		
3	B	134	Total	O	0	0
			134	134		
3	C	1	Total	O	0	0
			1	1		

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	3	Total 3	O 3	0	0
3	E	1	Total 1	O 1	0	0
3	F	4	Total 4	O 4	0	0

### 3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: Protein-tyrosine phosphatase yopH



- Molecule 1: Protein-tyrosine phosphatase yopH




- Molecule 2: Epidermal growth factor receptor derived peptide




- Molecule 2: Epidermal growth factor receptor derived peptide






- Molecule 2: Epidermal growth factor receptor derived peptide



- Molecule 2: Epidermal growth factor receptor derived peptide



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	54.13Å 47.17Å 71.82Å 104.45° 115.05° 89.97°	Depositor
Resolution (Å)	50.00 – 2.50	Depositor
% Data completeness (in resolution range)	85.5 (50.00-2.50)	Depositor
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	CNS	Depositor
R, $R_{free}$	0.178 , 0.229	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4831	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FTY, NH2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.37	0/2193	0.59	0/2962
1	B	0.37	0/2193	0.59	0/2962
2	C	0.53	0/23	0.40	0/28
2	D	0.41	0/36	0.54	0/46
2	E	0.52	0/23	0.47	0/28
2	F	0.39	0/36	0.51	0/46
All	All	0.37	0/4504	0.58	0/6072

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2166	0	2167	36	0
1	B	2166	0	2167	40	0
2	C	44	0	27	3	0
2	D	57	0	39	2	0
2	E	44	0	27	1	0
2	F	57	0	39	1	0
3	A	154	0	0	4	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	134	0	0	6	0
3	C	1	0	0	0	0
3	D	3	0	0	0	0
3	E	1	0	0	0	0
3	F	4	0	0	0	0
All	All	4831	0	4466	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 79 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:241:ARG:HD3	1:A:244:LEU:HD12	1.72	0.70
1:B:241:ARG:HD3	1:B:244:LEU:HD12	1.72	0.70
1:A:272:ARG:HG3	1:A:311:ILE:HD11	1.75	0.68
1:A:342:LYS:HD3	2:D:5:FTY:HE1	1.74	0.68
1:B:363:GLU:HB2	3:B:821:HOH:O	1.94	0.67

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	280/306 (92%)	263 (94%)	15 (5%)	2 (1%)	22	39
1	B	280/306 (92%)	266 (95%)	12 (4%)	2 (1%)	22	39
2	C	2/8 (25%)	1 (50%)	1 (50%)	0	100	100
2	D	4/8 (50%)	4 (100%)	0	0	100	100
2	E	2/8 (25%)	2 (100%)	0	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	F	4/8 (50%)	4 (100%)	0	0	100	100
All	All	572/644 (89%)	540 (94%)	28 (5%)	4 (1%)	22	39

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	442	GLY
1	A	445	VAL
1	B	445	VAL
1	B	442	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	238/255 (93%)	234 (98%)	4 (2%)	60	82
1	B	238/255 (93%)	234 (98%)	4 (2%)	60	82
2	C	3/4 (75%)	2 (67%)	1 (33%)	0	0
2	D	4/4 (100%)	4 (100%)	0	100	100
2	E	3/4 (75%)	3 (100%)	0	100	100
2	F	4/4 (100%)	4 (100%)	0	100	100
All	All	490/526 (93%)	481 (98%)	9 (2%)	59	81

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	237	GLN
2	C	106	LEU
1	B	319	GLN
1	A	319	GLN
1	B	251	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	420	ASN
1	B	435	GLN
1	B	294	GLN
1	A	294	GLN
1	B	420	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	FTY	C	105	2	14,18,19	2.32	3 (21%)	19,27,29	2.50	3 (15%)
2	FTY	D	5	2	14,18,19	1.96	3 (21%)	19,27,29	2.28	4 (21%)
2	FTY	E	105	2	14,18,19	2.18	3 (21%)	19,27,29	2.55	3 (15%)
2	FTY	F	5	2	14,18,19	2.02	3 (21%)	19,27,29	2.24	3 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FTY	C	105	2	-	0/15/21/23	0/1/1/1
2	FTY	D	5	2	-	5/15/21/23	0/1/1/1
2	FTY	E	105	2	-	0/15/21/23	0/1/1/1
2	FTY	F	5	2	-	6/15/21/23	0/1/1/1

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	105	FTY	C1-CZ	7.27	1.56	1.50
2	E	105	FTY	C1-CZ	6.59	1.56	1.50
2	F	5	FTY	C1-CZ	5.75	1.55	1.50
2	D	5	FTY	C1-CZ	5.57	1.55	1.50
2	E	105	FTY	P-O2P	-3.64	1.48	1.54

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	105	FTY	F2-C1-F1	-8.16	96.42	106.73
2	C	105	FTY	F2-C1-F1	-8.06	96.56	106.73
2	D	5	FTY	F2-C1-F1	-7.22	97.62	106.73
2	F	5	FTY	F2-C1-F1	-6.71	98.26	106.73
2	E	105	FTY	F2-C1-CZ	-5.03	104.01	110.49

There are no chirality outliers.

5 of 11 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	5	FTY	F1-C1-CZ-CE1
2	D	5	FTY	F1-C1-CZ-CE2
2	D	5	FTY	F2-C1-P-O1P
2	F	5	FTY	O-C-CA-CB
2	F	5	FTY	F1-C1-CZ-CE1

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	105	FTY	1	0
2	D	5	FTY	1	0
2	E	105	FTY	1	0

## 5.5 Carbohydrates

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

There are no ligands in this entry.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates ⓘ

EDS was not executed - this section is therefore empty.

### 6.4 Ligands ⓘ

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers ⓘ

EDS was not executed - this section is therefore empty.