



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 7, 2020 – 07:09 PM BST

PDB ID : 4XX6
Title : Crystal structure of a glycosylated endo-beta-1,4-xylanase (glycoside hydrolase family 10/GH10) enzyme from *Gloeophyllum trabeum*
Authors : Stogios, P.J.; Nocek, B.; Xu, X.; Cui, H.; Lowden, M.; Savchenko, A.
Deposited on : 2015-01-29
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	321	<div> <div style="width: 96%;"></div> <div>96%</div> </div>
1	B	321	<div> <div style="width: 95%;"></div> <div>95%</div> </div>
2	C	8	<div> <div style="width: 63%;"></div> <div>63%</div> </div>
3	D	9	<div> <div style="width: 56%;"></div> <div>56%</div> </div>
4	E	4	<div> <div style="width: 50%;"></div> <div>50%</div> </div>
5	F	6	<div> <div style="width: 33%;"></div> <div>33%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
6	G	8	 <div>75%25%</div>
7	H	2	 <div>100%</div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	MAN	F	6	-	-	-	X

2 Entry composition [i](#)


There are 15 unique types of molecules in this entry. The entry contains 6233 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-xylanase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	318	Total	C	N	O	S	0	2	0
			2515	1618	413	476	8			
1	B	321	Total	C	N	O	S	0	2	0
			2533	1627	416	482	8			

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	8	Total	C	N	O	0	0	0
			94	52	2	40			

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	D	9	Total	C	N	O	0	0	0
			105	58	2	45			

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.

e-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	E	4	Total	C	N	O	0	0	0
			50	28	2	20			

- Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-4)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	F	6	Total	C	N	O	0	0	0
			72	40	2	30			

- Molecule 6 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
6	G	8	Total	C	N	O	0	0	0
			94	52	2	40			

- Molecule 7 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
7	H	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 8 is UNKNOWN ATOM OR ION (three-letter code: UNX) (formula: X).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	B	1	Total	X			0	0
			1	1				
8	A	1	Total	X			0	0
			1	1				

- Molecule 9 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	A	1	Total	C	N	O	0	0
			14	8	1	5		
9	B	1	Total	C	N	O	0	0
			14	8	1	5		
9	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 10 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	B	1	Total	Mg	0	0
			1	1		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	1	Total	Mg	0	0
			1	1		

- Molecule 11 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
11	A	1	Total	C	O	0	0
			7	4	3		
11	B	1	Total	C	O	0	0
			7	4	3		
11	B	1	Total	C	O	0	0
			7	4	3		

- Molecule 12 is HEXAETHYLENE GLYCOL (three-letter code: P6G) (formula: $C_{12}H_{26}O_7$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
12	A	1	Total	C	O	0	0
			18	12	6		
12	A	1	Total	C	O	0	0
			13	8	5		

- Molecule 13 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
13	A	1	Total	C	O	0	0
			6	3	3		
13	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
13	A	1	Total	C	O	0	0
			6	3	3		
13	A	1	Total	C	O	0	0
			6	3	3		
13	B	1	Total	C	O	0	0
			6	3	3		
13	B	1	Total	C	O	0	0
			6	3	3		
13	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 14 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
14	B	1	Total	Cl	0	0
			1	1		


- Molecule 15 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
15	A	293	Total	O	0	13
			306	306		
15	B	288	Total	O	0	7
			295	295		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Beta-xylanase




- Molecule 1: Beta-xylanase



- Molecule 2: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




- Molecule 4: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose





- Molecule 5: alpha-D-mannopyranose-(1-4)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:



- Molecule 6: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:



- Molecule 7: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	52.41Å 99.34Å 147.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.20 – 1.95 30.20 – 1.95	Depositor EDS
% Data completeness (in resolution range)	91.7 (30.20-1.95) 86.6 (30.20-1.95)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.57 (at 1.95Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE: 1.9_1692)	Depositor
R, R_{free}	0.154 , 0.204 0.153 , 0.203	Depositor DCC
R_{free} test set	2000 reflections (3.92%)	wwPDB-VP
Wilson B-factor (Å ²)	18.0	Xtriage
Anisotropy	0.110	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 66.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6233	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.01% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, BMA, NAG, CL, UNX, P6G, PEG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.40	0/2596	0.53	0/3563
1	B	0.39	0/2615	0.53	0/3589
All	All	0.40	0/5211	0.53	0/7152

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2515	0	2370	8	0
1	B	2533	0	2382	11	0
2	C	94	0	79	0	0
3	D	105	0	88	2	0
4	E	50	0	43	3	0
5	F	72	0	61	1	0
6	G	94	0	79	0	0
7	H	28	0	25	0	0
8	A	1	0	0	0	0
8	B	1	0	0	0	0
9	A	14	0	13	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
9	B	28	0	26	0	0
10	A	1	0	0	0	0
10	B	1	0	0	0	0
11	A	7	0	10	0	0
11	B	14	0	20	3	0
12	A	31	0	39	2	0
13	A	24	0	32	1	0
13	B	18	0	24	2	0
14	B	1	0	0	0	0
15	A	306	0	0	1	0
15	B	295	0	0	0	1
All	All	6233	0	5291	25	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:269:ARG:HH22	13:B:424:GOL:H32	1.46	0.80
1:B:298:ALA:HA	11:B:422:PEG:H22	1.69	0.75
1:A:35:ARG:HE	12:A:426:P6G:H171	1.63	0.64
1:B:171:GLU:HG3	1:B:171:GLU:O	1.98	0.61
1:B:294:LYS:NZ	1:B:347:ALA:O	2.40	0.54
1:B:269:ARG:NH2	13:B:424:GOL:H32	2.20	0.54
5:F:5:MAN:H5	5:F:6:MAN:O5	2.10	0.51
1:B:40:PRO:HG2	11:B:422:PEG:H12	1.94	0.50
1:A:30:PRO:HA	1:A:31:PHE:HA	1.69	0.48
1:A:46:ALA:HB2	1:A:306:TRP:CE3	2.50	0.47
11:B:423:PEG:H31	11:B:423:PEG:H11	1.79	0.47
4:E:2:NAG:O3	4:E:3:BMA:H2	2.15	0.47
1:A:283:ALA:HB3	13:A:428:GOL:H32	1.98	0.46
1:B:70:ASN:HA	1:B:107:TYR:CE1	2.51	0.46
3:D:3:BMA:H3	3:D:9:MAN:H2	1.84	0.45
1:A:30:PRO:HA	1:A:31:PHE:HD1	1.83	0.44
1:A:343:GLN:HG2	15:A:771:HOH:O	2.19	0.42
12:A:426:P6G:H51	12:A:426:P6G:H82	1.56	0.42
1:B:60:LEU:HA	1:B:60:LEU:HD23	1.83	0.42
1:B:270:MET:SD	1:B:274:PRO:HB3	2.61	0.41
1:A:91:ASN:HD22	3:D:1:NAG:H83	1.85	0.41
4:E:3:BMA:H3	4:E:4:MAN:H2	1.54	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:166:ASN:HD22	4:E:1:NAG:H83	1.86	0.41
1:B:121:ALA:HA	1:B:122:PRO:HD3	1.93	0.40
1:B:224:LYS:HE3	1:B:258:GLY:O	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
15:B:611:HOH:O	15:B:680:HOH:O[1_655]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	318/321 (99%)	313 (98%)	5 (2%)	0	100	100
1	B	321/321 (100%)	315 (98%)	6 (2%)	0	100	100
All	All	639/642 (100%)	628 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	262/263 (100%)	262 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	265/263 (101%)	265 (100%)	0	100	100
All	All	527/526 (100%)	527 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

37 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	C	1	1,2	14,14,15	0.30	0	17,19,21	0.60	0
2	NAG	C	2	2	14,14,15	0.53	0	17,19,21	0.51	0
2	BMA	C	3	2	11,11,12	0.60	0	15,15,17	0.89	0
2	MAN	C	4	2	11,11,12	0.76	0	15,15,17	0.86	0
2	MAN	C	5	2	11,11,12	0.87	1 (9%)	15,15,17	0.97	1 (6%)
2	MAN	C	6	2	11,11,12	1.05	0	15,15,17	1.83	2 (13%)
2	MAN	C	7	2	11,11,12	0.68	0	15,15,17	1.13	1 (6%)
2	MAN	C	8	2	11,11,12	0.61	0	15,15,17	0.84	0
3	NAG	D	1	1,3	14,14,15	0.43	0	17,19,21	0.65	0
3	NAG	D	2	3	14,14,15	0.38	0	17,19,21	0.63	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	BMA	D	3	3	11,11,12	0.97	1 (9%)	15,15,17	0.99	1 (6%)
3	MAN	D	4	3	11,11,12	0.89	0	15,15,17	0.94	0
3	MAN	D	5	3	11,11,12	0.83	0	15,15,17	0.94	0
3	MAN	D	6	3	11,11,12	0.85	0	15,15,17	0.88	0
3	MAN	D	7	3	11,11,12	0.79	0	15,15,17	1.02	1 (6%)
3	MAN	D	8	3	11,11,12	0.65	0	15,15,17	0.82	0
3	MAN	D	9	3	11,11,12	1.00	0	15,15,17	0.92	0
4	NAG	E	1	1,4	14,14,15	0.56	0	17,19,21	0.76	0
4	NAG	E	2	4	14,14,15	0.39	0	17,19,21	0.60	0
4	BMA	E	3	4	11,11,12	1.09	1 (9%)	15,15,17	1.46	2 (13%)
4	MAN	E	4	4	11,11,12	0.82	1 (9%)	15,15,17	0.88	0
5	NAG	F	1	1,5	14,14,15	0.29	0	17,19,21	0.47	0
5	NAG	F	2	5	14,14,15	0.62	1 (7%)	17,19,21	0.71	0
5	BMA	F	3	5	11,11,12	1.02	1 (9%)	15,15,17	1.80	3 (20%)
5	MAN	F	4	5	11,11,12	0.72	0	15,15,17	0.83	0
5	MAN	F	5	5	11,11,12	1.25	2 (18%)	15,15,17	1.17	1 (6%)
5	MAN	F	6	5	11,11,12	0.85	0	15,15,17	0.85	0
6	NAG	G	1	1,6	14,14,15	0.52	0	17,19,21	0.44	0
6	NAG	G	2	6	14,14,15	0.18	0	17,19,21	0.62	0
6	BMA	G	3	6	11,11,12	0.63	0	15,15,17	0.75	0
6	MAN	G	4	6	11,11,12	0.88	0	15,15,17	1.00	0
6	MAN	G	5	6	11,11,12	0.78	1 (9%)	15,15,17	0.85	0
6	MAN	G	6	6	11,11,12	0.74	0	15,15,17	0.84	0
6	MAN	G	7	6	11,11,12	0.82	0	15,15,17	1.05	2 (13%)
6	MAN	G	8	6	11,11,12	0.66	0	15,15,17	0.87	0
7	NAG	H	1	1,7	14,14,15	0.33	0	17,19,21	0.39	0
7	NAG	H	2	7	14,14,15	0.22	0	17,19,21	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	C	2	2	-	0/6/23/26	0/1/1/1
2	BMA	C	3	2	-	0/2/19/22	0/1/1/1
2	MAN	C	4	2	-	0/2/19/22	0/1/1/1
2	MAN	C	5	2	-	2/2/19/22	0/1/1/1
2	MAN	C	6	2	-	0/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MAN	C	7	2	-	0/2/19/22	1/1/1/1
2	MAN	C	8	2	-	2/2/19/22	0/1/1/1
3	NAG	D	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	D	2	3	-	1/6/23/26	0/1/1/1
3	BMA	D	3	3	-	0/2/19/22	0/1/1/1
3	MAN	D	4	3	-	0/2/19/22	0/1/1/1
3	MAN	D	5	3	-	2/2/19/22	0/1/1/1
3	MAN	D	6	3	-	0/2/19/22	0/1/1/1
3	MAN	D	7	3	-	2/2/19/22	0/1/1/1
3	MAN	D	8	3	-	0/2/19/22	0/1/1/1
3	MAN	D	9	3	-	2/2/19/22	0/1/1/1
4	NAG	E	1	1,4	-	2/6/23/26	0/1/1/1
4	NAG	E	2	4	-	1/6/23/26	0/1/1/1
4	BMA	E	3	4	-	1/2/19/22	0/1/1/1
4	MAN	E	4	4	-	0/2/19/22	0/1/1/1
5	NAG	F	1	1,5	-	0/6/23/26	0/1/1/1
5	NAG	F	2	5	-	0/6/23/26	0/1/1/1
5	BMA	F	3	5	-	2/2/19/22	0/1/1/1
5	MAN	F	4	5	-	0/2/19/22	0/1/1/1
5	MAN	F	5	5	-	1/2/19/22	0/1/1/1
5	MAN	F	6	5	-	1/2/19/22	1/1/1/1
6	NAG	G	1	1,6	-	2/6/23/26	0/1/1/1
6	NAG	G	2	6	-	1/6/23/26	0/1/1/1
6	BMA	G	3	6	-	0/2/19/22	0/1/1/1
6	MAN	G	4	6	-	0/2/19/22	0/1/1/1
6	MAN	G	5	6	-	0/2/19/22	0/1/1/1
6	MAN	G	6	6	-	0/2/19/22	0/1/1/1
6	MAN	G	7	6	-	1/2/19/22	0/1/1/1
6	MAN	G	8	6	-	2/2/19/22	0/1/1/1
7	NAG	H	1	1,7	-	4/6/23/26	0/1/1/1
7	NAG	H	2	7	-	2/6/23/26	0/1/1/1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	F	5	MAN	O4-C4	2.58	1.49	1.43
4	E	3	BMA	O5-C5	2.44	1.48	1.43
3	D	3	BMA	C2-C3	2.33	1.55	1.52
5	F	5	MAN	C4-C5	2.18	1.57	1.53
5	F	2	NAG	O5-C1	-2.16	1.40	1.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	G	5	MAN	O5-C5	2.16	1.47	1.43
4	E	4	MAN	C1-C2	2.11	1.57	1.52
2	C	5	MAN	C2-C3	2.04	1.55	1.52
5	F	3	BMA	O5-C5	2.03	1.47	1.43

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	6	MAN	O3-C3-C2	5.36	120.25	109.99
5	F	3	BMA	C1-O5-C5	5.10	119.10	112.19
4	E	3	BMA	C1-C2-C3	-4.24	104.46	109.67
2	C	6	MAN	C1-C2-C3	-3.21	105.72	109.67
6	G	7	MAN	O2-C2-C3	-2.93	104.27	110.14
5	F	5	MAN	C1-O5-C5	2.92	116.15	112.19
5	F	3	BMA	O5-C1-C2	2.79	115.08	110.77
2	C	7	MAN	C1-O5-C5	2.72	115.87	112.19
3	D	7	MAN	O2-C2-C3	-2.58	104.98	110.14
3	D	3	BMA	C1-O5-C5	2.36	115.39	112.19
4	E	3	BMA	O3-C3-C2	2.24	114.28	109.99
2	C	5	MAN	O3-C3-C2	2.11	114.03	109.99
5	F	3	BMA	O6-C6-C5	-2.06	104.21	111.29
6	G	7	MAN	C1-O5-C5	2.01	114.91	112.19

There are no chirality outliers.

All (35) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	F	3	BMA	O5-C5-C6-O6
2	C	5	MAN	O5-C5-C6-O6
7	H	2	NAG	O5-C5-C6-O6
3	D	9	MAN	C4-C5-C6-O6
3	D	9	MAN	O5-C5-C6-O6
5	F	3	BMA	C4-C5-C6-O6
7	H	2	NAG	C4-C5-C6-O6
3	D	1	NAG	C8-C7-N2-C2
3	D	1	NAG	O7-C7-N2-C2
7	H	1	NAG	C8-C7-N2-C2
7	H	1	NAG	O7-C7-N2-C2
6	G	1	NAG	C8-C7-N2-C2
6	G	1	NAG	O7-C7-N2-C2
4	E	1	NAG	C8-C7-N2-C2
4	E	1	NAG	O7-C7-N2-C2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	D	7	MAN	O5-C5-C6-O6
5	F	5	MAN	O5-C5-C6-O6
3	D	5	MAN	C4-C5-C6-O6
6	G	8	MAN	O5-C5-C6-O6
2	C	5	MAN	C4-C5-C6-O6
2	C	8	MAN	O5-C5-C6-O6
3	D	5	MAN	O5-C5-C6-O6
3	D	7	MAN	C4-C5-C6-O6
2	C	1	NAG	C4-C5-C6-O6
5	F	6	MAN	O5-C5-C6-O6
6	G	8	MAN	C4-C5-C6-O6
7	H	1	NAG	C4-C5-C6-O6
6	G	7	MAN	O5-C5-C6-O6
4	E	3	BMA	C4-C5-C6-O6
2	C	1	NAG	O5-C5-C6-O6
3	D	2	NAG	C3-C2-N2-C7
6	G	2	NAG	C3-C2-N2-C7
7	H	1	NAG	O5-C5-C6-O6
2	C	8	MAN	C4-C5-C6-O6
4	E	2	NAG	O5-C5-C6-O6


All (2) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	7	MAN	C1-C2-C3-C4-C5-O5
5	F	6	MAN	C1-C2-C3-C4-C5-O5


9 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	1	NAG	1	0
3	D	3	BMA	1	0
3	D	9	MAN	1	0
5	F	5	MAN	1	0
4	E	2	NAG	1	0
4	E	4	MAN	1	0
4	E	1	NAG	1	0
5	F	6	MAN	1	0
4	E	3	BMA	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.




Oligosaccharide Chain D




Bond lengths



Bond angles




Torsions




Rings


Oligosaccharide Chain E




Bond lengths





Bond angles




Torsions



Rings

Oligosaccharide Chain F**Oligosaccharide Chain G**



5.6 Ligand geometry [i](#)

Of 20 ligands modelled in this entry, 2 are unknown and 3 are monoatomic - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
12	P6G	A	427	-	12,12,18	0.66	0	11,11,17	0.98	1 (9%)
13	GOL	B	426	-	5,5,5	0.35	0	5,5,5	0.32	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
11	PEG	A	425	-	6,6,6	0.65	0	5,5,5	0.20	0
13	GOL	B	425	-	5,5,5	0.37	0	5,5,5	0.23	0
13	GOL	A	429	-	5,5,5	0.34	0	5,5,5	0.32	0
9	NAG	A	423	1	14,14,15	0.33	0	17,19,21	0.41	0
12	P6G	A	426	-	17,17,18	0.58	0	16,16,17	1.84	4 (25%)
9	NAG	B	416	1	14,14,15	0.19	0	17,19,21	0.66	1 (5%)
11	PEG	B	422	-	6,6,6	0.65	0	5,5,5	0.24	0
11	PEG	B	423	-	6,6,6	0.65	0	5,5,5	0.33	0
13	GOL	B	424	-	5,5,5	0.37	0	5,5,5	0.19	0
9	NAG	B	419	1	14,14,15	0.30	0	17,19,21	0.55	0
13	GOL	A	430	-	5,5,5	0.36	0	5,5,5	0.28	0
13	GOL	A	428	-	5,5,5	0.35	0	5,5,5	0.32	0
13	GOL	A	431	-	5,5,5	0.35	0	5,5,5	0.21	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
12	P6G	A	427	-	-	1/10/10/16	-
13	GOL	B	426	-	-	2/4/4/4	-
11	PEG	A	425	-	-	0/4/4/4	-
13	GOL	B	425	-	-	2/4/4/4	-
13	GOL	A	429	-	-	2/4/4/4	-
9	NAG	A	423	1	-	2/6/23/26	0/1/1/1
12	P6G	A	426	-	-	6/15/15/16	-
9	NAG	B	416	1	-	4/6/23/26	0/1/1/1
11	PEG	B	422	-	-	0/4/4/4	-
11	PEG	B	423	-	-	2/4/4/4	-
13	GOL	B	424	-	-	4/4/4/4	-
9	NAG	B	419	1	-	2/6/23/26	0/1/1/1
13	GOL	A	430	-	-	2/4/4/4	-
13	GOL	A	428	-	-	0/4/4/4	-
13	GOL	A	431	-	-	2/4/4/4	-

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
12	A	426	P6G	O7-C8-C9	4.30	129.76	110.39
12	A	426	P6G	O10-C9-C8	-3.57	94.32	110.39
12	A	426	P6G	C8-O7-C6	-3.08	99.95	113.29
9	B	416	NAG	C1-O5-C5	2.26	115.26	112.19
12	A	427	P6G	O13-C12-C11	2.20	120.30	110.39
12	A	426	P6G	O16-C15-C14	-2.08	101.03	110.39

There are no chirality outliers.

All (31) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
13	B	426	GOL	O1-C1-C2-C3
13	A	429	GOL	O1-C1-C2-C3
13	B	424	GOL	O1-C1-C2-O2
13	B	424	GOL	O1-C1-C2-C3
13	B	424	GOL	C1-C2-C3-O3
13	A	430	GOL	O1-C1-C2-C3
11	B	423	PEG	C1-C2-O2-C3
9	A	423	NAG	C4-C5-C6-O6
9	B	416	NAG	C4-C5-C6-O6
9	B	416	NAG	O5-C5-C6-O6
9	A	423	NAG	O5-C5-C6-O6
9	B	416	NAG	C8-C7-N2-C2
9	B	416	NAG	O7-C7-N2-C2
9	B	419	NAG	C8-C7-N2-C2
9	B	419	NAG	O7-C7-N2-C2
13	B	425	GOL	O1-C1-C2-O2
13	B	425	GOL	O1-C1-C2-C3
13	A	431	GOL	O1-C1-C2-C3
11	B	423	PEG	O1-C1-C2-O2
12	A	426	P6G	C11-C12-O13-C14
13	B	426	GOL	O1-C1-C2-O2
13	A	429	GOL	O1-C1-C2-O2
13	B	424	GOL	O2-C2-C3-O3
13	A	430	GOL	O1-C1-C2-O2
13	A	431	GOL	O1-C1-C2-O2
12	A	426	P6G	C5-C6-O7-C8
12	A	427	P6G	C11-C12-O13-C14
12	A	426	P6G	O10-C11-C12-O13
12	A	426	P6G	O4-C5-C6-O7
12	A	426	P6G	O7-C8-C9-O10
12	A	426	P6G	O13-C14-C15-O16

There are no ring outliers.

5 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
12	A	426	P6G	2	0
11	B	422	PEG	2	0
11	B	423	PEG	1	0
13	B	424	GOL	2	0
13	A	428	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	318/321 (99%)	-0.13	3 (0%) 84 89	10, 19, 37, 99	0
1	B	321/321 (100%)	-0.02	11 (3%) 45 55	10, 21, 41, 64	0
All	All	639/642 (99%)	-0.08	14 (2%) 62 70	10, 20, 40, 99	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	31	PHE	10.5
1	A	30	PRO	9.6
1	B	127	LEU	5.1
1	A	347	ALA	3.4
1	B	347	ALA	2.8
1	B	171	GLU	2.8
1	B	27	PRO	2.6
1	B	130	GLU	2.6
1	B	126	THR	2.6
1	B	176	THR	2.6
1	B	122	PRO	2.6
1	B	131	ASP	2.5
1	B	128	THR	2.2
1	B	158	VAL	2.2

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	MAN	F	6	11/12	0.44	0.69	96,100,105,106	0
4	MAN	E	4	11/12	0.56	0.30	79,88,100,101	0
4	BMA	E	3	11/12	0.62	0.30	62,78,84,88	0
3	MAN	D	9	11/12	0.70	0.39	68,83,90,92	0
3	MAN	D	6	11/12	0.73	0.32	48,65,74,77	0
2	MAN	C	7	11/12	0.77	0.32	62,70,79,86	0
5	BMA	F	3	11/12	0.80	0.28	27,61,96,103	0
7	NAG	H	2	14/15	0.81	0.41	76,82,86,92	0
4	NAG	E	2	14/15	0.85	0.28	27,58,68,71	0
6	MAN	G	6	11/12	0.86	0.39	50,58,70,71	0
7	NAG	H	1	14/15	0.87	0.32	41,59,65,68	0
5	MAN	F	5	11/12	0.88	0.18	41,50,83,88	0
2	MAN	C	5	11/12	0.89	0.20	41,46,58,64	0
3	MAN	D	8	11/12	0.89	0.30	55,58,62,67	0
3	MAN	D	5	11/12	0.90	0.13	27,30,36,38	0
4	NAG	E	1	14/15	0.91	0.14	24,36,52,52	0
6	MAN	G	8	11/12	0.91	0.21	43,45,55,56	0
5	MAN	F	4	11/12	0.92	0.13	26,32,38,41	0
6	MAN	G	5	11/12	0.92	0.16	22,24,30,35	0
2	BMA	C	3	11/12	0.93	0.15	18,27,30,31	0
5	NAG	F	2	14/15	0.93	0.19	20,25,38,39	0
3	MAN	D	7	11/12	0.93	0.22	36,42,55,64	0
2	MAN	C	4	11/12	0.93	0.15	25,31,38,38	0
3	BMA	D	3	11/12	0.94	0.17	23,28,41,52	0
6	BMA	G	3	11/12	0.94	0.13	23,28,40,43	0
6	MAN	G	7	11/12	0.94	0.19	28,34,46,50	0
6	MAN	G	4	11/12	0.94	0.15	20,26,34,34	0
2	MAN	C	6	11/12	0.95	0.17	24,30,37,45	0
2	NAG	C	1	14/15	0.95	0.09	12,17,29,33	0
6	NAG	G	2	14/15	0.95	0.09	14,17,30,32	0
2	MAN	C	8	11/12	0.95	0.11	23,31,35,39	0
6	NAG	G	1	14/15	0.96	0.08	16,18,38,53	0
3	NAG	D	2	14/15	0.96	0.08	11,15,25,26	0
3	MAN	D	4	11/12	0.96	0.16	20,24,31,34	0
5	NAG	F	1	14/15	0.96	0.12	13,21,27,42	0
3	NAG	D	1	14/15	0.96	0.08	10,17,30,40	0
2	NAG	C	2	14/15	0.97	0.09	12,21,27,28	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain F:**


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain G:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
13	GOL	A	431	6/6	0.53	0.28	57,69,71,74	0
13	GOL	B	425	6/6	0.58	0.30	61,78,79,85	0
13	GOL	A	430	6/6	0.67	0.34	83,85,89,90	0
13	GOL	B	426	6/6	0.67	0.30	75,76,81,84	0
12	P6G	A	426	18/19	0.68	0.27	50,66,80,81	0
13	GOL	A	428	6/6	0.73	0.24	62,63,65,68	0
9	NAG	A	423	14/15	0.74	0.30	47,57,73,73	0
13	GOL	A	429	6/6	0.78	0.32	58,62,62,62	0
11	PEG	B	423	7/7	0.81	0.28	44,58,62,66	0

Continued on next page...



Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
8	UNX	A	401	1/1	0.84	0.72	30,30,30,30	0
9	NAG	B	419	14/15	0.86	0.26	39,54,71,77	0
13	GOL	B	424	6/6	0.87	0.12	58,59,59,60	0
8	UNX	B	401	1/1	0.89	0.62	30,30,30,30	0
9	NAG	B	416	14/15	0.90	0.40	40,62,68,72	0
11	PEG	A	425	7/7	0.91	0.25	40,47,50,53	0
12	P6G	A	427	13/19	0.91	0.18	36,44,54,56	0
11	PEG	B	422	7/7	0.92	0.24	42,50,63,67	0
10	MG	B	420	1/1	0.95	0.04	23,23,23,23	0
10	MG	A	424	1/1	0.97	0.09	21,21,21,21	0
14	CL	B	421	1/1	0.98	0.08	21,21,21,21	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around P6G A 426:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around P6G A 427:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.