



# Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 09:00 am BST

PDB ID : 5XX9  
Title : Crystal structure of Bacterioferritin  
Authors : Jobichen, C.; Rajesh, R.; Angayarkanni, J.; Sivaraman, J.  
Deposited on : 2017-07-01  
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	167	<div> <div>2%</div> <div> <div></div> <div>89%</div> <div>8%</div> <div>•</div> </div> </div>
1	B	167	<div> <div>%</div> <div> <div></div> <div>83%</div> <div>11%</div> <div>6%</div> <div>•</div> </div> </div>
1	C	167	<div> <div>2%</div> <div> <div></div> <div>84%</div> <div>11%</div> <div>•</div> </div> </div>
1	D	167	<div> <div>%</div> <div> <div></div> <div>82%</div> <div>12%</div> <div>6%</div> </div> </div>
1	E	167	<div> <div>3%</div> <div> <div></div> <div>83%</div> <div>13%</div> <div>•</div> </div> </div>
1	F	167	<div> <div>2%</div> <div> <div></div> <div>89%</div> <div>7%</div> <div>•</div> </div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 8454 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bacterioferritin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	162	Total	C	N	O	S	0	0	0
			1315	828	227	256	4			
1	B	157	Total	C	N	O	S	0	0	0
			1285	811	221	249	4			
1	C	162	Total	C	N	O	S	0	0	0
			1318	832	226	256	4			
1	D	157	Total	C	N	O	S	0	0	0
			1285	811	221	249	4			
1	E	161	Total	C	N	O	S	0	0	0
			1304	823	225	252	4			
1	F	161	Total	C	N	O	S	0	0	0
			1310	826	225	255	4			

- Molecule 2 is FE (II) ION (three-letter code: FE2) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	D	2	Total	Fe	0	0
			2	2		
2	E	3	Total	Fe	0	0
			3	3		
2	B	1	Total	Fe	0	0
			1	1		
2	C	2	Total	Fe	0	0
			2	2		
2	A	3	Total	Fe	0	0
			3	3		
2	F	1	Total	Fe	0	0
			1	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	121	Total 121	O 121	0	0
3	B	102	Total 102	O 102	0	0
3	C	94	Total 94	O 94	0	0
3	D	110	Total 110	O 110	0	0
3	E	90	Total 90	O 90	0	0
3	F	108	Total 108	O 108	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Bacterioferritin

Chain A: 




#### • Molecule 1: Bacterioferritin

Chain B: 





#### • Molecule 1: Bacterioferritin

Chain C: 




#### • Molecule 1: Bacterioferritin

Chain D: 




#### • Molecule 1: Bacterioferritin

Chain E: 



#### • Molecule 1: Bacterioferritin



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	123.76 Å   123.76 Å   172.30 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	43.76 – 2.60 43.76 – 2.60	Depositor EDS
% Data completeness (in resolution range)	97.0 (43.76-2.60) 85.6 (43.76-2.60)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.32 (at 2.61 Å)	Xtriage
Refinement program	PHENIX (dev_2733)	Depositor
R, $R_{free}$	0.161 , 0.208 0.163 , 0.209	Depositor DCC
$R_{free}$ test set	1906 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	15.8	Xtriage
Anisotropy	0.068	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 39.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.023 for $-1/2^*h+1/2^*k-1/2^*l, 1/2^*h-1/2^*k-1/2^*l, -h-k$ 0.004 for $-1/2^*h+1/2^*k+1/2^*l, 1/2^*h-1/2^*k+1/2^*l, h+k$ 0.011 for $-1/2^*h-1/2^*k+1/2^*l, -1/2^*h-1/2^*k-1/2^*l, h-k$ 0.015 for $-1/2^*h-1/2^*k-1/2^*l, -1/2^*h-1/2^*k+1/2^*l, -h+k$ 0.032 for $-h, k, -l$	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	8454	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.87% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FE2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.42	0/1336	0.57	1/1805 (0.1%)
1	B	0.43	0/1306	0.56	0/1764
1	C	0.42	0/1341	0.56	0/1814
1	D	0.41	0/1306	0.53	0/1764
1	E	0.42	0/1325	0.58	2/1790 (0.1%)
1	F	0.41	0/1332	0.56	1/1801 (0.1%)
All	All	0.42	0/7946	0.56	4/10738 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	161	PRO	N-CA-CB	5.92	110.40	103.30
1	E	159	PRO	N-CA-CB	5.92	110.40	103.30
1	A	159	PRO	N-CA-CB	5.32	109.69	103.30
1	F	159	PRO	N-CA-CB	5.31	109.67	103.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1315	0	1280	7	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1285	0	1262	11	0
1	C	1318	0	1289	11	0
1	D	1285	0	1262	11	0
1	E	1304	0	1268	13	0
1	F	1310	0	1276	6	0
2	A	3	0	0	0	0
2	B	1	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
2	E	3	0	0	0	0
2	F	1	0	0	0	0
3	A	121	0	0	3	2
3	B	102	0	0	1	0
3	C	94	0	0	2	2
3	D	110	0	0	3	0
3	E	90	0	0	2	0
3	F	108	0	0	2	0
All	All	8454	0	7637	58	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:156:GLN:O	3:E:401:HOH:O	1.81	0.97
1:F:53:ARG:NH2	3:F:301:HOH:O	1.97	0.93
1:B:50:ASP:OD1	1:B:53:ARG:NH1	2.15	0.78
1:A:94:GLU:OE2	3:A:301:HOH:O	2.02	0.76
1:E:100:ARG:NH2	3:E:402:HOH:O	2.05	0.72
1:F:161:PRO:HA	3:F:303:HOH:O	1.90	0.71
1:A:78:ARG:NH1	3:A:302:HOH:O	2.24	0.70
1:F:34:HIS:ND1	1:F:160:ASP:OD2	2.23	0.69
1:D:127:GLU:OE2	3:D:301:HOH:O	2.13	0.67
1:C:151:SER:O	3:C:303:HOH:O	2.16	0.59
1:B:83:VAL:HG11	1:B:150:LEU:HD11	1.85	0.57
1:C:159:PRO:HG2	1:C:161:PRO:HG3	1.88	0.55
1:A:100:ARG:NH2	3:A:306:HOH:O	2.40	0.54
1:C:28:HIS:O	1:C:32:GLN:HG3	2.07	0.54
1:A:28:HIS:O	1:A:32:GLN:HG3	2.08	0.53
1:E:47:GLU:HG3	1:E:130:HIS:NE2	2.24	0.53

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:37:TRP:CZ3	1:E:83:VAL:HG22	2.44	0.52
1:B:81:GLN:NE2	3:B:303:HOH:O	2.38	0.52
1:C:37:TRP:CZ3	1:C:83:VAL:HG22	2.44	0.52
1:B:149:TYR:O	1:B:152:THR:HB	2.10	0.52
1:C:128:GLU:O	1:C:131:ILE:HG22	2.08	0.52
1:B:28:HIS:O	1:B:32:GLN:HG3	2.10	0.51
1:E:39:LYS:NZ	1:E:155:GLU:OE2	2.29	0.51
1:C:1:MET:O	1:C:66:ASP:HB2	2.10	0.50
1:B:37:TRP:CZ3	1:B:83:VAL:HG13	2.46	0.50
1:C:47:GLU:HG3	1:C:130:HIS:NE2	2.27	0.50
1:D:35:LYS:O	3:D:303:HOH:O	2.20	0.48
1:B:47:GLU:HG3	1:B:130:HIS:NE2	2.28	0.48
1:E:117:ALA:O	1:E:121:GLU:HG3	2.13	0.48
1:C:34:HIS:ND1	1:C:160:ASP:OD2	2.40	0.47
1:D:94:GLU:OE2	3:D:301:HOH:O	2.20	0.47
1:E:28:HIS:O	1:E:32:GLN:HG3	2.15	0.47
1:A:128:GLU:O	1:A:131:ILE:HG22	2.15	0.47
1:A:47:GLU:HG3	1:A:130:HIS:NE2	2.30	0.47
1:C:76:HIS:ND1	3:C:301:HOH:O	2.05	0.47
1:D:47:GLU:HG3	1:D:130:HIS:NE2	2.30	0.47
1:A:13:GLU:HG2	1:A:100:ARG:HD3	1.96	0.46
1:B:155:GLU:O	1:B:156:GLN:HB2	2.17	0.45
1:F:128:GLU:O	1:F:131:ILE:HG22	2.16	0.45
1:B:117:ALA:O	1:B:121:GLU:HG3	2.16	0.45
1:F:123:ILE:O	1:F:127:GLU:HG2	2.15	0.45
1:F:117:ALA:O	1:F:121:GLU:HG3	2.17	0.44
1:E:1:MET:O	1:E:66:ASP:HB2	2.16	0.44
1:E:28:HIS:CD2	1:E:86:MET:HG2	2.53	0.44
1:E:123:ILE:O	1:E:127:GLU:HG2	2.18	0.43
1:B:76:HIS:ND1	1:E:9:GLU:OE1	2.49	0.43
1:D:123:ILE:O	1:D:127:GLU:HG2	2.18	0.43
1:D:68:LEU:HD22	1:D:69:PRO:HD2	2.00	0.43
1:D:155:GLU:O	1:D:157:THR:N	2.51	0.43
1:D:19:LEU:HA	1:D:22:ILE:HD12	2.00	0.42
1:C:156:GLN:O	1:C:157:THR:HG23	2.21	0.41
1:B:128:GLU:O	1:B:131:ILE:HG22	2.20	0.41
1:D:98:ILE:O	1:D:102:ARG:HG3	2.20	0.41
1:D:128:GLU:O	1:D:131:ILE:HG22	2.21	0.41
1:E:37:TRP:CH2	1:E:83:VAL:HG22	2.56	0.41
1:C:101:LEU:HD13	1:C:124:LEU:HA	2.02	0.41
1:E:13:GLU:HG2	1:E:100:ARG:HD3	2.03	0.40

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:117:ALA:O	1:D:121:GLU:HG3	2.21	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:318:HOH:O	3:C:362:HOH:O[3_555]	2.07	0.13
3:A:395:HOH:O	3:C:379:HOH:O[3_555]	2.18	0.02

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	160/167 (96%)	157 (98%)	1 (1%)	2 (1%)	12	24
1	B	155/167 (93%)	153 (99%)	2 (1%)	0	100	100
1	C	160/167 (96%)	156 (98%)	2 (1%)	2 (1%)	12	24
1	D	155/167 (93%)	153 (99%)	1 (1%)	1 (1%)	25	47
1	E	159/167 (95%)	156 (98%)	1 (1%)	2 (1%)	12	24
1	F	159/167 (95%)	155 (98%)	2 (1%)	2 (1%)	12	24
All	All	948/1002 (95%)	930 (98%)	9 (1%)	9 (1%)	17	35

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	160	ASP
1	A	161	PRO
1	D	156	GLN
1	C	156	GLN
1	F	157	THR
1	C	159	PRO

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	E	160	ASP
1	E	159	PRO
1	F	159	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	140/147 (95%)	139 (99%)	1 (1%)	84	94
1	B	138/147 (94%)	137 (99%)	1 (1%)	84	94
1	C	142/147 (97%)	141 (99%)	1 (1%)	84	94
1	D	138/147 (94%)	137 (99%)	1 (1%)	84	94
1	E	138/147 (94%)	136 (99%)	2 (1%)	67	85
1	F	140/147 (95%)	138 (99%)	2 (1%)	67	85
All	All	836/882 (95%)	828 (99%)	8 (1%)	76	90

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	150	LEU
1	B	150	LEU
1	C	150	LEU
1	D	150	LEU
1	E	150	LEU
1	E	156	GLN
1	F	150	LEU
1	F	160	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 12 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	162/167 (97%)	-0.57	4 (2%) 57 51	9, 16, 33, 59	0
1	B	157/167 (94%)	-0.57	1 (0%) 89 88	10, 15, 28, 48	0
1	C	162/167 (97%)	-0.50	4 (2%) 57 51	11, 17, 35, 58	0
1	D	157/167 (94%)	-0.51	1 (0%) 89 88	10, 16, 31, 43	0
1	E	161/167 (96%)	-0.48	5 (3%) 49 42	9, 16, 34, 54	0
1	F	161/167 (96%)	-0.50	3 (1%) 66 62	10, 16, 32, 53	0
All	All	960/1002 (95%)	-0.52	18 (1%) 66 62	9, 16, 34, 59	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	158	GLN	4.3
1	E	159	PRO	4.0
1	F	158	GLN	3.9
1	E	158	GLN	3.2
1	E	160	ASP	3.1
1	F	161	PRO	3.1
1	D	100	ARG	3.0
1	F	159	PRO	2.9
1	B	157	THR	2.9
1	E	161	PRO	2.9
1	C	159	PRO	2.9
1	E	157	THR	2.8
1	A	159	PRO	2.4
1	C	162	SER	2.4
1	A	158	GLN	2.2
1	A	161	PRO	2.1
1	C	1	MET	2.1
1	A	157	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	FE2	E	203	1/1	0.81	0.48	86,86,86,86	0
2	FE2	C	202	1/1	0.85	0.24	88,88,88,88	0
2	FE2	E	202	1/1	0.86	0.16	78,78,78,78	0
2	FE2	A	203	1/1	0.86	0.09	75,75,75,75	0
2	FE2	D	202	1/1	0.87	0.28	81,81,81,81	0
2	FE2	A	202	1/1	0.93	0.15	75,75,75,75	0
2	FE2	B	200	1/1	0.93	0.08	32,32,32,32	0
2	FE2	D	201	1/1	0.94	0.07	30,30,30,30	0
2	FE2	C	201	1/1	0.94	0.07	29,29,29,29	0
2	FE2	E	201	1/1	0.97	0.06	30,30,30,30	0
2	FE2	F	200	1/1	0.97	0.09	29,29,29,29	0
2	FE2	A	201	1/1	0.98	0.04	30,30,30,30	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.