



Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 07:05 pm BST

PDB ID : 5XYS
Title : Structure of 6-aminohexanoate-oligomer hydrolase from *Arthrobacter* sp. KI72., D122V muntant
Authors : Negoro, S.; Shibata, N.; Nagai, K.; Higuchi, Y.
Deposited on : 2017-07-10
Resolution : 1.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1202 (1.10-1.02)
Clashscore	141614	1252 (1.10-1.02)
Ramachandran outliers	138981	1204 (1.10-1.02)
Sidechain outliers	138945	1202 (1.10-1.02)
RSRZ outliers	127900	1178 (1.10-1.02)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	355	<div> <div>7%</div> <div> <div></div> <div>82%</div> <div>10%</div> <div>••</div> <div>7%</div> </div> </div>
1	B	355	<div> <div>5%</div> <div> <div></div> <div>82%</div> <div>10%</div> <div>••</div> <div>6%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PO4	A	403	-	-	X	-
3	PO4	B	401	-	-	X	X

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5728 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Endo-type 6-aminohexanoate oligomer hydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	331	Total	C	N	O	S	0	18	0
			2550	1600	465	475	10			
1	B	332	Total	C	N	O	S	0	18	0
			2545	1597	459	479	10			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	122	VAL	ASP	engineered mutation	UNP Q57326
B	122	VAL	ASP	engineered mutation	UNP Q57326

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	294	Total	O	0	0
			294	294		
4	B	317	Total	O	0	0
			317	317		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Endo-type 6-aminohexanoate oligomer hydrolase



- Molecule 1: Endo-type 6-aminohexanoate oligomer hydrolase



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	70.46Å 144.78Å 128.57Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	72.39 – 1.05 45.13 – 1.05	Depositor EDS
% Data completeness (in resolution range)	90.2 (72.39-1.05) 90.1 (45.13-1.05)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.60 (at 1.05Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.133 , 0.152 0.139 , 0.157	Depositor DCC
R_{free} test set	13830 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	18.1	Xtriage
Anisotropy	0.063	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 49.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	5728	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.85	5/2635 (0.2%)	1.01	12/3577 (0.3%)
1	B	0.99	11/2624 (0.4%)	1.03	18/3564 (0.5%)
All	All	0.92	16/5259 (0.3%)	1.02	30/7141 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
All	All	0	4

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	305[A]	MET	CG-SD	12.84	2.14	1.81
1	B	305[B]	MET	CG-SD	12.84	2.14	1.81
1	B	292	SER	CA-CB	8.45	1.65	1.52
1	A	88[A]	ARG	CG-CD	-8.18	1.31	1.51
1	A	88[B]	ARG	CG-CD	-8.18	1.31	1.51
1	B	66	GLU	CD-OE1	7.42	1.33	1.25
1	B	65	TYR	CE2-CZ	6.95	1.47	1.38
1	B	65	TYR	CZ-OH	-6.64	1.26	1.37
1	B	244	ARG	CA-CB	6.38	1.68	1.53
1	A	150[A]	ALA	CA-CB	6.32	1.65	1.52
1	A	150[B]	ALA	CA-CB	6.32	1.65	1.52
1	B	88[A]	ARG	CZ-NH2	-5.64	1.25	1.33
1	B	88[B]	ARG	CZ-NH2	-5.64	1.25	1.33
1	B	272	ILE	CA-CB	5.56	1.67	1.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	159	LYS	CE-NZ	5.36	1.62	1.49
1	A	49	GLU	CB-CG	-5.15	1.42	1.52

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	304	ASP	O-C-N	10.30	139.19	122.70
1	A	88[A]	ARG	CB-CG-CD	-9.74	86.28	111.60
1	A	88[B]	ARG	CB-CG-CD	-9.74	86.28	111.60
1	B	88[A]	ARG	NE-CZ-NH1	9.14	124.87	120.30
1	B	88[B]	ARG	NE-CZ-NH1	9.14	124.87	120.30
1	A	305[A]	MET	CG-SD-CE	8.77	114.23	100.20
1	A	305[B]	MET	CG-SD-CE	8.77	114.23	100.20
1	A	86	ASP	CB-CG-OD2	8.03	125.53	118.30
1	A	88[A]	ARG	NE-CZ-NH1	-7.02	116.79	120.30
1	A	88[B]	ARG	NE-CZ-NH1	-7.02	116.79	120.30
1	B	304	ASP	CA-C-N	-6.70	102.46	117.20
1	A	304	ASP	CB-CG-OD1	6.17	123.85	118.30
1	B	305[A]	MET	CB-CG-SD	-6.03	94.32	112.40
1	B	305[B]	MET	CB-CG-SD	-6.03	94.32	112.40
1	B	318	ILE	CB-CG1-CD1	-6.01	97.07	113.90
1	B	304	ASP	CB-CG-OD1	5.71	123.44	118.30
1	B	86[A]	ASP	CB-CG-OD2	5.68	123.41	118.30
1	B	86[B]	ASP	CB-CG-OD2	5.68	123.41	118.30
1	B	88[A]	ARG	NH1-CZ-NH2	-5.62	113.22	119.40
1	B	88[B]	ARG	NH1-CZ-NH2	-5.62	113.22	119.40
1	A	288	LYS	CD-CE-NZ	-5.58	98.85	111.70
1	B	65	TYR	CD1-CE1-CZ	5.32	124.59	119.80
1	A	305[A]	MET	CA-CB-CG	5.29	122.29	113.30
1	A	305[B]	MET	CA-CB-CG	5.29	122.29	113.30
1	B	305[A]	MET	C-N-CA	5.19	134.68	121.70
1	B	305[B]	MET	C-N-CA	5.19	134.68	121.70
1	B	65	TYR	CE1-CZ-CE2	-5.10	111.64	119.80
1	B	163	ARG	NE-CZ-NH2	-5.07	117.76	120.30
1	B	304	ASP	CB-CG-OD2	-5.06	113.75	118.30
1	A	156	TYR	CB-CG-CD2	-5.02	117.99	121.00

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	151[A]	ARG	Sidechain

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	A	88[A]	ARG	Sidechain
1	B	211	ARG	Sidechain
1	B	88[A]	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2550	0	2523	48	0
1	B	2545	0	2518	47	0
2	A	12	0	15	0	0
3	A	5	0	0	7	0
3	B	5	0	0	3	0
4	A	294	0	0	11	0
4	B	317	0	0	4	1
All	All	5728	0	5056	94	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (94) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:88[B]:ARG:CG	1:A:88[B]:ARG:CD	1.78	1.55
1:B:305[A]:MET:CG	1:B:305[A]:MET:SD	2.14	1.34
1:A:88[A]:ARG:NH1	3:A:403:PO4:O3	1.64	1.28
1:A:82:ARG:NH2	1:A:278:MET:O	1.78	1.15
1:B:82[B]:ARG:NH1	1:B:278:MET:O	1.89	1.05
1:A:149[B]:SER:OG	1:A:192[B]:TRP:CH2	2.13	1.02
1:A:167:GLU:HG2	4:A:712:HOH:O	1.61	1.00
1:A:29:VAL:CG2	1:A:152:SER:OG	2.12	0.98
1:A:88[A]:ARG:CZ	3:A:403:PO4:P	2.54	0.95
1:B:30:PHE:HE1	1:B:149[B]:SER:HG	0.99	0.94
1:B:30:PHE:HE1	1:B:149[B]:SER:OG	1.49	0.94
1:B:91[B]:ALA:HA	1:B:91[B]:ALA:N	1.24	0.89
1:A:29:VAL:HB	1:A:152:SER:OG	1.72	0.89

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:91[B]:ALA:HA	1:B:91[B]:ALA:C	1.89	0.88
1:A:239:GLN:OE1	4:A:501:HOH:O	1.91	0.88
1:A:88[A]:ARG:CZ	3:A:403:PO4:O3	2.25	0.84
1:A:149[B]:SER:OG	1:A:192[B]:TRP:CZ2	2.29	0.83
1:B:66:GLU:HG3	4:B:525:HOH:O	1.78	0.83
1:A:29:VAL:CB	1:A:152:SER:OG	2.27	0.81
1:A:221:VAL:HG21	1:A:305[A]:MET:HB2	1.65	0.79
1:A:67[B]:GLU:OE1	4:A:502:HOH:O	1.99	0.79
1:A:29:VAL:HG21	1:A:152:SER:OG	1.83	0.77
1:B:88[B]:ARG:NH2	3:B:401:PO4:P	2.57	0.77
1:A:88[A]:ARG:CZ	3:A:403:PO4:O1	2.33	0.76
1:B:221:VAL:HG21	1:B:305[A]:MET:HB2	1.67	0.75
1:A:88[A]:ARG:NH2	3:A:403:PO4:P	2.61	0.73
1:A:29:VAL:HB	1:A:152:SER:CB	2.19	0.72
1:B:90[B]:GLY:C	1:B:91[B]:ALA:HA	1.73	0.72
1:B:86[B]:ASP:OD1	1:B:288[B]:LYS:HD3	1.88	0.72
1:B:86[B]:ASP:CG	1:B:288[B]:LYS:HD3	2.10	0.72
1:B:86[B]:ASP:OD2	1:B:287:ALA:C	2.29	0.71
1:B:305[A]:MET:SD	1:B:305[A]:MET:CB	2.79	0.71
1:B:88[B]:ARG:CZ	3:B:401:PO4:O1	2.37	0.71
1:B:30:PHE:CE1	1:B:149[B]:SER:OG	2.33	0.69
1:A:36:ASP:OD2	4:A:503:HOH:O	2.14	0.66
1:A:167:GLU:CG	4:A:712:HOH:O	2.29	0.66
1:B:90[B]:GLY:HA2	4:B:539:HOH:O	1.95	0.65
1:B:82[B]:ARG:CZ	1:B:278:MET:O	2.44	0.65
1:A:156:TYR:OH	1:B:122[B]:VAL:HG21	1.97	0.64
1:A:88[A]:ARG:NE	3:A:403:PO4:O1	2.30	0.64
1:B:86[B]:ASP:OD2	1:B:287:ALA:CB	2.46	0.64
1:A:88[A]:ARG:NH2	3:A:403:PO4:O4	2.31	0.64
1:A:50[B]:MET:SD	1:A:52:ARG:CZ	2.87	0.62
1:B:91[B]:ALA:HA	1:B:92:VAL:N	2.11	0.61
1:B:82[B]:ARG:NH2	1:B:278:MET:O	2.33	0.61
1:B:88[B]:ARG:NH2	3:B:401:PO4:O4	2.33	0.60
1:B:86[B]:ASP:OD2	1:B:287:ALA:HB1	2.02	0.60
1:B:167:GLU:HG2	4:B:537:HOH:O	2.01	0.59
1:B:82[A]:ARG:NH1	1:B:280:PRO:N	2.51	0.59
1:A:130:HIS:HE1	4:A:709:HOH:O	1.85	0.59
1:A:328[A]:ARG:NH1	4:A:505:HOH:O	2.39	0.55
1:A:130:HIS:HD2	4:A:745:HOH:O	1.91	0.52
1:B:305[A]:MET:SD	1:B:305[A]:MET:HB3	2.50	0.51
1:A:167:GLU:HG3	4:A:605:HOH:O	2.11	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:277:ARG:HD2	1:B:319:ASP:OD1	2.11	0.49
1:A:88[B]:ARG:CG	1:A:88[B]:ARG:NE	2.70	0.49
1:B:76:HIS:HD2	4:B:734:HOH:O	1.95	0.49
1:A:221:VAL:HG21	1:A:305[B]:MET:HB2	1.95	0.48
1:A:192[B]:TRP:CH2	1:A:195:THR:HG21	2.48	0.48
1:B:82[A]:ARG:HH12	1:B:279:SER:C	2.17	0.47
1:A:26:GLY:O	1:A:192[B]:TRP:CD1	2.68	0.47
1:A:66:GLU:HG3	4:A:733:HOH:O	2.15	0.46
1:B:82[A]:ARG:HH11	1:B:280:PRO:N	2.13	0.46
1:A:88[A]:ARG:HD3	1:A:88[A]:ARG:HA	1.07	0.46
1:A:192[A]:TRP:CH2	1:A:195:THR:HG21	2.49	0.46
1:A:29:VAL:HG23	1:A:152:SER:OG	2.12	0.45
1:A:30:PHE:CD1	1:A:30:PHE:N	2.84	0.45
1:B:86[B]:ASP:OD1	1:B:288[B]:LYS:CD	2.61	0.45
1:B:91[B]:ALA:HB3	1:B:91[B]:ALA:H	1.72	0.45
1:A:183[A]:MET:HE3	1:A:183[A]:MET:HA	1.98	0.45
1:A:35:ASN:OD1	1:A:155:VAL:HA	2.17	0.45
1:A:26:GLY:O	1:A:192[A]:TRP:CD1	2.70	0.44
1:A:87[B]:ALA:HB1	1:A:92:VAL:HG21	2.00	0.44
1:B:88[A]:ARG:HG2	1:B:302:HIS:CE1	2.53	0.43
1:A:88[B]:ARG:HG2	1:A:88[B]:ARG:CD	2.22	0.43
1:B:305[A]:MET:CG	1:B:305[A]:MET:CE	2.96	0.42
1:A:244[A]:ARG:HG3	4:A:684:HOH:O	2.19	0.42
1:B:50[B]:MET:HB2	1:B:50[B]:MET:HE3	1.90	0.42
1:B:91[B]:ALA:HB2	1:B:91[B]:ALA:N	2.11	0.41
1:B:69:PRO:HB2	1:B:183[A]:MET:HG3	2.03	0.41
1:B:88[A]:ARG:HG2	1:B:302:HIS:ND1	2.36	0.41
1:A:183[A]:MET:CA	1:A:183[A]:MET:HE3	2.51	0.40
1:A:38:PHE:CD1	1:A:67[A]:GLU:HG2	2.56	0.40
1:B:60:ILE:HD11	1:B:213:LEU:HD21	2.04	0.40
1:A:206:ARG:NH2	1:A:211[B]:ARG:HH22	2.20	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:577:HOH:O	4:B:577:HOH:O[3_455]	2.10	0.10

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	345/355 (97%)	331 (96%)	14 (4%)	0	100	100
1	B	346/355 (98%)	330 (95%)	16 (5%)	0	100	100
All	All	691/710 (97%)	661 (96%)	30 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	257/262 (98%)	249 (97%)	8 (3%)	40	7
1	B	258/262 (98%)	251 (97%)	7 (3%)	44	9
All	All	515/524 (98%)	500 (97%)	15 (3%)	52	8

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	30	PHE
1	A	39	ASP
1	A	55	PHE
1	A	88[A]	ARG
1	A	88[B]	ARG
1	A	128	LEU
1	A	305[A]	MET
1	A	305[B]	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	55	PHE
1	B	128	LEU
1	B	209[A]	ASP
1	B	209[B]	ASP
1	B	305[A]	MET
1	B	305[B]	MET
1	B	318	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	130	HIS
1	A	257	GLN
1	A	284	ASN
1	B	76	HIS
1	B	251	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	A	402	-	5,5,5	1.19	1 (20%)	5,5,5	2.04	1 (20%)
3	PO4	B	401	-	4,4,4	0.40	0	6,6,6	0.98	0
2	GOL	A	401	-	5,5,5	0.63	0	5,5,5	0.92	0
3	PO4	A	403	-	4,4,4	0.66	0	6,6,6	1.25	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	402	-	-	2/4/4/4	-
2	GOL	A	401	-	-	0/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	402	GOL	O2-C2	-2.59	1.35	1.43

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	402	GOL	O2-C2-C1	-3.89	91.98	109.12
3	A	403	PO4	O2-P-O1	-2.35	102.30	110.89

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	402	GOL	O1-C1-C2-C3
2	A	402	GOL	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	401	PO4	3	0
3	A	403	PO4	7	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	331/355 (93%)	0.41	25 (7%) 13 15	15, 20, 36, 64	0
1	B	332/355 (93%)	0.20	18 (5%) 25 23	15, 19, 32, 45	0
All	All	663/710 (93%)	0.31	43 (6%) 18 19	15, 19, 33, 64	0

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	30	PHE	11.1
1	A	260	PRO	10.0
1	A	29	VAL	8.4
1	A	28	PRO	7.8
1	A	192[A]	TRP	7.2
1	B	29	VAL	6.9
1	B	18	VAL	6.2
1	B	30	PHE	6.1
1	A	259	PRO	5.5
1	B	134	PHE	5.2
1	A	134	PHE	5.1
1	B	31	GLY	4.9
1	B	28	PRO	4.9
1	B	19	ASP	4.8
1	B	192	TRP	4.7
1	A	31	GLY	4.7
1	A	322	THR	4.2
1	B	260	PRO	4.1
1	B	322	THR	4.0
1	A	239	GLN	3.9
1	A	258	VAL	3.8
1	A	323	THR	3.7
1	A	19	ASP	3.6
1	A	150[A]	ALA	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	86[A]	ASP	3.5
1	A	320	LEU	3.4
1	B	321	PRO	3.3
1	A	135	ALA	3.3
1	A	319	ASP	3.2
1	B	135	ALA	3.1
1	A	27	PRO	2.9
1	A	321	PRO	2.9
1	B	259	PRO	2.8
1	B	323	THR	2.7
1	B	20	PRO	2.6
1	B	27	PRO	2.4
1	A	39	ASP	2.3
1	A	330	ARG	2.3
1	A	255	ALA	2.2
1	A	94	LEU	2.2
1	A	325	GLY	2.2
1	B	330	ARG	2.1
1	A	149[A]	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	PO4	B	401	5/5	0.78	0.48	29,32,33,36	0
3	PO4	A	403	5/5	0.85	0.50	31,32,37,37	0
2	GOL	A	402	6/6	0.87	0.20	28,35,45,54	0
2	GOL	A	401	6/6	0.96	0.09	22,28,36,41	0

6.5 Other polymers [i](#)

There are no such residues in this entry.