



# Full wwPDB X-ray Structure Validation Report ⓘ

Dec 14, 2022 – 04:07 PM JST

PDB ID : 7XYX  
Title : Crystal structure of ZYG11B bound to CFLH degron  
Authors : Dong, C.; Yan, X.; Li, Y.  
Deposited on : 2022-06-02  
Resolution : 2.87 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.31.3
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.31.3


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 2.87 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2691 (2.90-2.86)
Clashscore	141614	2947 (2.90-2.86)
Ramachandran outliers	138981	2868 (2.90-2.86)
Sidechain outliers	138945	2871 (2.90-2.86)
RSRZ outliers	127900	2629 (2.90-2.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	250	
1	B	250	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3904 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein zyg-11 homolog B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	243	Total	C	N	O	S	0	0	0
			1943	1245	329	358	11			
1	A	244	Total	C	N	O	S	0	0	0
			1956	1253	332	361	10			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	479	CYS	-	expression tag	UNP Q9C0D3
B	480	PHE	-	expression tag	UNP Q9C0D3
B	481	LEU	-	expression tag	UNP Q9C0D3
B	482	HIS	-	expression tag	UNP Q9C0D3
B	483	VAL	-	expression tag	UNP Q9C0D3
B	484	GLY	-	expression tag	UNP Q9C0D3
A	479	CYS	-	expression tag	UNP Q9C0D3
A	480	PHE	-	expression tag	UNP Q9C0D3
A	481	LEU	-	expression tag	UNP Q9C0D3
A	482	HIS	-	expression tag	UNP Q9C0D3
A	483	VAL	-	expression tag	UNP Q9C0D3
A	484	GLY	-	expression tag	UNP Q9C0D3

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	3	Total	O	0	0
			3	3		
2	A	2	Total	O	0	0
			2	2		

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Protein zyg-11 homolog B

Chain B: 



#### • Molecule 1: Protein zyg-11 homolog B

Chain A: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	52.78Å 97.37Å 122.89Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.40 – 2.87 46.40 – 2.87	Depositor EDS
% Data completeness (in resolution range)	98.5 (46.40-2.87) 98.6 (46.40-2.87)	Depositor EDS
$R_{merge}$	0.26	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.51 (at 2.86Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, $R_{free}$	0.202 , 0.264 0.204 , 0.257	Depositor DCC
$R_{free}$ test set	680 reflections (4.57%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	68.6	Xtriage
Anisotropy	0.425	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 51.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3904	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	74.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.39% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.55	0/2003	0.68	1/2723 (0.0%)
1	B	0.51	0/1988	0.66	0/2701
All	All	0.53	0/3991	0.67	1/5424 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	619	LEU	CB-CA-C	-5.57	99.62	110.20

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1956	0	1925	38	0
1	B	1943	0	1916	36	0
2	A	2	0	0	0	0
2	B	3	0	0	1	0
All	All	3904	0	3841	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:617:TRP:CD1	1:A:619:LEU:O	2.36	0.78
1:A:617:TRP:HD1	1:A:619:LEU:O	1.74	0.69
1:B:617:TRP:CD2	1:B:624:ARG:HG3	2.29	0.67
1:A:555:SER:HA	1:A:558:GLN:HG3	1.80	0.64
1:A:491:LEU:HB3	1:A:534:HIS:CD2	2.33	0.64
1:A:588:SER:HB2	1:A:627:LEU:HD21	1.80	0.63
1:A:555:SER:HA	1:A:558:GLN:HE21	1.64	0.63
1:B:502:LYS:HG3	1:B:543:LEU:HD13	1.81	0.62
1:B:614:GLU:H	1:B:614:GLU:CD	2.02	0.61
1:A:515:LYS:HE3	1:A:560:LYS:NZ	2.16	0.61
1:B:541:LEU:HD21	1:B:578:LEU:HD23	1.84	0.59
1:B:614:GLU:HG3	1:B:624:ARG:NH2	2.18	0.58
1:B:649:ARG:NH2	2:B:801:HOH:O	1.98	0.57
1:A:545:MET:SD	1:A:583:PHE:HA	2.44	0.57
1:A:515:LYS:HE3	1:A:560:LYS:HZ1	1.70	0.57
1:A:602:ALA:O	1:A:606:ILE:HG13	2.05	0.56
1:A:631:LEU:O	1:A:635:ILE:HG13	2.07	0.55
1:B:491:LEU:HD13	1:B:534:HIS:ND1	2.23	0.54
1:B:636:LEU:HD21	1:B:668:LEU:HD21	1.90	0.54
1:A:545:MET:SD	1:A:583:PHE:HD1	2.31	0.53
1:A:558:GLN:O	1:A:562:LEU:HD12	2.09	0.53
1:A:491:LEU:HD22	1:A:534:HIS:CG	2.44	0.53
1:B:567:ASN:OD1	1:A:479:CYS:HB3	2.08	0.53
1:B:491:LEU:HD13	1:B:534:HIS:CE1	2.44	0.52
1:B:690:ILE:HD11	1:B:722:LEU:HG	1.92	0.51
1:B:503:GLN:O	1:B:507:GLN:HG3	2.11	0.51
1:B:588:SER:HB3	1:B:627:LEU:HD21	1.93	0.50
1:A:712:GLN:O	1:A:716:VAL:HG13	2.11	0.50
1:A:528:SER:OG	1:A:531:THR:HG23	2.13	0.49
1:A:606:ILE:HG22	1:A:610:ILE:HD12	1.96	0.48
1:B:504:LYS:HG3	1:B:514:LEU:HB2	1.94	0.47
1:B:525:THR:HG22	1:B:531:THR:HG22	1.96	0.47
1:A:519:SER:OG	1:A:560:LYS:HE2	2.13	0.47
1:B:510:VAL:HG11	1:B:554:GLU:HG3	1.95	0.47
1:A:687:SER:O	1:A:691:GLU:HG3	2.15	0.47
1:B:502:LYS:HG3	1:B:543:LEU:CD1	2.43	0.46
1:B:580:TRP:N	1:B:619:LEU:HD11	2.31	0.46
1:A:701:ILE:HG23	1:A:707:THR:HG21	1.98	0.46
1:A:518:LEU:HD23	1:A:518:LEU:HA	1.76	0.45
1:B:542:GLU:H	1:B:542:GLU:CD	2.18	0.45
1:B:519:SER:OG	1:B:560:LYS:HD3	2.16	0.45
1:B:607:ALA:HA	1:B:666:VAL:HG22	1.99	0.45

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:660:CYS:HB3	1:A:663:THR:OG1	2.16	0.45
1:B:487:LEU:H	1:B:487:LEU:HG	1.40	0.45
1:A:491:LEU:HB3	1:A:534:HIS:NE2	2.31	0.45
1:A:657:LEU:HB3	1:A:666:VAL:HG12	1.99	0.44
1:A:690:ILE:HG12	1:A:695:LEU:HD11	2.00	0.44
1:B:614:GLU:HG3	1:B:624:ARG:CZ	2.47	0.44
1:B:649:ARG:HG2	1:A:480:PHE:CE1	2.53	0.44
1:B:597:GLU:H	1:B:597:GLU:CD	2.21	0.43
1:A:641:PRO:HG2	1:A:676:HIS:CE1	2.53	0.43
1:B:570:GLU:HG2	1:B:608:HIS:CE1	2.53	0.43
1:A:617:TRP:CD2	1:A:624:ARG:HG3	2.54	0.43
1:A:718:ILE:O	1:A:722:LEU:HD12	2.19	0.42
1:A:497:LEU:HD22	1:A:517:THR:HG23	2.02	0.42
1:B:617:TRP:NE1	1:B:620:SER:O	2.51	0.42
1:B:491:LEU:HD13	1:B:534:HIS:CG	2.54	0.42
1:A:645:MET:H	1:A:645:MET:HG2	1.65	0.42
1:B:499:GLN:O	1:B:503:GLN:HG3	2.20	0.42
1:B:677:VAL:HB	1:B:685:TYR:CD2	2.55	0.42
1:B:610:ILE:HD13	1:B:628:LEU:HD13	2.02	0.42
1:A:536:ILE:HD13	1:A:536:ILE:HA	1.78	0.42
1:B:697:HIS:O	1:B:701:ILE:HG12	2.20	0.41
1:A:710:HIS:O	1:A:714:ILE:HG12	2.20	0.41
1:B:523:ASN:HB3	1:A:486:GLN:OE1	2.21	0.41
1:A:635:ILE:HD12	1:A:668:LEU:HD23	2.02	0.41
1:B:617:TRP:CE2	1:B:624:ARG:HG3	2.55	0.41
1:B:668:LEU:HD12	1:B:668:LEU:HA	1.82	0.40
1:A:558:GLN:OE1	1:A:590:LEU:HD21	2.21	0.40
1:B:535:PHE:CD2	1:B:568:ILE:HD11	2.56	0.40
1:A:597:GLU:HG3	1:A:645:MET:HE2	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	240/250 (96%)	227 (95%)	13 (5%)	0	100	100
1	B	237/250 (95%)	227 (96%)	10 (4%)	0	100	100
All	All	477/500 (95%)	454 (95%)	23 (5%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	220/227 (97%)	205 (93%)	15 (7%)	16	40
1	B	219/227 (96%)	205 (94%)	14 (6%)	17	43
All	All	439/454 (97%)	410 (93%)	29 (7%)	16	41

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	487	LEU
1	B	508	ASN
1	B	510	VAL
1	B	538	ASN
1	B	539	GLN
1	B	556	SER
1	B	566	ASN
1	B	582	ASP
1	B	588	SER
1	B	620	SER
1	B	626	SER
1	B	649	ARG
1	B	707	THR
1	B	722	LEU
1	A	508	ASN
1	A	510	VAL
1	A	511	ASP
1	A	512	THR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	576	SER
1	A	615	GLN
1	A	621	ARG
1	A	626	SER
1	A	633	SER
1	A	649	ARG
1	A	662	THR
1	A	664	PRO
1	A	687	SER
1	A	721	SER
1	A	722	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	523	ASN
1	A	558	GLN
1	A	652	ASN
1	A	696	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	244/250 (97%)	-0.02	0 <span>100</span> <span>100</span>	45, 74, 102, 129	0
1	B	243/250 (97%)	0.20	9 (3%) <span>41</span> <span>37</span>	45, 71, 111, 152	0
All	All	487/500 (97%)	0.09	9 (1%) <span>68</span> <span>67</span>	45, 72, 107, 152	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	723	GLU	4.5
1	B	706	HIS	3.8
1	B	707	THR	3.1
1	B	722	LEU	3.1
1	B	719	LEU	2.9
1	B	485	ALA	2.5
1	B	701	ILE	2.4
1	B	483	VAL	2.3
1	B	486	GLN	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.