



Full wwPDB X-ray Structure Validation Report ⓘ

May 18, 2020 – 09:36 pm BST

PDB ID : 1XZ8
Title : Pyrr, The Regulator Of The Pyrimidine Biosynthetic Operon In Bacillus caldolyticus, Nucleotide-bound form
Authors : Chander, P.; Halbig, K.M.; Miller, J.K.; Fields, C.J.; Bonner, H.K.; Grabner, G.K.; Switzer, R.L.; Smith, J.L.
Deposited on : 2004-11-11
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	179	<div> <div>4%</div> <div> <div></div> <div>49%</div> <div>39%</div> <div>• 9%</div> </div> </div>
1	B	179	<div> <div>3%</div> <div> <div></div> <div>54%</div> <div>35%</div> <div>• 10%</div> </div> </div>

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 2654 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called PyrR bifunctional protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	162	Total	C	N	O	S	0	0	0
			1268	793	235	235	5			
1	B	161	Total	C	N	O	S	0	0	0
			1260	789	234	232	5			

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mg	0	0
			1	1		
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is GUANOSINE-5'-MONOPHOSPHATE (three-letter code: 5GP) (formula: C₁₀H₁₄N₅O₈P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			24	10	5	8	1		

- Molecule 4 is GUANOSINE-3'-MONOPHOSPHATE (three-letter code: 3GP) (formula: $C_{10}H_{14}N_5O_8P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			24	10	5	8	1		

- Molecule 5 is URIDINE-5'-MONOPHOSPHATE (three-letter code: U5P) (formula: $C_9H_{13}N_2O_9P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	B	1	Total	C	N	O	P	0	0
			21	9	2	9	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	28	Total	O	0	0
			28	28		
6	B	27	Total	O	0	0
			27	27		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: PyrR bifunctional protein



• Molecule 1: PyrR bifunctional protein



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	128.54Å 128.54Å 128.76Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.80 42.11 – 2.80	Depositor EDS
% Data completeness (in resolution range)	(Not available) (50.00-2.80) 99.7 (42.11-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.05 (at 2.81Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.220 , 0.300 0.208 , 0.290	Depositor DCC
R_{free} test set	1053 reflections (10.31%)	wwPDB-VP
Wilson B-factor (Å ²)	48.2	Xtriage
Anisotropy	0.102	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 50.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	2654	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 3GP, MG, 5GP, U5P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.36	0/1280	0.64	0/1727
1	B	0.38	0/1272	0.64	0/1716
All	All	0.37	0/2552	0.64	0/3443

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	4	ALA	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1268	0	1324	68	0
1	B	1260	0	1320	51	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	24	0	12	6	0
4	A	24	0	12	3	0
5	B	21	0	10	2	0
6	A	28	0	0	3	0
6	B	27	0	0	2	0
All	All	2654	0	2678	119	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

All (119) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:3:LYS:HE2	1:A:178:GLU:HB2	1.44	0.95
1:B:99:ASN:HD21	1:B:130:GLN:NE2	1.78	0.81
1:A:32:GLY:HA2	1:A:98:ARG:HE	1.48	0.78
1:B:30:ILE:HG21	1:B:53:ILE:HG21	1.66	0.78
1:B:7:MET:HB2	1:B:174:VAL:HB	1.65	0.77
1:B:69:THR:HG23	1:B:72:ARG:CZ	2.18	0.74
3:A:200:5GP:H2'	4:A:300:3GP:O2'	1.89	0.73
1:A:112:THR:OG1	3:A:200:5GP:H5'1	1.89	0.72
1:B:2:GLN:HG3	1:B:177:HIS:CE1	2.27	0.69
1:B:108:PHE:HB3	5:B:400:U5P:O3P	1.97	0.65
1:A:124:GLY:O	1:A:126:PRO:HD3	1.96	0.65
1:B:121:MET:HE2	1:B:129:ILE:HD12	1.77	0.65
1:A:43:ILE:O	1:A:47:ARG:HG3	1.97	0.65
1:A:179:LYS:HA	1:A:179:LYS:HE3	1.78	0.64
1:B:24:ILE:HD13	1:B:57:GLU:HG2	1.80	0.64
1:A:39:LYS:O	1:A:43:ILE:HG13	1.97	0.63
3:A:200:5GP:N3	4:A:300:3GP:O2'	2.31	0.60
1:B:10:GLN:HG3	1:B:13:ARG:HH12	1.64	0.60
1:B:98:ARG:HH11	1:B:98:ARG:HG2	1.66	0.60
1:A:96:THR:O	1:A:98:ARG:HG2	2.02	0.59
1:B:51:GLU:O	1:B:55:GLN:HG3	2.02	0.59
1:A:3:LYS:CE	1:A:178:GLU:HB2	2.27	0.59
1:B:35:LEU:HD23	1:B:101:ILE:HB	1.84	0.59
1:A:104:ASP:O	1:A:133:VAL:HA	2.03	0.58
1:A:12:ILE:HD12	1:A:172:ASP:HB3	1.84	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:112:THR:HB	6:A:326:HOH:O	2.04	0.58
1:A:41:ARG:NH1	4:A:300:3GP:H5'1	2.19	0.58
1:B:69:THR:HG23	1:B:72:ARG:NE	2.19	0.58
1:A:54:GLU:OE1	1:A:60:SER:HB3	2.03	0.57
1:A:94:PRO:HB3	1:A:98:ARG:NH1	2.20	0.57
1:B:30:ILE:HG21	1:B:53:ILE:CG2	2.34	0.57
1:A:34:VAL:HG21	1:A:94:PRO:HB2	1.85	0.57
1:B:70:LEU:HD13	1:B:70:LEU:O	2.05	0.56
1:A:95:VAL:HG13	1:A:126:PRO:HG3	1.88	0.56
1:A:43:ILE:HD13	1:A:65:GLU:HB3	1.88	0.56
1:B:69:THR:HG23	1:B:72:ARG:NH2	2.20	0.56
1:A:127:ALA:O	1:A:128:ARG:HB3	2.05	0.55
1:B:99:ASN:ND2	1:B:128:ARG:HD2	2.21	0.55
1:A:1:MET:O	1:A:1:MET:HG3	2.07	0.55
1:A:3:LYS:HE2	1:A:178:GLU:CB	2.28	0.54
1:B:179:LYS:HB3	6:B:421:HOH:O	2.06	0.54
1:B:14:ARG:HG2	1:B:14:ARG:HH11	1.72	0.54
1:A:153:VAL:O	1:A:155:THR:HG23	2.08	0.54
1:B:99:ASN:HD21	1:B:130:GLN:HE21	1.52	0.54
1:B:96:THR:HA	1:B:125:ARG:O	2.08	0.54
1:A:96:THR:HA	1:A:125:ARG:O	2.07	0.54
1:B:104:ASP:O	1:B:133:VAL:HA	2.07	0.54
1:A:108:PHE:CE2	1:A:139:HIS:HB2	2.43	0.53
1:B:153:VAL:O	1:B:155:THR:HG23	2.09	0.53
1:A:145:ARG:HD2	1:B:145:ARG:HB3	1.90	0.52
1:B:99:ASN:ND2	1:B:130:GLN:NE2	2.55	0.51
1:A:112:THR:CB	3:A:200:5GP:H5'1	2.39	0.51
1:A:145:ARG:HB3	1:B:145:ARG:HG2	1.92	0.50
1:A:24:ILE:HD13	1:A:57:GLU:HG2	1.93	0.50
1:B:41:ARG:HG3	1:B:41:ARG:HH11	1.76	0.50
1:B:43:ILE:CD1	1:B:65:GLU:HB3	2.41	0.50
1:B:17:THR:HA	1:B:52:ARG:HD3	1.93	0.50
1:B:43:ILE:HD11	1:B:65:GLU:HB3	1.94	0.50
1:A:109:THR:HG23	3:A:200:5GP:O2P	2.11	0.50
1:A:72:ARG:N	1:A:72:ARG:HD2	2.27	0.50
1:B:99:ASN:HD21	1:B:130:GLN:HE22	1.56	0.50
1:A:107:LEU:HB2	1:A:133:VAL:HG11	1.95	0.49
1:A:7:MET:HB2	1:A:174:VAL:HB	1.94	0.49
1:B:91:VAL:N	1:B:92:PRO:HD3	2.28	0.49
1:A:12:ILE:CD1	1:A:172:ASP:HB3	2.43	0.49
1:A:130:GLN:HE21	1:A:148:PHE:HE1	1.60	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:113:VAL:HG23	6:A:326:HOH:O	2.13	0.48
1:B:99:ASN:ND2	1:B:130:GLN:HE21	2.11	0.48
1:B:156:SER:OG	1:B:159:GLU:HG3	2.13	0.48
1:A:69:THR:C	1:A:71:TYR:H	2.16	0.48
1:B:125:ARG:NH1	1:B:125:ARG:HB2	2.28	0.48
1:B:39:LYS:O	1:B:43:ILE:HG13	2.14	0.48
1:A:104:ASP:OD2	1:A:105:ASP:N	2.47	0.48
1:A:48:ARG:O	1:A:51:GLU:HB3	2.13	0.48
1:A:108:PHE:HB3	3:A:200:5GP:O2P	2.13	0.48
1:B:4:ALA:O	1:B:175:SER:HB2	2.14	0.48
1:A:114:ARG:HG2	1:A:114:ARG:HH11	1.78	0.47
1:A:49:LEU:O	1:A:53:ILE:HG13	2.13	0.47
1:A:108:PHE:HZ	1:A:157:ARG:NH1	2.13	0.47
1:A:21:HIS:CE1	1:A:56:ILE:HD13	2.49	0.47
1:B:157:ARG:HD3	6:B:419:HOH:O	2.14	0.47
1:A:52:ARG:HG3	1:A:52:ARG:HH11	1.80	0.47
1:B:160:LEU:HD22	5:B:400:U5P:O4	2.14	0.47
1:A:109:THR:OG1	1:A:111:ARG:HG3	2.15	0.46
1:B:41:ARG:HG3	1:B:41:ARG:NH1	2.30	0.46
1:B:27:ASN:OD1	1:B:99:ASN:ND2	2.49	0.46
1:B:10:GLN:HG3	1:B:13:ARG:NH1	2.30	0.46
1:A:97:GLU:HA	1:A:127:ALA:HB2	1.97	0.46
1:B:24:ILE:HG12	1:B:30:ILE:HG23	1.98	0.45
1:A:1:MET:O	1:A:2:GLN:C	2.52	0.45
1:A:51:GLU:HG2	1:A:55:GLN:HE21	1.81	0.45
1:A:43:ILE:CD1	1:A:65:GLU:HB3	2.46	0.45
1:A:114:ARG:HG2	1:A:114:ARG:NH1	2.31	0.45
1:B:12:ILE:O	1:B:16:LEU:HG	2.17	0.45
1:A:108:PHE:CZ	1:A:157:ARG:NH1	2.85	0.45
1:A:156:SER:OG	1:A:159:GLU:HG3	2.17	0.44
1:A:178:GLU:HG3	1:A:179:LYS:N	2.32	0.44
1:A:30:ILE:HG21	1:A:53:ILE:HG21	2.00	0.44
1:A:95:VAL:HG12	1:A:96:THR:N	2.33	0.43
1:A:71:TYR:HE1	1:A:73:ASP:HB3	1.84	0.43
1:A:131:LEU:HB3	1:A:146:ALA:HA	2.00	0.43
1:A:68:ILE:HG22	1:A:68:ILE:O	2.18	0.43
1:B:22:GLU:HG2	1:B:26:ARG:NH2	2.34	0.43
1:A:130:GLN:NE2	1:A:148:PHE:CE1	2.87	0.42
1:A:27:ASN:OD1	1:A:128:ARG:NH2	2.46	0.42
1:B:104:ASP:OD2	1:B:105:ASP:N	2.52	0.42
1:A:100:VAL:O	1:A:129:ILE:HA	2.20	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:41:ARG:HB3	1:B:165:LEU:HD12	2.02	0.42
1:B:93:PHE:HA	1:B:94:PRO:HD3	1.87	0.42
1:A:67:ASP:O	1:A:69:THR:N	2.52	0.42
1:B:44:TYR:O	1:B:48:ARG:HG3	2.19	0.42
1:A:91:VAL:HG13	6:A:301:HOH:O	2.19	0.42
1:B:102:LEU:N	1:B:130:GLN:O	2.51	0.41
1:A:130:GLN:NE2	1:A:148:PHE:HE1	2.18	0.41
1:B:96:THR:O	1:B:97:GLU:HB2	2.20	0.41
1:A:145:ARG:HH11	1:A:145:ARG:HG2	1.86	0.41
1:A:1:MET:O	1:A:1:MET:CG	2.70	0.40
1:A:52:ARG:NH1	1:A:52:ARG:HG3	2.36	0.40
1:A:37:GLY:HA3	1:A:43:ILE:HA	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	158/179 (88%)	146 (92%)	8 (5%)	4 (2%)	5	19
1	B	157/179 (88%)	142 (90%)	14 (9%)	1 (1%)	25	56
All	All	315/358 (88%)	288 (91%)	22 (7%)	5 (2%)	9	31

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	68	ILE
1	A	2	GLN
1	A	128	ARG
1	B	119	ALA
1	A	95	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	138/154 (90%)	132 (96%)	6 (4%)	29	62
1	B	137/154 (89%)	133 (97%)	4 (3%)	42	76
All	All	275/308 (89%)	265 (96%)	10 (4%)	35	69

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	ARG
1	A	31	ASP
1	A	123	LEU
1	A	135	VAL
1	A	145	ARG
1	A	179	LYS
1	B	123	LEU
1	B	135	VAL
1	B	140	ARG
1	B	145	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	2	GLN
1	A	10	GLN
1	A	55	GLN
1	A	130	GLN
1	A	177	HIS
1	B	130	GLN
1	B	177	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
5	U5P	B	400	2	18,22,22	2.68	5 (27%)	21,33,33	1.76	3 (14%)
4	3GP	A	300	2	22,26,26	2.52	7 (31%)	26,40,40	2.98	8 (30%)
3	5GP	A	200	-	22,26,26	1.65	3 (13%)	27,40,40	2.31	8 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	U5P	B	400	2	-	3/8/26/26	0/2/2/2
4	3GP	A	300	2	-	1/7/27/27	0/3/3/3
3	5GP	A	200	-	-	2/6/26/26	0/3/3/3

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	400	U5P	C6-N1	7.86	1.45	1.35
4	A	300	3GP	P-O1P	6.93	1.72	1.50
4	A	300	3GP	P-O2P	-5.92	1.32	1.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	400	U5P	C4-N3	5.82	1.43	1.33
3	A	200	5GP	C6-N1	5.00	1.41	1.33
5	B	400	U5P	P-O3P	3.21	1.67	1.54
4	A	300	3GP	C6-N1	3.20	1.38	1.33
4	A	300	3GP	P-O3'	2.86	1.64	1.59
4	A	300	3GP	O4'-C1'	2.55	1.44	1.41
4	A	300	3GP	C2'-C1'	-2.55	1.49	1.53
5	B	400	U5P	C6-C5	-2.54	1.32	1.38
4	A	300	3GP	C2-N1	2.28	1.39	1.35
5	B	400	U5P	O2'-C2'	2.20	1.48	1.43
3	A	200	5GP	C8-N7	-2.17	1.30	1.34
3	A	200	5GP	P-O5'	-2.11	1.53	1.60

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	300	3GP	C5-C6-N1	-8.73	111.50	123.43
3	A	200	5GP	C5-C6-N1	-8.36	112.00	123.43
4	A	300	3GP	O4'-C4'-C5'	6.56	123.38	109.21
4	A	300	3GP	C6-N1-C2	5.70	124.98	115.93
3	A	200	5GP	C6-N1-C2	5.10	124.04	115.93
4	A	300	3GP	O3P-P-O1P	-4.16	94.40	110.68
5	B	400	U5P	O5'-P-O1P	4.10	117.98	106.47
5	B	400	U5P	C5-C4-N3	-3.82	114.90	123.31
4	A	300	3GP	O2P-P-O1P	3.66	125.02	110.68
4	A	300	3GP	C2-N3-C4	-3.02	111.91	115.36
3	A	200	5GP	C2-N3-C4	-2.74	112.22	115.36
3	A	200	5GP	O4'-C4'-C5'	-2.60	100.81	109.37
3	A	200	5GP	N2-C2-N3	2.54	121.93	117.79
4	A	300	3GP	C4-C5-N7	2.44	111.94	109.40
3	A	200	5GP	N2-C2-N1	-2.35	113.59	117.25
3	A	200	5GP	C4-C5-N7	2.31	111.81	109.40
4	A	300	3GP	C6-C5-C4	-2.16	118.73	120.80
5	B	400	U5P	O3P-P-O5'	-2.10	101.15	106.73
3	A	200	5GP	N3-C2-N1	-2.06	124.47	127.22

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	400	U5P	O4'-C1'-N1-C6
3	A	200	5GP	O4'-C4'-C5'-O5'

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	200	5GP	C3'-C4'-C5'-O5'
4	A	300	3GP	C3'-O3'-P-O1P
5	B	400	U5P	O4'-C4'-C5'-O5'
5	B	400	U5P	C3'-C4'-C5'-O5'


There are no ring outliers.

3 monomers are involved in 9 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	400	U5P	2	0
4	A	300	3GP	3	0
3	A	200	5GP	6	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand U5P B 400



Ligand 3GP A 300





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	162/179 (90%)	0.14	7 (4%) 35 25	22, 37, 67, 99	0
1	B	161/179 (89%)	0.12	5 (3%) 49 39	21, 36, 61, 90	0
All	All	323/358 (90%)	0.13	12 (3%) 41 31	21, 37, 65, 99	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	73	ASP	6.9
1	B	69	THR	5.8
1	A	69	THR	5.4
1	B	71	TYR	4.6
1	B	70	LEU	4.0
1	B	4	ALA	2.8
1	A	70	LEU	2.8
1	A	72	ARG	2.7
1	A	157	ARG	2.6
1	B	1	MET	2.2
1	A	71	TYR	2.1
1	A	178	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MG	A	180	1/1	0.72	0.16	43,43,43,43	0
5	U5P	B	400	21/21	0.82	0.29	60,76,79,81	0
4	3GP	A	300	24/24	0.87	0.28	40,46,56,58	0
2	MG	B	180	1/1	0.89	0.15	43,43,43,43	0
3	5GP	A	200	24/24	0.90	0.27	43,46,55,58	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around U5P B 400:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 3GP A 300:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 5GP A 200:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.