



Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 07:10 am BST

PDB ID : 4XZA
Title : The crystal structure of Erve virus nucleoprotein
Authors : Guo, Y.; Wang, W.; Liu, X.; Wang, X.; Wang, J.; Huo, T.; Liu, B.
Deposited on : 2015-02-04
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	482	<div> <div>2%</div> <div> <div></div> <div>65%</div> <div>10%</div> <div>24%</div> </div> </div>
1	B	482	<div> <div>2%</div> <div> <div></div> <div>70%</div> <div>5%</div> <div>24%</div> </div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6429 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nucleoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	366	Total	C	N	O	S	0	0	0
			2884	1840	494	535	15			
1	B	365	Total	C	N	O	S	0	0	0
			2870	1828	493	534	15			

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	337	Total	O	0	0
			337	337		
2	B	338	Total	O	0	0
			338	338		

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Chain A:

2% 65% 10% 24%

PRO LYS VAL GLN ALA ASP PHE GLY GLU TRP GLY ALA ASP ASP GLN SER ARG VAL GLN ASP VAL ILE GLU ILE HIS GLN VAL LEU LEU LYS SER ASP ILE ILE ALA GLU THR THR ALA THR THR SER SER LEU LEU GLY GLY ALA F291 F292 K293 N294 Q295 I314 I322 K341 K345 R345 VAL

H1 E2 D6 F7 S8 GLY ARG D11 D14 R15 R18 F21 P22 D23 V24 Y33 E47 Q48 R51 Q52 A53 K54 L67 T71 K36 E87 K91 S102 G142 Y143 Q145 L146 E147 E148 T149 Q155 V156 I157 T169 R181 G191

Chain B:

Residue	Amino Acid
M1	MET
G9	ASN
ARG	ASN
D14	GLN
D14	SER
R18	ALA
R18	LEU
R18	ILE
S27	ASN
S27	PRO
Y33	ALA
Y33	TRP
Y33	GLY
Q48	ASP
A49	ILE
A50	ASP
R51	LYS
Q52	LYS
A53	ASN
A53	LYS
T71	ASN
T71	GLY
E87	LEU
E87	MET
N113	LEU
N113	LEU
Q116	THR
Q116	THR
K119	GLY
K119	ILE
E148	ALA
E148	LYS
S154	LEU
Q155	ARG
V156	GLU
V156	LEU
Q159	TYR
Q159	GLY
R180	PRO
R181	ALA
R181	ALA
GLY	MET
GLY	VAL
GLY	VAL
PRO	LYS
PRO	VAL
LYS	GLN
ARG	GLN
GLY	ALA
VAL	ALA
VAL	ASP
GLU	LYS
HIS	PHE
VAL	GLY
GLU	GLU
TRP	TRP
GLY	GLY
LYS	LYS
VAL	ALA
ASP	PRO

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	46.08Å 104.29Å 70.54Å 90.00° 92.38° 90.00°	Depositor
Resolution (Å)	46.04 – 1.80 46.04 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.5 (46.04-1.80) 96.6 (46.04-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.41 (at 1.79Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.168 , 0.202 0.171 , 0.202	Depositor DCC
R_{free} test set	1992 reflections (3.25%)	wwPDB-VP
Wilson B-factor (Å ²)	23.4	Xtriage
Anisotropy	0.318	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 39.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.036 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6429	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/2943	0.50	0/3978
1	B	0.37	0/2928	0.49	0/3957
All	All	0.36	0/5871	0.50	0/7935

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2884	0	2884	37	1
1	B	2870	0	2869	23	0
2	A	337	0	0	13	6
2	B	338	0	0	12	5
All	All	6429	0	5753	58	6

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:365:MET:SD	2:B:782:HOH:O	2.14	1.03

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:345:ARG:HD3	2:A:509:HOH:O	1.65	0.93
1:A:416:MET:SD	2:A:579:HOH:O	2.28	0.90
1:A:295:GLN:OE1	2:A:666:HOH:O	1.88	0.89
1:A:147:GLU:OE2	2:A:501:HOH:O	1.93	0.86
1:B:181:ARG:O	2:B:806:HOH:O	1.93	0.85
1:A:341:LYS:NZ	2:A:502:HOH:O	2.11	0.83
1:A:291:VAL:O	2:A:789:HOH:O	1.99	0.81
1:A:87:GLU:OE1	2:A:836:HOH:O	2.01	0.79
1:B:18:ARG:NH2	2:B:503:HOH:O	2.21	0.73
1:B:27:SER:OG	2:B:501:HOH:O	1.99	0.71
1:B:87:GLU:OE2	2:B:695:HOH:O	2.08	0.70
1:A:314:ILE:HD13	1:A:322:LEU:HD22	1.76	0.66
1:B:294:ASN:O	2:B:781:HOH:O	2.14	0.66
1:A:11:ASP:N	2:A:816:HOH:O	2.29	0.65
1:A:155:GLN:HG3	1:A:466:LYS:HB2	1.80	0.64
1:B:181:ARG:HD2	2:B:833:HOH:O	1.98	0.63
1:B:14:ASP:OD2	1:B:18:ARG:NH1	2.33	0.62
1:A:396:GLU:CD	1:A:396:GLU:H	2.04	0.61
2:A:801:HOH:O	1:B:365:MET:HE1	2.01	0.60
1:A:21:PHE:HB3	1:A:24:VAL:HG13	1.83	0.59
1:B:156:VAL:HG11	1:B:469:ALA:HA	1.85	0.57
1:B:155:GLN:HG2	2:B:772:HOH:O	2.05	0.56
1:B:156:VAL:O	2:B:772:HOH:O	2.18	0.56
1:B:1:MET:O	2:B:760:HOH:O	2.18	0.55
1:A:54:LYS:HG2	2:A:832:HOH:O	2.06	0.55
1:A:396:GLU:OE2	2:A:503:HOH:O	2.18	0.54
1:B:294:ASN:HB3	1:B:463:PHE:O	2.07	0.53
1:A:47:GLU:HB2	2:A:809:HOH:O	2.08	0.53
1:A:181:ARG:NH1	1:B:368:ILE:O	2.42	0.52
1:A:345:ARG:NH2	1:A:348:LEU:HG	2.24	0.52
1:B:71:THR:HB	1:B:407:ASN:HB2	1.91	0.52
1:B:155:GLN:HG3	1:B:466:LYS:O	2.10	0.51
1:A:15:ARG:HG2	1:A:18:ARG:NH2	2.26	0.51
1:A:169:THR:HG21	1:A:424:VAL:HG13	1.92	0.50
1:B:154:SER:OG	1:B:155:GLN:N	2.46	0.49
1:A:71:THR:HB	1:A:407:ASN:HB2	1.96	0.48
1:B:349:LYS:C	2:B:779:HOH:O	2.52	0.48
1:A:345:ARG:CD	2:A:509:HOH:O	2.40	0.47
1:B:33:TYR:CG	1:B:148:GLU:HG3	2.50	0.47
1:A:6:ASP:HB2	1:A:86:LYS:HD3	1.97	0.47
1:B:1:MET:HE2	1:B:393:SER:HB2	1.96	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:449:HIS:HD2	1:A:450:GLN:NE2	2.11	0.47
1:A:48:GLN:O	1:A:52:GLN:HG2	2.14	0.47
1:A:369:HIS:CE1	1:B:368:ILE:HD11	2.51	0.46
1:B:113:ASN:OD1	1:B:116:GLN:NE2	2.49	0.45
1:A:33:TYR:CG	1:A:148:GLU:HG3	2.53	0.44
1:A:143:TYR:CE2	1:A:149:THR:HB	2.52	0.44
1:A:449:HIS:HD2	1:A:450:GLN:HE21	1.67	0.42
1:A:67:LEU:HD21	1:A:442:VAL:HB	2.02	0.42
1:A:142:GLY:O	1:A:145:GLN:HG3	2.19	0.42
1:A:156:VAL:HG11	1:A:469:ALA:HA	2.02	0.42
1:A:15:ARG:HG2	1:A:18:ARG:HH21	1.85	0.42
1:B:119:LYS:HE2	2:B:747:HOH:O	2.18	0.41
1:A:14:ASP:HB3	1:A:18:ARG:NH1	2.35	0.41
1:A:87:GLU:O	1:A:91:LYS:HG2	2.20	0.41
1:A:345:ARG:CZ	1:A:348:LEU:HG	2.50	0.40
1:A:371:HIS:HB3	1:A:374:ILE:HG12	2.04	0.40

All (6) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:560:HOH:O	2:B:602:HOH:O[2_-1648]	1.90	0.30
1:A:345:ARG:NH1	2:A:501:HOH:O[1_-455]	2.02	0.18
2:A:542:HOH:O	2:B:572:HOH:O[2_-1748]	2.12	0.08
2:A:580:HOH:O	2:B:583:HOH:O[2_-1647]	2.12	0.08
2:A:717:HOH:O	2:B:611:HOH:O[2_-1747]	2.13	0.07
2:A:596:HOH:O	2:B:612:HOH:O[2_-1648]	2.18	0.02

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	360/482 (75%)	350 (97%)	9 (2%)	1 (0%)	41	27
1	B	359/482 (74%)	351 (98%)	8 (2%)	0	100	100
All	All	719/964 (75%)	701 (98%)	17 (2%)	1 (0%)	51	36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	348	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	314/404 (78%)	308 (98%)	6 (2%)	57	46
1	B	312/404 (77%)	310 (99%)	2 (1%)	86	84
All	All	626/808 (78%)	618 (99%)	8 (1%)	69	62

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	GLU
1	A	23	ASP
1	A	102	SER
1	A	291	VAL
1	A	352	LYS
1	A	360	ASP
1	B	352	LYS
1	B	360	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	366/482 (75%)	-0.21	12 (3%) 46 40	17, 27, 45, 61	0
1	B	365/482 (75%)	-0.14	10 (2%) 54 49	16, 26, 48, 63	0
All	All	731/964 (75%)	-0.17	22 (3%) 50 44	16, 26, 48, 63	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	51	ARG	4.5
1	A	1	MET	3.7
1	B	9	GLY	3.4
1	A	8	SER	3.4
1	B	433	SER	3.1
1	B	48	GLN	3.0
1	B	1	MET	2.8
1	B	53	ALA	2.8
1	A	346	ILE	2.6
1	A	294	ASN	2.6
1	A	51	ARG	2.5
1	A	48	GLN	2.4
1	B	477	PHE	2.4
1	B	159	GLN	2.4
1	B	50	ALA	2.3
1	A	292	PHE	2.3
1	B	180	ARG	2.3
1	A	291	VAL	2.3
1	A	347	GLY	2.2
1	A	293	LYS	2.2
1	A	53	ALA	2.0
1	A	157	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.