



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 13, 2020 – 10:03 AM BST

PDB ID : 1Y42
Title : Crystal structure of a C-terminally truncated CYT-18 protein
Authors : Paukstelis, P.J.; Coon, R.; Madabusi, L.; Nowakowski, J.; Monzingo, A.; Robertus, J.; Lambowitz, A.M.
Deposited on : 2004-11-29
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.14.4.dev1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.14.4.dev1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	X	392	<div> <div>8%</div> <div>80%</div> <div>13%</div> <div>6%</div> </div>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3161 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Tyrosyl-tRNA synthetase, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	X	370	Total	C	N	O	S	0	0	0
			2950	1877	514	544	15			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
X	32	MET	-	INITIATING METHIONINE	UNP P12063

- Molecule 2 is TYROSINE (three-letter code: TYR) (formula: $C_9H_{11}NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	X	1	Total	C	N	O	0	0
			13	9	1	3		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	X	198	Total 198	O 198	0	0

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Chain X:
-
- 8% 80% 13% 6%
- MET LEU ARG ARG GLU PHE GLY P39 K40 Y41 T42 N66 T67 K84 P85 H86 G101 I102 D103 A106 P107 S108 L109 L115 W122 G135 T138 I141 G142 D143 P144 T145 G146 ARG LEU LYS SER SER ASP HIS LEU SER SER SER D158 Y168 Q169 Q180 R184 Y186 L209 H217 A218 L219 R220 S226 R227 D228 T229 V230 K231 N232 R233 K234 T235 Q236 G237 D238 G239 V240 A243 Q251 K266 Q267 Y274 I278 D293 P294 Y299 D307 V310 L317 T318 D319 F325 S328 A329 N331 A332 I333 D336 F337 Y338 S353 D354 Q355 E356 F366 D382 A393 R394 G419 GLN MET THR ILE

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	104.88Å 73.21Å 56.79Å 90.00° 111.35° 90.00°	Depositor
Resolution (Å)	20.00 – 1.95 19.78 – 1.95	Depositor EDS
% Data completeness (in resolution range)	95.0 (20.00-1.95) 95.0 (19.78-1.95)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.93 (at 1.94Å)	Xtriage
Refinement program	REFMAC 5.2.0003	Depositor
R, R_{free}	0.174 , 0.235 0.182 , 0.236	Depositor DCC
R_{free} test set	1936 reflections (6.97%)	wwPDB-VP
Wilson B-factor (Å ²)	22.2	Xtriage
Anisotropy	0.197	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 51.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3161	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	X	0.88	0/3023	0.92	11/4089 (0.3%)

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	X	319	ASP	CB-CG-OD2	7.56	125.10	118.30
1	X	336	ASP	CB-CG-OD2	7.24	124.81	118.30
1	X	158	ASP	CB-CG-OD2	6.76	124.39	118.30
1	X	184	ARG	NE-CZ-NH1	6.64	123.62	120.30
1	X	382	ASP	CB-CG-OD2	6.51	124.16	118.30
1	X	143	ASP	CB-CG-OD2	6.41	124.07	118.30
1	X	293	ASP	CB-CG-OD2	6.23	123.91	118.30
1	X	307	ASP	CB-CG-OD2	5.83	123.55	118.30
1	X	354	ASP	CB-CG-OD2	5.57	123.31	118.30
1	X	319	ASP	CB-CG-OD1	-5.27	113.56	118.30
1	X	228	ASP	CB-CG-OD2	5.26	123.04	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	X	2950	0	2874	32	0
2	X	13	0	8	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	X	198	0	0	7	0
All	All	3161	0	2882	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:180:GLN:O	1:X:184:ARG:HG2	1.90	0.71
1:X:109:LEU:H	1:X:169:GLN:HE22	1.43	0.66
1:X:394:ARG:NH1	3:X:873:HOH:O	2.29	0.65
1:X:86:HIS:HE1	3:X:709:HOH:O	1.83	0.62
1:X:109:LEU:H	1:X:169:GLN:NE2	1.99	0.61
1:X:84:LYS:HB2	1:X:84:LYS:NZ	2.16	0.60
1:X:158:ASP:N	3:X:886:HOH:O	2.35	0.60
1:X:103:ASP:OD1	1:X:138:THR:HG23	2.03	0.58
1:X:251:GLN:HE21	1:X:267:GLN:HE22	1.56	0.54
1:X:353:SER:OG	1:X:356:GLU:HG2	2.08	0.54
1:X:328:SER:HB2	1:X:333:ILE:HD11	1.90	0.52
1:X:84:LYS:CB	1:X:84:LYS:NZ	2.73	0.51
1:X:66:ASN:HD22	1:X:67:THR:N	2.08	0.51
1:X:328:SER:O	1:X:329:ALA:HB3	2.11	0.51
1:X:141:ILE:HG23	1:X:243:ALA:HB1	1.94	0.49
1:X:115:LEU:HD22	1:X:317:LEU:HD22	1.94	0.49
1:X:317:LEU:HB3	1:X:325:PHE:HB2	1.96	0.48
1:X:122:TRP:CZ2	1:X:366:PHE:HA	2.49	0.47
1:X:331:ASN:N	3:X:870:HOH:O	2.48	0.46
1:X:106:ALA:HB1	1:X:107:PRO:HD2	1.98	0.45
1:X:86:HIS:HD2	3:X:756:HOH:O	1.99	0.45
1:X:101:GLY:HA3	2:X:701:TYR:CE1	2.53	0.44
1:X:251:GLN:HG3	2:X:701:TYR:CZ	2.53	0.44
1:X:274:TYR:O	1:X:278:ILE:HG12	2.18	0.43
1:X:266:MET:HA	1:X:310:VAL:O	2.19	0.43
1:X:184:ARG:HB2	1:X:186:TYR:CE1	2.54	0.42
1:X:101:GLY:HA3	2:X:701:TYR:CD1	2.54	0.41
1:X:330:GLY:HA2	3:X:870:HOH:O	2.20	0.41
1:X:135:GLY:CA	1:X:138:THR:HG22	2.51	0.41
1:X:84:LYS:HZ2	1:X:84:LYS:HB2	1.86	0.41
1:X:145:THR:HA	1:X:229:THR:HG21	2.01	0.40
1:X:394:ARG:NE	3:X:895:HOH:O	2.42	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	X	366/392 (93%)	358 (98%)	4 (1%)	4 (1%)	14 5

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	X	238	ASP
1	X	232	ASN
1	X	235	THR
1	X	329	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	X	305/333 (92%)	294 (96%)	11 (4%)	35 23

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	X	66	ASN
1	X	84	LYS
1	X	86	HIS
1	X	184	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	X	209	LEU
1	X	219	LEU
1	X	220	ARG
1	X	234	MET
1	X	240	VAL
1	X	317	LEU
1	X	355	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	X	51	ASN
1	X	66	ASN
1	X	86	HIS
1	X	169	GLN
1	X	176	ASN
1	X	206	GLN
1	X	236	GLN
1	X	267	GLN
1	X	273	GLN
1	X	276	ASN
1	X	339	GLN
1	X	355	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	TYR	X	701	-	10,13,13	1.20	3 (30%)	12,17,17	0.73	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TYR	X	701	-	-	0/4/8/8	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	X	701	TYR	CE1-CD1	2.29	1.42	1.38
2	X	701	TYR	CE1-CZ	2.08	1.42	1.38
2	X	701	TYR	CE2-CD2	2.05	1.42	1.38

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	X	701	TYR	3	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	X	370/392 (94%)	0.19	32 (8%) 10 16	14, 23, 48, 79	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	X	236	GLN	7.2
1	X	235	THR	5.9
1	X	329	ALA	5.7
1	X	226	SER	5.4
1	X	232	ASN	5.3
1	X	158	ASP	5.1
1	X	39	PRO	4.5
1	X	237	GLY	4.4
1	X	330	GLY	4.4
1	X	233	LYS	4.2
1	X	230	VAL	4.2
1	X	146	GLY	4.1
1	X	145	THR	3.9
1	X	228	ASP	3.8
1	X	234	MET	3.7
1	X	238	ASP	3.5
1	X	41	TYR	3.4
1	X	142	GLY	3.1
1	X	229	THR	3.1
1	X	299	TYR	3.0
1	X	331	ASN	2.9
1	X	294	PRO	2.8
1	X	231	LYS	2.7
1	X	144	PRO	2.5
1	X	40	LYS	2.5
1	X	338	TYR	2.3
1	X	42	THR	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	X	293	ASP	2.1
1	X	143	ASP	2.0
1	X	393	ALA	2.0
1	X	168	TYR	2.0
1	X	217	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	TYR	X	701	13/13	0.64	0.26	31,34,39,39	0

6.5 Other polymers [i](#)

There are no such residues in this entry.