



## wwPDB EM Validation Summary Report ⓘ

Dec 4, 2022 – 12:57 AM JST

PDB ID : 7Y5B  
EMDB ID : EMD-33615  
Title : Cryo-EM structure of F-ATP synthase from Mycolicibacterium smegmatis (rotational state 1)  
Authors : Saw, W.-G.; Wong, C.F.; Grueber, G.  
Deposited on : 2022-06-16  
Resolution : 4.40 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis	:	0.0.1.dev43
Mogul	:	1.8.5 (274361), CSD as541be (2020)
MolProbity	:	4.02b-467
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ	:	1.9.9
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.31.3

# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:  
*ELECTRON MICROSCOPY*










The reported resolution of this entry is 4.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	548	 94% 6%
1	B	548	 5% 92% 8%
2	C	548	 5% 94% 6%
3	D	481	 6% 96% .
3	E	481	 96% .
3	F	481	 96% .
4	G	307	 7% 95% 5%
5	H	121	 16% 96% . .
6	a	252	 42% 82% 18%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
7	b	170	
8	d	445	
9	1	86	
9	2	86	
9	3	86	
9	4	86	
9	5	86	
9	6	86	
9	7	86	
9	8	86	
9	9	86	

## 2 Entry composition

There are 12 unique types of molecules in this entry. The entry contains 35959 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATP synthase subunit alpha.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	516	Total	C	N	O	S	0	0
			3843	2418	673	742	10		
1	B	505	Total	C	N	O	S	0	0
			3790	2386	659	735	10		

- Molecule 2 is a protein called ATP synthase subunit alpha.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	C	514	Total	C	N	O	S	0	0
			3822	2407	672	733	10		

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	1	UNK	MET	conflict	UNP A0R202
C	2	UNK	ALA	conflict	UNP A0R202
C	3	UNK	GLU	conflict	UNP A0R202
C	4	UNK	LEU	conflict	UNP A0R202
C	5	UNK	THR	conflict	UNP A0R202
C	6	UNK	ILE	conflict	UNP A0R202
C	7	UNK	SER	conflict	UNP A0R202
C	8	UNK	ALA	conflict	UNP A0R202
C	9	UNK	ALA	conflict	UNP A0R202
C	10	UNK	ASP	conflict	UNP A0R202
C	11	UNK	ILE	conflict	UNP A0R202
C	12	UNK	GLU	conflict	UNP A0R202
C	13	UNK	GLY	conflict	UNP A0R202
C	14	UNK	ALA	conflict	UNP A0R202
C	15	UNK	ILE	conflict	UNP A0R202
C	16	UNK	GLU	conflict	UNP A0R202
C	17	UNK	ASP	conflict	UNP A0R202
C	18	UNK	TYR	conflict	UNP A0R202
C	19	UNK	VAL	conflict	UNP A0R202

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	20	UNK	SER	conflict	UNP A0R202
C	21	UNK	SER	conflict	UNP A0R202
C	22	UNK	PHE	conflict	UNP A0R202

- Molecule 3 is a protein called ATP synthase subunit beta.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	D	464	Total	C	N	O	S	0	0
			3503	2218	607	665	13		
3	E	464	Total	C	N	O	S	0	0
			3508	2222	605	668	13		
3	F	464	Total	C	N	O	S	0	0
			3528	2231	608	676	13		

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	-5	MET	-	initiating methionine	UNP A0R200
D	-4	HIS	-	expression tag	UNP A0R200
D	-3	HIS	-	expression tag	UNP A0R200
D	-2	HIS	-	expression tag	UNP A0R200
D	-1	HIS	-	expression tag	UNP A0R200
D	0	HIS	-	expression tag	UNP A0R200
D	1	HIS	-	expression tag	UNP A0R200
E	-5	MET	-	initiating methionine	UNP A0R200
E	-4	HIS	-	expression tag	UNP A0R200
E	-3	HIS	-	expression tag	UNP A0R200
E	-2	HIS	-	expression tag	UNP A0R200
E	-1	HIS	-	expression tag	UNP A0R200
E	0	HIS	-	expression tag	UNP A0R200
E	1	HIS	-	expression tag	UNP A0R200
F	-5	MET	-	initiating methionine	UNP A0R200
F	-4	HIS	-	expression tag	UNP A0R200
F	-3	HIS	-	expression tag	UNP A0R200
F	-2	HIS	-	expression tag	UNP A0R200
F	-1	HIS	-	expression tag	UNP A0R200
F	0	HIS	-	expression tag	UNP A0R200
F	1	HIS	-	expression tag	UNP A0R200

- Molecule 4 is a protein called ATP synthase gamma chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	G	292	Total	C	N	O	S	0	0
			2220	1384	407	423	6		

- Molecule 5 is a protein called ATP synthase epsilon chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	H	117	Total	C	N	O	S	0	0
			862	539	159	163	1		

- Molecule 6 is a protein called ATP synthase subunit a.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	a	207	Total	C	N	O	S	0	0
			1604	1093	252	254	5		

- Molecule 7 is a protein called ATP synthase subunit b.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	b	145	Total	C	N	O	S	0	0
			1055	656	192	205	2		

- Molecule 8 is a protein called ATP synthase subunit b-delta.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	d	431	Total	C	N	O	S	0	0
			3010	1875	559	571	5		

- Molecule 9 is a protein called ATP synthase subunit c.


Mol	Chain	Residues	Atoms					AltConf	Trace
9	1	81	Total	C	N	O	S	0	0
			560	370	92	96	2		
9	2	81	Total	C	N	O	S	0	0
			552	366	90	94	2		
9	3	80	Total	C	N	O	S	0	0
			548	362	89	95	2		
9	4	81	Total	C	N	O	S	0	0
			564	372	92	98	2		
9	5	81	Total	C	N	O	S	0	0
			554	367	89	96	2		
9	6	81	Total	C	N	O	S	0	0
			556	368	90	96	2		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf	Trace
9	7	81	Total	C	N	O	S	0	0
			556	368	91	95	2		
9	8	83	Total	C	N	O	S	0	0
			573	379	94	98	2		
9	9	81	Total	C	N	O	S	0	0
			564	372	92	98	2		

- Molecule 10 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula:  $C_{10}H_{16}N_5O_{13}P_3$ ) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					AltConf
10	A	1	Total	C	N	O	P	0
			31	10	5	13	3	
10	B	1	Total	C	N	O	P	0
			31	10	5	13	3	
10	C	1	Total	C	N	O	P	0
			31	10	5	13	3	
10	E	1	Total	C	N	O	P	0
			31	10	5	13	3	
10	F	1	Total	C	N	O	P	0
			31	10	5	13	3	

- Molecule 11 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
11	A	1	Total	Mg	0
			1	1	
11	B	1	Total	Mg	0
			1	1	
11	C	1	Total	Mg	0
			1	1	
11	E	1	Total	Mg	0
			1	1	
11	F	1	Total	Mg	0
			1	1	

- Molecule 12 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula:  $C_{10}H_{15}N_5O_{10}P_2$ ) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					AltConf
12	D	1	Total	C	N	O	P	0
			27	10	5	10	2	




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ATP synthase subunit alpha




- Molecule 1: ATP synthase subunit alpha




- Molecule 2: ATP synthase subunit alpha




- Molecule 3: ATP synthase subunit beta







- Molecule 3: ATP synthase subunit beta




- Molecule 3: ATP synthase subunit beta






- Molecule 4: ATP synthase gamma chain




- Molecule 5: ATP synthase epsilon chain






- Molecule 6: ATP synthase subunit a









- Molecule 7: ATP synthase subunit b





- Molecule 8: ATP synthase subunit b-delta




- Molecule 9: ATP synthase subunit c




- Molecule 9: ATP synthase subunit c



- Molecule 9: ATP synthase subunit c






- Molecule 9: ATP synthase subunit c

Chain 4: 50% 94% 6%




- Molecule 9: ATP synthase subunit c

Chain 5: 56% 94% 6%




- Molecule 9: ATP synthase subunit c


Chain 6: 56% 94% 6%




- Molecule 9: ATP synthase subunit c

Chain 7: 45% 93% 6%







● Molecule 9: ATP synthase subunit c



● Molecule 9: ATP synthase subunit c



## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	22396	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	38.4	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	130000	Depositor
Image detector	GATAN K2 QUANTUM (4k x 4k)	Depositor
Maximum map value	0.065	Depositor
Minimum map value	-0.030	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.001	Depositor
Recommended contour level	0.015	Depositor
Map size ( $\text{\AA}$ )	484.0, 484.0, 484.0	wwPDB
Map dimensions	440, 440, 440	wwPDB
Map angles ( $^\circ$ )	90.0, 90.0, 90.0	wwPDB
Pixel spacing ( $\text{\AA}$ )	1.1, 1.1, 1.1	Depositor

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ATP, ADP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.25	0/3903	0.50	0/5289
1	B	0.25	0/3849	0.49	0/5211
2	C	0.25	0/3826	0.47	0/5182
3	D	0.25	0/3568	0.50	0/4845
3	E	0.25	0/3573	0.49	0/4850
3	F	0.25	0/3593	0.50	0/4875
4	G	0.26	0/2249	0.52	0/3040
5	H	0.25	0/873	0.58	0/1190
6	a	0.27	0/1651	0.51	0/2258
7	b	0.24	0/1064	0.49	0/1438
8	d	0.24	0/3036	0.49	0/4132
9	1	0.27	0/571	0.49	0/774
9	2	0.27	0/563	0.51	0/764
9	3	0.27	0/559	0.48	0/758
9	4	0.27	0/575	0.46	0/779
9	5	0.28	0/565	0.49	0/767
9	6	0.27	0/567	0.49	0/769
9	7	0.29	0/567	0.53	0/769
9	8	0.28	0/584	0.47	0/793
9	9	0.31	0/575	0.54	0/779
All	All	0.25	0/36311	0.50	0/49262

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

Due to software issues we are unable to calculate clashes - this section is therefore empty.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	514/548 (94%)	494 (96%)	20 (4%)	0	100	100
1	B	501/548 (91%)	484 (97%)	17 (3%)	0	100	100
2	C	499/548 (91%)	492 (99%)	7 (1%)	0	100	100
3	D	462/481 (96%)	448 (97%)	14 (3%)	0	100	100
3	E	462/481 (96%)	448 (97%)	14 (3%)	0	100	100
3	F	462/481 (96%)	448 (97%)	14 (3%)	0	100	100
4	G	288/307 (94%)	280 (97%)	8 (3%)	0	100	100
5	H	115/121 (95%)	112 (97%)	3 (3%)	0	100	100
6	a	203/252 (81%)	193 (95%)	10 (5%)	0	100	100
7	b	143/170 (84%)	138 (96%)	5 (4%)	0	100	100
8	d	423/445 (95%)	418 (99%)	5 (1%)	0	100	100
9	1	79/86 (92%)	79 (100%)	0	0	100	100
9	2	79/86 (92%)	78 (99%)	1 (1%)	0	100	100
9	3	78/86 (91%)	78 (100%)	0	0	100	100
9	4	79/86 (92%)	76 (96%)	3 (4%)	0	100	100
9	5	79/86 (92%)	77 (98%)	2 (2%)	0	100	100
9	6	79/86 (92%)	78 (99%)	1 (1%)	0	100	100
9	7	79/86 (92%)	78 (99%)	1 (1%)	0	100	100
9	8	81/86 (94%)	78 (96%)	3 (4%)	0	100	100
9	9	79/86 (92%)	77 (98%)	2 (2%)	0	100	100
All	All	4784/5156 (93%)	4654 (97%)	130 (3%)	0	100	100



There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	399/441 (90%)	399 (100%)	0	100	100
1	B	397/441 (90%)	395 (100%)	2 (0%)	88	93
2	C	392/424 (92%)	392 (100%)	0	100	100
3	D	367/394 (93%)	365 (100%)	2 (0%)	88	93
3	E	368/394 (93%)	368 (100%)	0	100	100
3	F	374/394 (95%)	374 (100%)	0	100	100
4	G	221/238 (93%)	221 (100%)	0	100	100
5	H	80/96 (83%)	79 (99%)	1 (1%)	69	82
6	a	163/200 (82%)	163 (100%)	0	100	100
7	b	105/124 (85%)	103 (98%)	2 (2%)	57	75
8	d	269/347 (78%)	266 (99%)	3 (1%)	73	85
9	1	50/56 (89%)	49 (98%)	1 (2%)	55	73
9	2	48/56 (86%)	48 (100%)	0	100	100
9	3	48/56 (86%)	48 (100%)	0	100	100
9	4	51/56 (91%)	51 (100%)	0	100	100
9	5	49/56 (88%)	49 (100%)	0	100	100
9	6	49/56 (88%)	49 (100%)	0	100	100
9	7	49/56 (88%)	48 (98%)	1 (2%)	55	73
9	8	51/56 (91%)	51 (100%)	0	100	100
9	9	51/56 (91%)	51 (100%)	0	100	100
All	All	3581/3997 (90%)	3569 (100%)	12 (0%)	92	95

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
8	d	20	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
8	d	57	MET
9	7	52	ARG
8	d	149	ARG
3	D	258	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
9	6	71	ASN
9	9	37	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 5 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
10	ATP	C	601	11	26,33,33	0.61	0	31,52,52	0.74	2 (6%)
10	ATP	F	601	11	26,33,33	0.61	0	31,52,52	0.75	2 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
10	ATP	B	601	11	26,33,33	0.61	0	31,52,52	0.73	2 (6%)
10	ATP	E	601	11	26,33,33	0.61	0	31,52,52	0.74	2 (6%)
10	ATP	A	601	11	26,33,33	0.61	0	31,52,52	0.73	2 (6%)
12	ADP	D	601	-	24,29,29	0.95	1 (4%)	29,45,45	1.42	4 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	ATP	C	601	11	-	4/18/38/38	0/3/3/3
10	ATP	F	601	11	-	5/18/38/38	0/3/3/3
10	ATP	B	601	11	-	5/18/38/38	0/3/3/3
10	ATP	E	601	11	-	2/18/38/38	0/3/3/3
10	ATP	A	601	11	-	3/18/38/38	0/3/3/3
12	ADP	D	601	-	-	2/12/32/32	0/3/3/3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
12	D	601	ADP	C5-C4	2.47	1.47	1.40

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
12	D	601	ADP	C3'-C2'-C1'	3.47	106.19	100.98
12	D	601	ADP	PA-O3A-PB	-3.30	121.50	132.83
12	D	601	ADP	N3-C2-N1	-3.11	123.81	128.68
12	D	601	ADP	C4-C5-N7	-2.54	106.76	109.40
10	E	601	ATP	C5-C6-N6	2.31	123.87	120.35

There are no chirality outliers.

5 of 21 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
10	A	601	ATP	O4'-C4'-C5'-O5'
10	B	601	ATP	C5'-O5'-PA-O3A
10	B	601	ATP	C3'-C4'-C5'-O5'

*Continued on next page...*


*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
10	C	601	ATP	PB-O3B-PG-O2G
10	F	601	ATP	C5'-O5'-PA-O2A


There are no ring outliers.


No monomer is involved in short contacts.



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.













## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-33615. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X




Y




Z


#### 6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

## 6.2 Central slices [i](#)


### 6.2.1 Primary map



X Index: 220




Y Index: 220




Z Index: 220


### 6.2.2 Raw map



X Index: 220



Y Index: 220




Z Index: 220


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 255




Y Index: 220




Z Index: 260


### 6.3.2 Raw map



X Index: 211



Y Index: 219



Z Index: 249

The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal surface views [i](#)

### 6.4.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.015. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

### 6.4.2 Raw map



X



Y



Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


## 6.5 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


### 6.5.1 emd\_33615\_msk\_1.map [i](#)



X



Y




Z

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

## 7.2 Volume estimate [i](#)




The volume at the recommended contour level is 254 nm<sup>3</sup>; this corresponds to an approximate mass of 229 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



### 7.3 Rotationally averaged power spectrum ⓘ




\*Reported resolution corresponds to spatial frequency of 0.227 Å<sup>-1</sup>

## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of 0.227 Å<sup>-1</sup>

## 8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	4.40	-	-
Author-provided FSC curve	4.40	6.82	4.47
Unmasked-calculated*	7.56	9.78	7.96

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 7.56 differs from the reported value 4.4 by more than 10 %

## 9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-33615 and PDB model 7Y5B. Per-residue inclusion information can be found in section [3](#) on page [9](#).

### 9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.015 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.015).











































## 9.4 Atom inclusion [i](#)



At the recommended contour level, 81% of all backbone atoms, 69% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.015) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.6916	 0.3950
1	 0.4739	 0.2920
2	 0.3583	 0.2330
3	 0.3554	 0.2560
4	 0.3918	 0.2680
5	 0.3739	 0.2610
6	 0.3539	 0.2520
7	 0.4192	 0.2960
8	 0.5053	 0.2880
9	 0.4866	 0.2950
A	 0.8069	 0.4490
B	 0.7606	 0.4340
C	 0.7877	 0.4400
D	 0.7582	 0.4270
E	 0.7974	 0.4600
F	 0.8037	 0.4570
G	 0.7224	 0.4130
H	 0.6468	 0.3900
a	 0.4058	 0.2510
b	 0.5130	 0.2890
d	 0.6710	 0.3410

