



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 25, 2020 – 02:17 PM BST

PDB ID : 2Y6T
Title : Molecular Recognition of Chymotrypsin by the Serine Protease Inhibitor Ecotin from *Yersinia pestis*
Authors : Clark, E.A.; Walker, N.; Ford, D.C.; Cooper, I.A.; Oyston, P.C.F.; Acharya, K.R.
Deposited on : 2011-01-26
Resolution : 2.74 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.13
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.13


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.74 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1271 (2.76-2.72)
Clashscore	141614	1322 (2.76-2.72)
Ramachandran outliers	138981	1297 (2.76-2.72)
Sidechain outliers	138945	1298 (2.76-2.72)
RSRZ outliers	127900	1243 (2.76-2.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	245	<div> <div>%</div> <div> <div></div> <div>90%</div> <div>7%</div> <div>••</div> </div> </div>
1	B	245	<div> <div>2%</div> <div> <div></div> <div>91%</div> <div>7%</div> <div>•</div> </div> </div>
1	C	245	<div> <div>%</div> <div> <div></div> <div>91%</div> <div>7%</div> <div>•</div> </div> </div>
1	D	245	<div> <div></div> <div> <div></div> <div>89%</div> <div>9%</div> <div>•</div> </div> </div>
2	E	148	<div> <div>3%</div> <div> <div></div> <div>83%</div> <div>11%</div> <div>• 5%</div> </div> </div>
2	F	148	<div> <div>4%</div> <div> <div></div> <div>84%</div> <div>11%</div> <div>••</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	G	148	<div><div></div><div>3%</div><div>82%</div><div>11%</div><div>• 5%</div></div>
2	H	148	<div><div></div><div>5%</div><div>78%</div><div>11%</div><div>• 7%</div></div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 11529 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CHYMOTRYPSINOGEN A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	240	Total	C	N	O	S	0	0	1
			1756	1102	298	344	12			
1	B	240	Total	C	N	O	S	0	0	1
			1756	1102	298	344	12			
1	C	240	Total	C	N	O	S	0	0	1
			1756	1102	298	344	12			
1	D	240	Total	C	N	O	S	0	0	1
			1756	1102	298	344	12			

- Molecule 2 is a protein called ECOTIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	141	Total	C	N	O	S	0	0	0
			1120	708	195	209	8			
2	F	142	Total	C	N	O	S	0	0	0
			1128	712	197	211	8			
2	G	140	Total	C	N	O	S	0	0	0
			1111	703	193	207	8			
2	H	138	Total	C	N	O	S	0	0	0
			1093	693	189	203	8			

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	4	Total	O	0	0
			4	4		
4	B	6	Total	O	0	0
			6	6		
4	C	5	Total	O	0	0
			5	5		
4	D	4	Total	O	0	0
			4	4		
4	E	3	Total	O	0	0
			3	3		
4	F	4	Total	O	0	0
			4	4		
4	G	6	Total	O	0	0
			6	6		
4	H	1	Total	O	0	0
			1	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: CHYMOTRYPSINOGEN A




- Molecule 1: CHYMOTRYPSINOGEN A




- Molecule 1: CHYMOTRYPSINOGEN A




- Molecule 1: CHYMOTRYPSINOGEN A




- Molecule 2: ECOTIN




- Molecule 2: ECOTIN



• Molecule 2: ECOTIN



• Molecule 2: ECOTIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	97.31Å 48.28Å 174.64Å 90.00° 103.96° 90.00°	Depositor
Resolution (Å)	169.48 – 2.74 41.49 – 2.74	Depositor EDS
% Data completeness (in resolution range)	97.3 (169.48-2.74) 97.4 (41.49-2.74)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.24 (at 2.73Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.244 , 0.314 0.240 , 0.308	Depositor DCC
R_{free} test set	2080 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	39.9	Xtriage
Anisotropy	0.560	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 28.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.018 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	11529	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.35	0/1790	0.50	0/2439
1	B	0.35	0/1790	0.49	0/2439
1	C	0.35	0/1790	0.48	0/2439
1	D	0.35	0/1790	0.48	0/2439
2	E	0.34	0/1142	0.50	0/1542
2	F	0.34	0/1150	0.50	0/1553
2	G	0.36	0/1133	0.53	1/1530 (0.1%)
2	H	0.34	0/1115	0.50	0/1505
All	All	0.35	0/11700	0.49	1/15886 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	130	ASP	CB-CG-OD2	5.21	122.99	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1756	0	1730	12	0
1	B	1756	0	1730	8	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1756	0	1730	8	0
1	D	1756	0	1730	7	0
2	E	1120	0	1125	12	0
2	F	1128	0	1131	14	0
2	G	1111	0	1117	11	0
2	H	1093	0	1102	14	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
3	C	5	0	0	0	0
3	D	5	0	0	0	0
4	A	4	0	0	0	0
4	B	6	0	0	0	0
4	C	5	0	0	0	0
4	D	4	0	0	0	0
4	E	3	0	0	0	0
4	F	4	0	0	0	0
4	G	6	0	0	0	0
4	H	1	0	0	0	0
All	All	11529	0	11395	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:77:ASN:HD21	2:E:111:MET:H	1.30	0.79
2:F:146:VAL:HG13	2:F:150:VAL:HB	1.66	0.77
2:H:77:ASN:HD21	2:H:111:MET:H	1.31	0.76
1:D:81:GLN:HE22	1:D:113:SER:H	1.41	0.67
1:A:32:SER:HB2	1:A:40:HIS:ND1	2.10	0.67
2:F:77:ASN:HD21	2:F:111:MET:H	1.43	0.65
2:G:77:ASN:HB2	2:G:109:THR:OG1	1.97	0.65
2:E:64:VAL:HG13	2:E:134:GLN:HB2	1.81	0.62
2:F:31:GLN:HB2	2:F:32:PRO:CD	2.29	0.62
1:C:41:PHE:HA	2:G:112:ALA:HB3	1.82	0.61
1:A:21:GLU:OE2	1:A:154:ARG:HG3	2.01	0.60
2:F:31:GLN:HB2	2:F:32:PRO:HD2	1.83	0.59
1:A:239:GLN:HA	1:A:239:GLN:HE21	1.67	0.58
2:H:77:ASN:HD21	2:H:111:MET:N	2.01	0.57
1:A:191:CYS:HA	2:E:110:MET:HE2	1.87	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:98:THR:HA	2:G:80:MET:HE3	1.88	0.56
1:B:41:PHE:HA	2:F:112:ALA:HB3	1.88	0.55
2:E:155:ARG:HD3	2:F:159:ALA:HB1	1.89	0.55
2:G:74:VAL:HG23	2:G:120:GLN:HG3	1.88	0.54
2:G:118:LYS:N	2:G:119:PRO:HD3	2.22	0.54
2:E:146:VAL:HG13	2:E:150:VAL:HB	1.90	0.54
1:B:219:THR:HG22	1:B:219:THR:O	2.09	0.52
1:C:99:ILE:HD11	2:G:78:ARG:HG2	1.91	0.52
2:H:105:GLN:HG2	2:H:106:PRO:HD2	1.92	0.52
2:G:123:PHE:HE1	2:G:151:GLU:HG3	1.76	0.51
1:B:201:CYS:SG	1:B:210:VAL:HG21	2.51	0.51
2:H:77:ASN:ND2	2:H:111:MET:H	2.06	0.51
1:B:57:HIS:HD2	2:F:111:MET:HE2	1.77	0.49
1:C:193:GLY:HA2	2:G:112:ALA:HB2	1.93	0.49
2:G:61:ARG:HB2	2:G:159:ALA:HB3	1.94	0.49
1:C:105:LEU:HD11	1:C:238:VAL:HG13	1.95	0.48
2:F:77:ASN:HD21	2:F:111:MET:CB	2.26	0.48
2:H:118:LYS:H	2:H:118:LYS:HD3	1.80	0.47
1:B:240:GLN:HG3	1:B:241:THR:N	2.29	0.46
1:A:191:CYS:HA	2:E:110:MET:CE	2.45	0.46
2:H:100:MET:O	2:H:147:PRO:HD3	2.15	0.46
1:D:83:LEU:HD22	1:D:108:LEU:HD13	1.97	0.46
2:E:77:ASN:ND2	2:E:111:MET:H	2.05	0.46
2:F:146:VAL:CG1	2:F:150:VAL:HB	2.41	0.46
1:A:143:LEU:HD22	1:A:150:ASN:HB2	1.98	0.46
2:H:114:PRO:O	2:H:117:SER:HB3	2.16	0.46
1:D:164:SER:OG	1:D:167:ASN:HB2	2.15	0.45
1:A:238:VAL:O	1:A:242:LEU:HG	2.15	0.45
1:A:27:TRP:CZ2	1:A:137:VAL:HG11	2.52	0.45
2:E:48:ARG:HD2	2:F:169:LYS:O	2.17	0.45
2:G:146:VAL:HG13	2:G:150:VAL:HB	1.98	0.45
1:B:97:LEU:HD23	2:F:78:ARG:HH22	1.81	0.44
1:A:154:ARG:H	1:A:154:ARG:HD3	1.82	0.44
2:H:74:VAL:HG21	2:H:122:LYS:HG2	2.00	0.44
2:F:77:ASN:HD21	2:F:111:MET:HB2	1.83	0.43
1:D:160:LEU:HB2	1:D:184:GLY:HA2	2.00	0.43
2:H:81:LEU:HD22	2:H:124:VAL:HG12	2.00	0.43
1:A:51:TRP:CD2	1:A:242:LEU:HD22	2.54	0.43
2:F:69:GLY:HA3	2:F:124:VAL:O	2.18	0.43
2:H:156:ILE:H	2:H:156:ILE:HG13	1.70	0.43
2:G:85:LEU:HD21	2:G:98:LEU:HD23	2.01	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:65:VAL:HG11	1:C:82:LYS:HG3	2.00	0.43
2:H:146:VAL:HG13	2:H:150:VAL:HB	2.00	0.43
2:H:161:GLU:CD	2:H:161:GLU:H	2.22	0.43
1:C:212:ILE:HB	1:C:229:ALA:HB3	2.00	0.42
1:B:138:THR:HG22	1:B:160:LEU:HD22	2.01	0.42
2:E:74:VAL:HB	2:E:79:HIS:CE1	2.54	0.42
1:D:219:THR:HG22	1:D:219:THR:O	2.20	0.42
1:A:71:PHE:O	1:A:154:ARG:HA	2.19	0.42
1:D:212:ILE:HB	1:D:229:ALA:HB3	2.02	0.42
2:E:105:GLN:HG2	2:E:106:PRO:HD2	2.02	0.41
2:H:64:VAL:HG13	2:H:134:GLN:HB2	2.02	0.41
1:B:169:LYS:C	1:B:171:TYR:H	2.23	0.41
1:C:211:GLY:HA2	1:C:229:ALA:O	2.20	0.41
2:E:81:LEU:HA	2:E:81:LEU:HD12	1.97	0.41
2:F:128:LEU:HB2	2:F:132:ALA:HB2	2.02	0.41
1:A:172:TRP:CZ3	2:E:107:ALA:HB2	2.55	0.41
1:D:105:LEU:HD11	1:D:238:VAL:HA	2.02	0.41
2:H:148:GLN:HB3	2:H:148:GLN:HE21	1.67	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	234/245 (96%)	224 (96%)	10 (4%)	0	100	100
1	B	234/245 (96%)	216 (92%)	17 (7%)	1 (0%)	34	55
1	C	234/245 (96%)	223 (95%)	11 (5%)	0	100	100
1	D	234/245 (96%)	222 (95%)	11 (5%)	1 (0%)	34	55
2	E	139/148 (94%)	124 (89%)	14 (10%)	1 (1%)	22	40
2	F	140/148 (95%)	134 (96%)	5 (4%)	1 (1%)	22	40

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	G	138/148 (93%)	130 (94%)	8 (6%)	0	100	100
2	H	136/148 (92%)	131 (96%)	5 (4%)	0	100	100
All	All	1489/1572 (95%)	1404 (94%)	81 (5%)	4 (0%)	41	61

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	E	43	GLU
1	D	63	SER
2	F	31	GLN
1	B	99	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	195/200 (98%)	189 (97%)	6 (3%)	40	60
1	B	195/200 (98%)	191 (98%)	4 (2%)	53	72
1	C	195/200 (98%)	188 (96%)	7 (4%)	35	55
1	D	195/200 (98%)	186 (95%)	9 (5%)	27	46
2	E	123/130 (95%)	120 (98%)	3 (2%)	49	68
2	F	124/130 (95%)	120 (97%)	4 (3%)	39	59
2	G	122/130 (94%)	120 (98%)	2 (2%)	62	78
2	H	120/130 (92%)	113 (94%)	7 (6%)	20	35
All	All	1269/1320 (96%)	1227 (97%)	42 (3%)	38	59

All (42) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	96	SER
1	A	154	ARG
1	A	165	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	232	THR
1	A	239	GLN
1	A	240	GLN
1	B	165	ASN
1	B	192	MET
1	B	195	SER
1	B	232	THR
1	C	92	SER
1	C	165	ASN
1	C	192	MET
1	C	195	SER
1	C	219	THR
1	C	232	THR
1	C	234	LEU
1	D	80	ILE
1	D	92	SER
1	D	151	THR
1	D	165	ASN
1	D	167	ASN
1	D	191	CYS
1	D	195	SER
1	D	232	THR
1	D	234	LEU
2	E	116	ASP
2	E	146	VAL
2	E	161	GLU
2	F	64	VAL
2	F	135	ARG
2	F	140	LEU
2	F	162	ASP
2	G	118	LYS
2	G	146	VAL
2	H	61	ARG
2	H	81	LEU
2	H	118	LYS
2	H	128	LEU
2	H	146	VAL
2	H	148	GLN
2	H	161	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (26) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	165	ASN
1	A	167	ASN
1	A	239	GLN
1	B	18	ASN
1	B	165	ASN
1	C	165	ASN
1	C	167	ASN
1	D	81	GLN
1	D	101	ASN
1	D	204	ASN
1	D	240	GLN
2	E	29	GLN
2	E	56	GLN
2	E	77	ASN
2	E	105	GLN
2	F	31	GLN
2	F	56	GLN
2	F	77	ASN
2	F	79	HIS
2	G	79	HIS
2	G	105	GLN
2	G	134	GLN
2	H	77	ASN
2	H	105	GLN
2	H	127	ASN
2	H	148	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	D	1246	-	4,4,4	0.16	0	6,6,6	0.11	0
3	SO4	B	1246	-	4,4,4	0.12	0	6,6,6	0.09	0
3	SO4	C	1246	-	4,4,4	0.14	0	6,6,6	0.11	0
3	SO4	A	1246	-	4,4,4	0.14	0	6,6,6	0.10	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	240/245 (97%)	-0.02	3 (1%) 77 82	17, 29, 44, 50	0
1	B	240/245 (97%)	-0.02	4 (1%) 70 76	24, 34, 44, 51	0
1	C	240/245 (97%)	0.17	2 (0%) 86 89	22, 34, 47, 51	0
1	D	240/245 (97%)	0.05	1 (0%) 92 95	20, 32, 47, 53	0
2	E	141/148 (95%)	0.26	4 (2%) 53 60	26, 38, 49, 57	0
2	F	142/148 (95%)	0.36	6 (4%) 36 39	30, 43, 52, 54	0
2	G	140/148 (94%)	0.14	4 (2%) 51 58	19, 36, 50, 52	0
2	H	138/148 (93%)	0.41	7 (5%) 28 31	34, 45, 51, 52	0
All	All	1521/1572 (96%)	0.14	31 (2%) 65 72	17, 36, 49, 57	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	150	ASN	3.8
2	F	45	GLY	3.3
1	A	147	THR	3.1
2	E	30	GLN	3.0
1	C	149	ALA	3.0
1	C	39	PHE	2.8
2	H	102	LYS	2.8
2	G	117	SER	2.7
1	B	203	LYS	2.7
2	E	31	GLN	2.6
1	B	76	SER	2.6
2	H	169	LYS	2.5
2	F	44	LYS	2.5
2	F	101	ASP	2.5
2	G	115	GLU	2.4
2	H	57	LYS	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	E	162	ASP	2.4
2	F	29	GLN	2.3
1	A	201	CYS	2.3
2	F	31	GLN	2.3
2	H	62	PHE	2.3
2	G	116	ASP	2.3
2	F	103	ILE	2.3
1	A	167	ASN	2.2
1	B	175	LYS	2.2
2	H	101	ASP	2.2
1	D	174	THR	2.1
2	H	155	ARG	2.1
2	H	145	TYR	2.1
2	E	161	GLU	2.0
2	G	118	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	SO4	C	1246	5/5	0.80	0.25	78,78,78,78	0
3	SO4	B	1246	5/5	0.91	0.16	70,71,71,71	0
3	SO4	A	1246	5/5	0.93	0.18	59,59,59,59	0
3	SO4	D	1246	5/5	0.98	0.09	38,38,38,38	0

6.5 Other polymers [i](#)

There are no such residues in this entry.