



wwPDB X-ray Structure Validation Summary Report ⓘ

May 29, 2020 – 06:38 am BST

PDB ID : 4Y8M
Title : Yeast 20S proteasome beta7-delta7_Cter mutant
Authors : Huber, E.M.; Groll, M.
Deposited on : 2015-02-16
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	250	<div> <div>2%</div> <div>98%</div> <div>•</div> </div>
1	O	250	<div> <div>2%</div> <div>97%</div> <div>•</div> </div>
2	B	258	<div> <div>2%</div> <div>86%</div> <div>8% • 5%</div> </div>
2	P	258	<div> <div>3%</div> <div>85%</div> <div>8% • 5%</div> </div>
3	C	254	<div> <div>3%</div> <div>85%</div> <div>9% • 6%</div> </div>
3	Q	254	<div> <div>6%</div> <div>85%</div> <div>8% • 6%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	D	260	
4	R	260	
5	E	234	
5	S	234	
6	F	288	
6	T	288	
7	G	252	
7	U	252	
8	H	232	
8	V	232	
9	I	205	
9	W	205	
10	J	198	
10	X	198	
11	K	212	
11	Y	212	
12	L	222	
12	Z	222	
13	M	239	
13	a	239	
14	N	196	
14	b	196	

2 Entry composition

There are 17 unique types of molecules in this entry. The entry contains 49570 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proteasome subunit alpha type-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	250	Total	C	N	O	S	0	0	0
			1915	1219	315	377	4			
1	O	250	Total	C	N	O	S	0	0	0
			1915	1219	315	377	4			

- Molecule 2 is a protein called Proteasome subunit alpha type-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	244	Total	C	N	O	S	0	0	0
			1904	1201	321	379	3			
2	P	244	Total	C	N	O	S	0	0	0
			1904	1201	321	379	3			

- Molecule 3 is a protein called Proteasome subunit alpha type-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	240	Total	C	N	O	S	0	0	0
			1881	1176	329	372	4			
3	Q	240	Total	C	N	O	S	0	0	0
			1881	1176	329	372	4			

- Molecule 4 is a protein called Proteasome subunit alpha type-5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	235	Total	C	N	O	S	0	0	0
			1813	1136	304	366	7			
4	R	235	Total	C	N	O	S	0	0	0
			1813	1136	304	366	7			

- Molecule 5 is a protein called Proteasome subunit alpha type-6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	231	Total	C	N	O	S	0	0	0
			1773	1114	307	348	4			
5	S	231	Total	C	N	O	S	0	0	0
			1773	1114	307	348	4			

- Molecule 6 is a protein called Probable proteasome subunit alpha type-7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	F	243	Total	C	N	O	S	0	0	0
			1892	1203	329	356	4			
6	T	243	Total	C	N	O	S	0	0	0
			1892	1203	329	356	4			

- Molecule 7 is a protein called Proteasome subunit alpha type-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	G	241	Total	C	N	O	S	0	0	0
			1907	1214	320	365	8			
7	U	241	Total	C	N	O	S	0	0	0
			1907	1214	320	365	8			

- Molecule 8 is a protein called Proteasome subunit beta type-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	H	226	Total	C	N	O	S	0	0	0
			1719	1082	298	332	7			
8	V	226	Total	C	N	O	S	0	0	0
			1719	1082	298	332	7			

- Molecule 9 is a protein called Proteasome subunit beta type-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	I	204	Total	C	N	O	S	0	0	0
			1581	1010	258	305	8			
9	W	204	Total	C	N	O	S	0	0	0
			1581	1010	258	305	8			

- Molecule 10 is a protein called Proteasome subunit beta type-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	J	195	Total	C	N	O	S	0	0	0
			1561	992	264	299	6			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	X	195	Total	C	N	O	S	0	0	0
			1561	992	264	299	6			

- Molecule 11 is a protein called Proteasome subunit beta type-5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
11	K	212	Total	C	N	O	S	0	0	0
			1644	1045	280	312	7			
11	Y	212	Total	C	N	O	S	0	0	0
			1644	1045	280	312	7			

- Molecule 12 is a protein called Proteasome subunit beta type-6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
12	L	222	Total	C	N	O	S	0	0	0
			1757	1115	303	335	4			
12	Z	222	Total	C	N	O	S	0	0	0
			1757	1115	303	335	4			

- Molecule 13 is a protein called Proteasome subunit beta type-7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
13	M	222	Total	C	N	O	S	0	0	0
			1736	1098	297	334	7			
13	a	222	Total	C	N	O	S	0	0	0
			1736	1098	297	334	7			

- Molecule 14 is a protein called Proteasome subunit beta type-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
14	N	196	Total	C	N	O	S	0	0	0
			1512	955	250	300	7			
14	b	196	Total	C	N	O	S	0	0	0
			1512	955	250	300	7			

- Molecule 15 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
15	G	1	Total	Mg	0	0
			1	1		
15	K	1	Total	Mg	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
15	H	1	Total 1	Mg 1	0	0
15	I	1	Total 1	Mg 1	0	0
15	V	1	Total 1	Mg 1	0	0
15	W	1	Total 1	Mg 1	0	0
15	Z	1	Total 1	Mg 1	0	0
15	N	1	Total 1	Mg 1	0	0
15	Y	1	Total 1	Mg 1	0	0

- Molecule 16 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
16	G	1	Total 1	Cl 1	0	0
16	U	1	Total 1	Cl 1	0	0

- Molecule 17 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
17	A	13	Total 13	O 13	0	0
17	B	12	Total 12	O 12	0	0
17	C	9	Total 9	O 9	0	0
17	D	9	Total 9	O 9	0	0
17	E	11	Total 11	O 11	0	0
17	F	11	Total 11	O 11	0	0
17	G	15	Total 15	O 15	0	0
17	H	11	Total 11	O 11	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
17	I	11	Total 11	O 11	0	0
17	J	15	Total 15	O 15	0	0
17	K	19	Total 19	O 19	0	0
17	L	16	Total 16	O 16	0	0
17	M	21	Total 21	O 21	0	0
17	N	14	Total 14	O 14	0	0
17	O	12	Total 12	O 12	0	0
17	P	9	Total 9	O 9	0	0
17	Q	7	Total 7	O 7	0	0
17	R	5	Total 5	O 5	0	0
17	S	5	Total 5	O 5	0	0
17	T	10	Total 10	O 10	0	0
17	U	20	Total 20	O 20	0	0
17	V	9	Total 9	O 9	0	0
17	W	14	Total 14	O 14	0	0
17	X	12	Total 12	O 12	0	0
17	Y	21	Total 21	O 21	0	0
17	Z	20	Total 20	O 20	0	0
17	a	19	Total 19	O 19	0	0
17	b	19	Total 19	O 19	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Proteasome subunit alpha type-2




- Molecule 1: Proteasome subunit alpha type-2




- Molecule 2: Proteasome subunit alpha type-3



- Molecule 2: Proteasome subunit alpha type-3




- Molecule 3: Proteasome subunit alpha type-4



LYS
LYS
SER
ASN
ASN
HIS


• Molecule 3: Proteasome subunit alpha type-4

Chain Q: 

MET SER G1 R4 Q17 K35 N38 S48 T49 L50 K51 L52 S60 V124 Q147 T148 E149 P150 S158 A159 Q160 V169 K175 K180 E181 P182 P183 V201 Q202 T203 Q204 A205 K206 Q236 E237 K238 Q239 E240 GLN GLN GLN ASP LYS LYS

LYS
SER
ASN
HIS


• Molecule 4: Proteasome subunit alpha type-5

Chain D: 

MET PHE LEU THR ARG SER GLU TTR D1 Q15 L40 L51 A88 A91 H91 I99 E117 GLY ALA SER GLY GLU GLU ARG L125 D143 N160 L176 W179 L190 L193 E202 I214 L235 K236 E242 SER PRO GLU GLU ALA ASP VAL MET

MET
SER

• Molecule 4: Proteasome subunit alpha type-5

Chain R: 

MET PHE LEU THR ARG SER GLU TTR D1 L40 L51 A88 H91 I99 E117 GLY ALA SER GLY GLU GLU ARG L125 D143 N160 L176 W179 L190 L193 E202 I214 L235 K236 E242 SER PRO GLU GLU ALA ASP VAL MET


SER

• Molecule 5: Proteasome subunit alpha type-6

Chain E: 

MET PHE ARG R3 T9 F12 K29 L55 H68 L71 L87 N99 L102 A107 Q116 Y122 T174 L175 F178 N184 L188 D202 V207 D208 K231 Y232 I233






• Molecule 5: Proteasome subunit alpha type-6

Chain S: 

MET PHE ARG R3 T9 F12 K29 L55 H68 L71 L87 N99 L102 A107 Q116 R173 T174 L175 F178 I179 K180 N184 L188 D202 V207 D208 K231 Y232 I233

• Molecule 6: Probable proteasome subunit alpha type-7

Chain F: 






- Molecule 9: Proteasome subunit beta type-3




- Molecule 10: Proteasome subunit beta type-4



- Molecule 10: Proteasome subunit beta type-4




- Molecule 11: Proteasome subunit beta type-5



- Molecule 11: Proteasome subunit beta type-5



- Molecule 12: Proteasome subunit beta type-6





- Molecule 12: Proteasome subunit beta type-6

Chain Z:  96% .




- Molecule 13: Proteasome subunit beta type-7

Chain M:  87% 5% 7%




- Molecule 13: Proteasome subunit beta type-7

Chain a:  90% 7%




- Molecule 14: Proteasome subunit beta type-1

Chain N:  97% . .



- Molecule 14: Proteasome subunit beta type-1

Chain b:  98% .



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	135.40Å 302.25Å 144.59Å 90.00° 112.75° 90.00°	Depositor
Resolution (Å)	15.00 – 2.80 15.00 – 2.80	Depositor EDS
% Data completeness (in resolution range)	95.8 (15.00-2.80) 95.8 (15.00-2.80)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.53 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.197 , 0.218 0.200 , 0.220	Depositor DCC
R_{free} test set	12479 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	59.1	Xtriage
Anisotropy	0.042	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 42.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	49570	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.27	0/1952	0.47	0/2642
1	O	0.27	0/1952	0.47	0/2642
2	B	0.28	0/1934	0.50	0/2618
2	P	0.28	0/1934	0.50	0/2618
3	C	0.28	0/1910	0.51	0/2586
3	Q	0.28	0/1910	0.51	0/2586
4	D	0.27	0/1837	0.48	0/2475
4	R	0.27	0/1837	0.48	0/2475
5	E	0.27	0/1800	0.48	0/2433
5	S	0.27	0/1800	0.48	0/2433
6	F	0.28	0/1932	0.46	0/2609
6	T	0.27	0/1932	0.46	0/2609
7	G	0.28	0/1945	0.47	0/2634
7	U	0.27	0/1945	0.47	0/2634
8	H	0.26	0/1750	0.48	0/2373
8	V	0.30	0/1750	0.48	0/2373
9	I	0.28	0/1611	0.48	0/2174
9	W	0.28	0/1611	0.48	0/2174
10	J	0.27	0/1589	0.49	0/2142
10	X	0.27	0/1589	0.49	0/2142
11	K	0.27	0/1681	0.49	1/2274 (0.0%)
11	Y	0.27	0/1681	0.49	1/2274 (0.0%)
12	L	0.28	0/1795	0.48	0/2420
12	Z	0.28	0/1795	0.48	0/2420
13	M	0.31	0/1766	0.52	0/2398
13	a	0.31	0/1766	0.52	0/2398
14	N	0.29	0/1541	0.47	0/2087
14	b	0.27	0/1541	0.48	0/2087
All	All	0.28	0/50086	0.49	2/67730 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
11	K	4	LEU	CA-CB-CG	5.22	127.30	115.30
11	Y	4	LEU	CA-CB-CG	5.04	126.90	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1915	0	1929	2	0
1	O	1915	0	1929	4	0
2	B	1904	0	1904	11	0
2	P	1904	0	1904	12	0
3	C	1881	0	1895	8	0
3	Q	1881	0	1895	7	0
4	D	1813	0	1797	6	0
4	R	1813	0	1797	4	0
5	E	1773	0	1775	7	0
5	S	1773	0	1775	4	0
6	F	1892	0	1883	4	0
6	T	1892	0	1883	3	0
7	G	1907	0	1901	5	0
7	U	1907	0	1901	6	0
8	H	1719	0	1719	7	0
8	V	1719	0	1719	6	0
9	I	1581	0	1574	6	0
9	W	1581	0	1574	7	0
10	J	1561	0	1569	10	0
10	X	1561	0	1569	10	0
11	K	1644	0	1595	6	0
11	Y	1644	0	1595	5	0
12	L	1757	0	1711	1	0
12	Z	1757	0	1711	1	0
13	M	1736	0	1737	5	0
13	a	1736	0	1737	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
14	N	1512	0	1481	1	0
14	b	1512	0	1481	0	0
15	G	1	0	0	0	0
15	H	1	0	0	0	0
15	I	1	0	0	0	0
15	K	1	0	0	0	0
15	N	1	0	0	0	0
15	V	1	0	0	0	0
15	W	1	0	0	0	0
15	Y	1	0	0	0	0
15	Z	1	0	0	0	0
16	G	1	0	0	0	0
16	U	1	0	0	0	0
17	A	13	0	0	0	0
17	B	12	0	0	0	0
17	C	9	0	0	0	0
17	D	9	0	0	0	0
17	E	11	0	0	1	0
17	F	11	0	0	0	0
17	G	15	0	0	0	0
17	H	11	0	0	0	0
17	I	11	0	0	0	0
17	J	15	0	0	0	0
17	K	19	0	0	0	0
17	L	16	0	0	0	0
17	M	21	0	0	0	0
17	N	14	0	0	0	0
17	O	12	0	0	0	0
17	P	9	0	0	2	0
17	Q	7	0	0	0	0
17	R	5	0	0	0	0
17	S	5	0	0	0	0
17	T	10	0	0	0	0
17	U	20	0	0	0	0
17	V	9	0	0	0	0
17	W	14	0	0	0	0
17	X	12	0	0	0	0
17	Y	21	0	0	0	0
17	Z	20	0	0	0	0
17	a	19	0	0	0	0
17	b	19	0	0	0	0
All	All	49570	0	48940	129	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 129 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:V:52:THR:O	8:V:56:THR:OG1	1.98	0.82
8:H:52:THR:O	8:H:56:THR:OG1	1.98	0.80
2:B:12:PHE:H	3:C:17:GLN:HE22	1.46	0.64
3:Q:51:LYS:O	3:Q:52:LEU:HB2	2.03	0.59
3:C:51:LYS:O	3:C:52:LEU:HB2	2.02	0.59

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	248/250 (99%)	238 (96%)	9 (4%)	1 (0%)	34	66
1	O	248/250 (99%)	238 (96%)	9 (4%)	1 (0%)	34	66
2	B	242/258 (94%)	236 (98%)	2 (1%)	4 (2%)	9	29
2	P	242/258 (94%)	236 (98%)	2 (1%)	4 (2%)	9	29
3	C	238/254 (94%)	232 (98%)	3 (1%)	3 (1%)	12	36
3	Q	238/254 (94%)	232 (98%)	3 (1%)	3 (1%)	12	36
4	D	231/260 (89%)	229 (99%)	2 (1%)	0	100	100
4	R	231/260 (89%)	229 (99%)	2 (1%)	0	100	100
5	E	229/234 (98%)	222 (97%)	7 (3%)	0	100	100
5	S	229/234 (98%)	222 (97%)	7 (3%)	0	100	100
6	F	241/288 (84%)	237 (98%)	4 (2%)	0	100	100
6	T	241/288 (84%)	237 (98%)	4 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
7	G	239/252 (95%)	239 (100%)	0	0	100	100
7	U	239/252 (95%)	239 (100%)	0	0	100	100
8	H	224/232 (97%)	218 (97%)	6 (3%)	0	100	100
8	V	224/232 (97%)	218 (97%)	6 (3%)	0	100	100
9	I	202/205 (98%)	197 (98%)	5 (2%)	0	100	100
9	W	202/205 (98%)	197 (98%)	5 (2%)	0	100	100
10	J	193/198 (98%)	188 (97%)	5 (3%)	0	100	100
10	X	193/198 (98%)	188 (97%)	5 (3%)	0	100	100
11	K	210/212 (99%)	206 (98%)	4 (2%)	0	100	100
11	Y	210/212 (99%)	206 (98%)	4 (2%)	0	100	100
12	L	220/222 (99%)	216 (98%)	4 (2%)	0	100	100
12	Z	220/222 (99%)	216 (98%)	4 (2%)	0	100	100
13	M	220/239 (92%)	210 (96%)	8 (4%)	2 (1%)	17	46
13	a	220/239 (92%)	211 (96%)	9 (4%)	0	100	100
14	N	194/196 (99%)	190 (98%)	4 (2%)	0	100	100
14	b	194/196 (99%)	190 (98%)	4 (2%)	0	100	100
All	All	6262/6600 (95%)	6117 (98%)	127 (2%)	18 (0%)	41	72

5 of 18 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	2	THR
2	B	51	VAL
2	B	221	ASP
3	C	202	GLN
13	M	220	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	209/209 (100%)	205 (98%)	4 (2%)	57	85
1	O	209/209 (100%)	205 (98%)	4 (2%)	57	85
2	B	203/216 (94%)	195 (96%)	8 (4%)	32	66
2	P	203/216 (94%)	195 (96%)	8 (4%)	32	66
3	C	212/226 (94%)	199 (94%)	13 (6%)	18	48
3	Q	212/226 (94%)	199 (94%)	13 (6%)	18	48
4	D	194/215 (90%)	181 (93%)	13 (7%)	16	43
4	R	194/215 (90%)	181 (93%)	13 (7%)	16	43
5	E	190/193 (98%)	178 (94%)	12 (6%)	18	46
5	S	190/193 (98%)	178 (94%)	12 (6%)	18	46
6	F	201/239 (84%)	191 (95%)	10 (5%)	24	56
6	T	201/239 (84%)	191 (95%)	10 (5%)	24	56
7	G	206/210 (98%)	195 (95%)	11 (5%)	22	54
7	U	206/210 (98%)	195 (95%)	11 (5%)	22	54
8	H	185/190 (97%)	181 (98%)	4 (2%)	52	83
8	V	185/190 (97%)	181 (98%)	4 (2%)	52	83
9	I	172/173 (99%)	168 (98%)	4 (2%)	50	82
9	W	172/173 (99%)	168 (98%)	4 (2%)	50	82
10	J	173/175 (99%)	169 (98%)	4 (2%)	50	82
10	X	173/175 (99%)	169 (98%)	4 (2%)	50	82
11	K	169/169 (100%)	160 (95%)	9 (5%)	22	54
11	Y	169/169 (100%)	160 (95%)	9 (5%)	22	54
12	L	185/185 (100%)	179 (97%)	6 (3%)	39	73
12	Z	185/185 (100%)	178 (96%)	7 (4%)	33	67
13	M	190/203 (94%)	184 (97%)	6 (3%)	39	73
13	a	190/203 (94%)	184 (97%)	6 (3%)	39	73
14	N	162/162 (100%)	157 (97%)	5 (3%)	40	74
14	b	162/162 (100%)	158 (98%)	4 (2%)	47	80
All	All	5302/5530 (96%)	5084 (96%)	218 (4%)	30	64

5 of 218 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
13	M	2	GLN
3	Q	38	ASN
12	Z	3	ASN
13	M	104	ARG
1	O	122	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 117 such sidechains are listed below:

Mol	Chain	Res	Type
12	L	70	ASN
2	P	176	GLN
12	Z	49	ASN
13	M	48	ASN
1	O	94	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 11 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	250/250 (100%)	-0.47	6 (2%) 59 49	42, 54, 87, 126	0
1	O	250/250 (100%)	-0.41	5 (2%) 65 56	47, 63, 104, 132	0
2	B	244/258 (94%)	-0.39	5 (2%) 65 56	42, 60, 100, 150	0
2	P	244/258 (94%)	-0.35	9 (3%) 41 31	48, 63, 106, 153	0
3	C	240/254 (94%)	-0.29	8 (3%) 46 36	43, 64, 122, 148	0
3	Q	240/254 (94%)	-0.07	14 (5%) 23 15	48, 76, 154, 178	0
4	D	235/260 (90%)	-0.46	2 (0%) 84 80	47, 66, 97, 138	0
4	R	235/260 (90%)	-0.33	4 (1%) 70 63	50, 70, 111, 142	0
5	E	231/234 (98%)	-0.32	1 (0%) 92 91	51, 70, 107, 150	0
5	S	231/234 (98%)	-0.19	4 (1%) 70 63	53, 78, 117, 160	0
6	F	243/288 (84%)	-0.47	3 (1%) 79 73	46, 62, 112, 139	0
6	T	243/288 (84%)	-0.33	6 (2%) 57 47	49, 73, 123, 151	0
7	G	241/252 (95%)	-0.56	0 100 100	36, 56, 87, 137	0
7	U	241/252 (95%)	-0.44	1 (0%) 92 91	48, 61, 89, 132	0
8	H	226/232 (97%)	-0.37	7 (3%) 49 39	37, 56, 84, 147	0
8	V	226/232 (97%)	-0.30	6 (2%) 54 44	45, 59, 89, 159	0
9	I	204/205 (99%)	-0.71	1 (0%) 91 88	40, 51, 77, 100	0
9	W	204/205 (99%)	-0.71	3 (1%) 73 68	42, 53, 81, 105	0
10	J	195/198 (98%)	-0.57	2 (1%) 82 77	40, 54, 78, 134	0
10	X	195/198 (98%)	-0.55	3 (1%) 73 68	41, 55, 80, 142	0
11	K	212/212 (100%)	-0.60	0 100 100	37, 54, 75, 94	0
11	Y	212/212 (100%)	-0.58	0 100 100	43, 55, 76, 97	0
12	L	222/222 (100%)	-0.56	2 (0%) 84 80	42, 57, 84, 122	0
12	Z	222/222 (100%)	-0.59	1 (0%) 91 88	42, 56, 85, 121	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
13	M	222/239 (92%)	-0.57	4 (1%) 68 61	39, 56, 81, 100	0
13	a	222/239 (92%)	-0.56	3 (1%) 75 70	40, 56, 78, 97	0
14	N	196/196 (100%)	-0.65	2 (1%) 82 77	39, 51, 79, 108	0
14	b	196/196 (100%)	-0.60	2 (1%) 82 77	41, 53, 79, 114	0
All	All	6322/6600 (95%)	-0.46	104 (1%) 72 66	36, 60, 103, 178	0

The worst 5 of 104 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
8	V	224	GLN	7.1
3	Q	236	GLN	6.1
3	Q	49	THR	5.6
2	P	221	ASP	5.5
2	P	220	ASN	5.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
15	MG	Z	301	1/1	0.81	0.51	78,78,78,78	0
15	MG	H	301	1/1	0.81	0.13	57,57,57,57	0
15	MG	I	301	1/1	0.88	0.11	50,50,50,50	0
15	MG	K	301	1/1	0.94	0.12	53,53,53,53	0
15	MG	V	301	1/1	0.97	0.06	57,57,57,57	0
15	MG	W	301	1/1	0.97	0.17	57,57,57,57	0
15	MG	G	301	1/1	0.97	0.05	47,47,47,47	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
16	CL	U	301	1/1	0.98	0.23	30,30,30,30	0
15	MG	N	201	1/1	0.98	0.14	52,52,52,52	0
16	CL	G	302	1/1	0.98	0.25	30,30,30,30	0
15	MG	Y	301	1/1	0.99	0.07	50,50,50,50	0

6.5 Other polymers [i](#)

There are no such residues in this entry.