



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 14, 2020 – 12:15 AM BST

PDB ID : 5YBG
Title : Crystal structure of the GluA2o LBD in complex with glutamate and LY451395
Authors : Sogabe, S.; Igaki, S.; Hirokawa, A.; Zama, Y.; Lane, W.; Snell, G.
Deposited on : 2017-09-04
Resolution : 1.52 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.14.4.dev1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.14.4.dev1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.52 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4009 (1.54-1.50)
Clashscore	141614	4249 (1.54-1.50)
Ramachandran outliers	138981	4148 (1.54-1.50)
Sidechain outliers	138945	4146 (1.54-1.50)
RSRZ outliers	127900	3943 (1.54-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	263	<div> <div>13%</div> <div> <div></div> <div>94%</div> <div>••</div> </div> </div>
1	B	263	<div> <div>2%</div> <div> <div></div> <div>94%</div> <div>5% •</div> </div> </div>
1	C	263	<div> <div>2%</div> <div> <div></div> <div>93%</div> <div>6% •</div> </div> </div>
1	D	263	<div> <div>•</div> <div> <div></div> <div>91%</div> <div>8% •</div> </div> </div>
1	E	263	<div> <div>7%</div> <div> <div></div> <div>93%</div> <div>6% •</div> </div> </div>
1	F	263	<div> <div>3%</div> <div> <div></div> <div>92%</div> <div>6% ••</div> </div> </div>

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 13863 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamate receptor 2, Glutamate receptor 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	260	Total	C	N	O	S	0	3	0
			2053	1307	345	386	15			
1	B	260	Total	C	N	O	S	0	4	0
			2051	1306	342	387	16			
1	C	259	Total	C	N	O	S	0	6	0
			2063	1315	346	387	15			
1	D	260	Total	C	N	O	S	0	5	0
			2060	1311	347	387	15			
1	E	261	Total	C	N	O	S	0	5	0
			2065	1316	346	388	15			
1	F	260	Total	C	N	O	S	0	4	0
			2053	1308	342	387	16			

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	411	GLY	-	expression tag	UNP P42262
A	412	SER	-	expression tag	UNP P42262
A	641	GLY	-	linker	UNP P42262
A	642	THR	-	linker	UNP P42262
B	411	GLY	-	expression tag	UNP P42262
B	412	SER	-	expression tag	UNP P42262
B	641	GLY	-	linker	UNP P42262
B	642	THR	-	linker	UNP P42262
C	411	GLY	-	expression tag	UNP P42262
C	412	SER	-	expression tag	UNP P42262
C	641	GLY	-	linker	UNP P42262
C	642	THR	-	linker	UNP P42262
D	411	GLY	-	expression tag	UNP P42262
D	412	SER	-	expression tag	UNP P42262
D	641	GLY	-	linker	UNP P42262
D	642	THR	-	linker	UNP P42262
E	411	GLY	-	expression tag	UNP P42262

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	412	SER	-	expression tag	UNP P42262
E	641	GLY	-	linker	UNP P42262
E	642	THR	-	linker	UNP P42262
F	411	GLY	-	expression tag	UNP P42262
F	412	SER	-	expression tag	UNP P42262
F	641	GLY	-	linker	UNP P42262
F	642	THR	-	linker	UNP P42262

- Molecule 2 is GLUTAMIC ACID (three-letter code: GLU) (formula: C₅H₉NO₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			10	5	1	4		
2	B	1	Total	C	N	O	0	0
			10	5	1	4		
2	C	1	Total	C	N	O	0	0
			10	5	1	4		
2	D	1	Total	C	N	O	0	0
			10	5	1	4		
2	E	1	Total	C	N	O	0	0
			10	5	1	4		
2	F	1	Total	C	N	O	0	0
			10	5	1	4		

- Molecule 3 is N-[(2R)-2-[4-[4-[2-(methylsulfonylamino)ethyl]phenyl]phenyl]propyl]propane-2-sulfonamide (three-letter code: 8SO) (formula: C₂₁H₃₀N₂O₄S₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	1
			58	42	4	8	4		
3	C	1	Total	C	N	O	S	0	1
			58	42	4	8	4		
3	E	1	Total	C	N	O	S	0	1
			58	42	4	8	4		

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	3	Total	Zn	0	0
			3	3		
4	E	1	Total	Zn	0	0
			1	1		
4	B	1	Total	Zn	0	0
			1	1		
4	C	2	Total	Zn	0	0
			2	2		
4	A	2	Total	Zn	0	0
			2	2		
4	F	1	Total	Zn	0	0
			1	1		

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	C	1	Total	C	O	0	0
			4	2	2		
5	C	1	Total	C	O	0	0
			4	2	2		
5	D	1	Total	C	O	0	0
			4	2	2		
5	E	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	E	1	Total	C	O	0	0
			6	3	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	153	Total	O	0	0
			153	153		
7	B	194	Total	O	0	0
			194	194		
7	C	296	Total	O	0	0
			296	296		
7	D	296	Total	O	0	0
			296	296		
7	E	142	Total	O	0	0
			142	142		
7	F	161	Total	O	0	0
			161	161		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glutamate receptor 2,Glutamate receptor 2




- Molecule 1: Glutamate receptor 2,Glutamate receptor 2




- Molecule 1: Glutamate receptor 2,Glutamate receptor 2




- Molecule 1: Glutamate receptor 2,Glutamate receptor 2




- Molecule 1: Glutamate receptor 2,Glutamate receptor 2





● Molecule 1: Glutamate receptor 2,Glutamate receptor 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	113.72Å 161.91Å 46.96Å 90.00° 89.97° 90.00°	Depositor
Resolution (Å)	40.00 – 1.52 41.93 – 1.52	Depositor EDS
% Data completeness (in resolution range)	99.4 (40.00-1.52) 99.4 (41.93-1.52)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.05 (at 1.52Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.202 , 0.233 0.204 , 0.206	Depositor DCC
R_{free} test set	12981 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	20.1	Xtriage
Anisotropy	0.141	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 29.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.469 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	13863	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ZN, 8SO, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.51	0/2098	0.71	1/2819 (0.0%)
1	B	0.50	0/2101	0.68	0/2823
1	C	0.58	0/2117	0.78	0/2844
1	D	0.58	0/2115	0.79	0/2841
1	E	0.55	2/2120 (0.1%)	0.70	0/2849
1	F	0.50	0/2101	0.69	0/2823
All	All	0.54	2/12652 (0.0%)	0.73	1/16999 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	699	GLU	CD-OE2	6.72	1.33	1.25
1	E	699	GLU	CG-CD	5.35	1.59	1.51

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	699	GLU	OE1-CD-OE2	-5.10	117.18	123.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2053	0	2096	10	0
1	B	2051	0	2093	9	0
1	C	2063	0	2117	13	0
1	D	2060	0	2103	20	0
1	E	2065	0	2115	9	0
1	F	2053	0	2097	12	0
2	A	10	0	5	0	0
2	B	10	0	5	1	0
2	C	10	0	5	1	0
2	D	10	0	5	1	0
2	E	10	0	5	1	0
2	F	10	0	5	1	0
3	A	58	0	0	5	0
3	C	58	0	0	1	0
3	E	58	0	0	2	0
4	A	2	0	0	0	0
4	B	1	0	0	0	0
4	C	2	0	0	0	0
4	D	3	0	0	0	0
4	E	1	0	0	0	0
4	F	1	0	0	0	0
5	A	4	0	3	0	0
5	C	8	0	6	0	0
5	D	4	0	3	0	0
5	E	4	0	3	0	0
6	A	6	0	8	1	0
6	E	6	0	8	2	0
7	A	153	0	0	1	0
7	B	194	0	0	2	0
7	C	296	0	0	3	0
7	D	296	0	0	5	0
7	E	142	0	0	2	0
7	F	161	0	0	1	0
All	All	13863	0	12682	82	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 82 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:736[B]:ARG:HH21	1:C:736[B]:ARG:HG2	0.89	1.02

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:736[B]:ARG:HH11	1:A:736[B]:ARG:HG2	0.89	1.01
1:D:736[B]:ARG:NH1	1:D:736[B]:ARG:HG2	1.64	1.01
1:A:736[B]:ARG:HG2	1:A:736[B]:ARG:NH1	1.69	1.00
1:D:736[B]:ARG:HH11	1:D:736[B]:ARG:CG	1.75	0.99

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	261/263 (99%)	256 (98%)	5 (2%)	0	100	100
1	B	262/263 (100%)	260 (99%)	2 (1%)	0	100	100
1	C	263/263 (100%)	258 (98%)	5 (2%)	0	100	100
1	D	263/263 (100%)	260 (99%)	3 (1%)	0	100	100
1	E	264/263 (100%)	258 (98%)	6 (2%)	0	100	100
1	F	262/263 (100%)	258 (98%)	4 (2%)	0	100	100
All	All	1575/1578 (100%)	1550 (98%)	25 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	222/221 (100%)	219 (99%)	3 (1%)	67	41
1	B	223/221 (101%)	221 (99%)	2 (1%)	78	60
1	C	225/221 (102%)	223 (99%)	2 (1%)	78	60
1	D	224/221 (101%)	223 (100%)	1 (0%)	91	82
1	E	225/221 (102%)	222 (99%)	3 (1%)	69	43
1	F	223/221 (101%)	220 (99%)	3 (1%)	69	43
All	All	1342/1326 (101%)	1328 (99%)	14 (1%)	73	56

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	519	LEU
1	D	519	LEU
1	F	432	ASN
1	C	477	ASP
1	E	664	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	775	ASN
1	D	785	ASN
1	F	432	ASN
1	F	765	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 29 ligands modelled in this entry, 10 are monoatomic - leaving 19 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	8SO	A	902[B]	-	29,30,30	1.68	3 (10%)	37,43,43	2.92	5 (13%)
6	GOL	A	905	-	5,5,5	0.14	0	5,5,5	0.56	0
5	ACT	C	805	-	1,3,3	1.39	0	0,3,3	0.00	-
6	GOL	E	905	-	5,5,5	0.16	0	5,5,5	0.46	0
2	GLU	B	901	-	2,9,9	0.09	0	2,11,11	0.29	0
2	GLU	A	901	-	2,9,9	0.17	0	2,11,11	0.10	0
2	GLU	D	901	-	2,9,9	0.11	0	2,11,11	0.68	0
3	8SO	C	802[B]	-	29,30,30	1.77	3 (10%)	37,43,43	3.50	9 (24%)
2	GLU	F	901	-	2,9,9	0.23	0	2,11,11	0.29	0
2	GLU	E	901	-	2,9,9	0.19	0	2,11,11	0.08	0
3	8SO	E	902[B]	-	29,30,30	1.68	3 (10%)	37,43,43	2.91	5 (13%)
3	8SO	E	902[A]	-	29,30,30	1.60	3 (10%)	37,43,43	3.22	3 (8%)
3	8SO	C	802[A]	-	29,30,30	1.68	3 (10%)	37,43,43	3.13	8 (21%)
5	ACT	D	904	-	1,3,3	1.45	0	0,3,3	0.00	-
2	GLU	C	801	-	2,9,9	0.46	0	2,11,11	0.56	0
3	8SO	A	902[A]	-	29,30,30	1.69	3 (10%)	37,43,43	3.07	5 (13%)
5	ACT	C	806	-	1,3,3	1.14	0	0,3,3	0.00	-
5	ACT	A	904	4	1,3,3	1.47	0	0,3,3	0.00	-
5	ACT	E	904	4	1,3,3	1.31	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	A	905	-	-	2/4/4/4	-
6	GOL	E	905	-	-	2/4/4/4	-
2	GLU	B	901	-	-	0/3/9/9	-
2	GLU	A	901	-	-	0/3/9/9	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLU	D	901	-	-	0/3/9/9	-
2	GLU	F	901	-	-	0/3/9/9	-
2	GLU	E	901	-	-	0/3/9/9	-
3	8SO	C	802[A]	-	-	7/25/27/27	0/2/2/2
3	8SO	E	902[B]	-	-	5/25/27/27	0/2/2/2
3	8SO	E	902[A]	-	-	7/25/27/27	0/2/2/2
2	GLU	C	801	-	-	0/3/9/9	-
3	8SO	A	902[A]	-	-	8/25/27/27	0/2/2/2
3	8SO	A	902[B]	-	-	7/25/27/27	0/2/2/2
3	8SO	C	802[B]	-	-	0/25/27/27	0/2/2/2

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	802[B]	8SO	S28-N22	6.31	1.73	1.60
3	C	802[A]	8SO	S28-N22	6.18	1.73	1.60
3	A	902[A]	8SO	S28-N22	6.01	1.72	1.60
3	A	902[B]	8SO	S28-N22	5.96	1.72	1.60
3	E	902[A]	8SO	S28-N22	5.94	1.72	1.60

The worst 5 of 35 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	802[B]	8SO	C16-S28-N22	-18.54	96.41	107.76
3	E	902[A]	8SO	C16-S28-N22	-17.94	96.77	107.76
3	A	902[A]	8SO	C16-S28-N22	-16.22	97.83	107.76
3	C	802[A]	8SO	C16-S28-N22	-16.19	97.84	107.76
3	A	902[B]	8SO	C16-S28-N22	-15.30	98.39	107.76

There are no chirality outliers.

5 of 38 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	E	905	GOL	C1-C2-C3-O3
3	E	902[A]	8SO	C18-N22-S28-C16
3	E	902[A]	8SO	C18-N22-S28-O24
3	E	902[A]	8SO	C18-N22-S28-O25
3	E	902[B]	8SO	C7-C12-C20-C19





There are no ring outliers.


12 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	902[B]	8SO	4	0
6	A	905	GOL	1	0
6	E	905	GOL	2	0
2	B	901	GLU	1	0
2	D	901	GLU	1	0
2	F	901	GLU	1	0
2	E	901	GLU	1	0
3	E	902[B]	8SO	1	0
3	E	902[A]	8SO	1	0
3	C	802[A]	8SO	1	0
2	C	801	GLU	1	0
3	A	902[A]	8SO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	260/263 (98%)	0.64	33 (12%) 3 3	16, 29, 54, 70	0
1	B	260/263 (98%)	0.14	4 (1%) 73 78	16, 28, 49, 68	0
1	C	259/263 (98%)	-0.05	5 (1%) 66 71	12, 19, 35, 59	0
1	D	260/263 (98%)	-0.10	3 (1%) 79 82	12, 19, 35, 61	0
1	E	261/263 (99%)	0.43	19 (7%) 15 16	16, 28, 53, 66	0
1	F	260/263 (98%)	0.22	7 (2%) 54 59	17, 28, 49, 78	0
All	All	1560/1578 (98%)	0.21	71 (4%) 32 35	12, 25, 49, 78	0

The worst 5 of 71 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	789	TYR	10.8
1	F	792	GLY	6.9
1	A	792	GLY	6.7
1	E	792	GLY	5.9
1	A	794	CYS	5.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	ACT	C	805	4/4	0.71	0.16	25,27,32,36	0
6	GOL	E	905	6/6	0.79	0.26	26,33,39,44	0
5	ACT	C	806	4/4	0.81	0.18	33,33,40,47	0
6	GOL	A	905	6/6	0.84	0.26	27,35,37,41	0
3	8SO	E	902[A]	29/29	0.88	0.17	18,23,30,34	29
3	8SO	E	902[B]	29/29	0.88	0.17	23,25,37,39	29
4	ZN	B	902	1/1	0.88	0.13	46,46,46,46	0
3	8SO	A	902[B]	29/29	0.89	0.17	19,24,29,32	29
3	8SO	A	902[A]	29/29	0.89	0.17	25,29,33,34	29
5	ACT	D	904	4/4	0.90	0.13	23,28,29,35	0
5	ACT	A	904	4/4	0.92	0.15	32,40,40,41	0
5	ACT	E	904	4/4	0.93	0.14	35,40,43,44	0
3	8SO	C	802[B]	29/29	0.93	0.13	14,20,25,28	29
3	8SO	C	802[A]	29/29	0.93	0.13	17,20,37,41	29
4	ZN	F	902	1/1	0.94	0.09	46,46,46,46	0
2	GLU	A	901	10/10	0.95	0.08	17,20,22,22	0
4	ZN	E	903	1/1	0.96	0.06	28,28,28,28	0
2	GLU	B	901	10/10	0.97	0.08	15,16,19,20	0
2	GLU	E	901	10/10	0.97	0.06	18,20,23,23	0
4	ZN	C	804	1/1	0.97	0.06	21,21,21,21	0
2	GLU	F	901	10/10	0.98	0.07	15,16,20,20	0
4	ZN	A	903	1/1	0.98	0.05	29,29,29,29	0
4	ZN	A	906	1/1	0.98	0.07	18,18,18,18	0
2	GLU	D	901	10/10	0.98	0.07	12,13,13,13	0
4	ZN	D	903	1/1	0.98	0.06	21,21,21,21	0
4	ZN	C	803	1/1	0.98	0.05	21,21,21,21	0
2	GLU	C	801	10/10	0.98	0.08	11,13,13,14	0
4	ZN	D	902	1/1	0.99	0.05	21,21,21,21	0
4	ZN	D	905	1/1	0.99	0.06	18,18,18,18	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 8SO E 902 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 8SO E 902 (B):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 8SO A 902 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 8SO A 902 (A):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 8SO C 802 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around 8SO C 802 (A):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.