



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 23, 2020 – 07:07 pm BST

PDB ID : 4YCO  
Title : E. coli dihydrouridine synthase C (DusC) in complex with tRNAPhe  
Authors : Byrne, R.T.; Jenkins, H.T.; Peters, D.T.; Whelan, F.; Stowell, J.; Aziz, N.;  
Kasatsky, P.; Rodnina, M.V.; Koonin, E.V.; Konevega, A.L.; Antson, A.A.  
Deposited on : 2015-02-20  
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)
RNA backbone	3102	1000 (2.54-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	321	<div> <div></div> <div>90% 6% .</div> </div>
1	B	321	<div> <div>2%</div> <div>90% 6% .</div> </div>
1	C	321	<div> <div></div> <div>93% . .</div> </div>
2	D	76	<div> <div>3%</div> <div>72% 18% 7% .</div> </div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	E	76	<div><div>5%</div><div><div></div><div>67%</div><div>18%</div><div>•</div><div>12%</div></div></div>
2	F	76	<div><div>%</div><div><div></div><div>62%</div><div>25%</div><div>7%</div><div>7%</div></div></div>

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 12665 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called tRNA-dihydrouridine synthase C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	311	Total	C	N	O	S	0	7	0
			2473	1566	450	446	11			
1	B	309	Total	C	N	O	S	0	5	0
			2447	1550	443	443	11			
1	C	311	Total	C	N	O	S	0	7	0
			2477	1568	451	447	11			

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	GLY	-	expression tag	UNP P33371
A	-4	SER	-	expression tag	UNP P33371
A	-3	HIS	-	expression tag	UNP P33371
A	-2	MET	-	expression tag	UNP P33371
A	-1	ALA	-	expression tag	UNP P33371
A	0	SER	-	expression tag	UNP P33371
A	98	ALA	CYS	engineered mutation	UNP P33371
B	-5	GLY	-	expression tag	UNP P33371
B	-4	SER	-	expression tag	UNP P33371
B	-3	HIS	-	expression tag	UNP P33371
B	-2	MET	-	expression tag	UNP P33371
B	-1	ALA	-	expression tag	UNP P33371
B	0	SER	-	expression tag	UNP P33371
B	98	ALA	CYS	engineered mutation	UNP P33371
C	-5	GLY	-	expression tag	UNP P33371
C	-4	SER	-	expression tag	UNP P33371
C	-3	HIS	-	expression tag	UNP P33371
C	-2	MET	-	expression tag	UNP P33371
C	-1	ALA	-	expression tag	UNP P33371
C	0	SER	-	expression tag	UNP P33371
C	98	ALA	CYS	engineered mutation	UNP P33371


- Molecule 2 is a RNA chain called tRNAp<sup>he</sup>.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	74	Total	C	N	O	P	0	0	0
			1581	704	282	521	74			
2	E	67	Total	C	N	O	P	0	0	0
			1433	638	255	473	67			
2	F	71	Total	C	N	O	P	0	0	0
			1519	676	271	501	71			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	3	G	C	engineered mutation	GB 731469900
D	70	C	G	engineered mutation	GB 731469900
E	3	G	C	engineered mutation	GB 731469900
E	70	C	G	engineered mutation	GB 731469900
F	3	G	C	engineered mutation	GB 731469900
F	70	C	G	engineered mutation	GB 731469900

- Molecule 3 is FLAVIN MONONUCLEOTIDE (three-letter code: FMN) (formula:  $C_{17}H_{21}N_4O_9P$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			31	17	4	9	1		
3	B	1	Total	C	N	O	P	0	0
			31	17	4	9	1		
3	C	1	Total	C	N	O	P	0	0
			31	17	4	9	1		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total 1	Mg 1	0	0
4	D	8	Total 8	Mg 8	0	0
4	F	8	Total 8	Mg 8	0	0
4	E	8	Total 8	Mg 8	0	0

- Molecule 5 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	D	1	Total 1	Mn 1	0	0
5	F	1	Total 1	Mn 1	0	0
5	E	1	Total 1	Mn 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	122	Total 122	O 122	0	0
6	B	89	Total 89	O 89	0	0
6	C	148	Total 148	O 148	0	1
6	D	95	Total 95	O 95	0	0
6	E	67	Total 67	O 67	0	0
6	F	93	Total 93	O 93	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: tRNA-dihydrouridine synthase C




- Molecule 1: tRNA-dihydrouridine synthase C




- Molecule 1: tRNA-dihydrouridine synthase C




- Molecule 2: tRNAp<sup>he</sup>



- Molecule 2: tRNAp<sup>he</sup>



- Molecule 2: tRNAp<sup>he</sup>





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	100.58Å 176.90Å 238.41Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.21 – 2.10 49.21 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.9 (49.21-2.10) 99.9 (49.21-2.10)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.01 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.192 , 0.223 0.194 , 0.223	Depositor DCC
$R_{free}$ test set	1241 reflections (1.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	39.5	Xtriage
Anisotropy	0.453	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 40.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.019 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.029 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	12665	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.19% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FMN, MG, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.53	0/2542	0.73	2/3449 (0.1%)
1	B	0.48	0/2510	0.65	0/3407
1	C	0.52	0/2546	0.72	1/3454 (0.0%)
2	D	0.45	1/1766 (0.1%)	0.74	1/2750 (0.0%)
2	E	0.44	1/1601 (0.1%)	0.72	2/2493 (0.1%)
2	F	0.44	1/1697 (0.1%)	0.75	2/2643 (0.1%)
All	All	0.49	3/12662 (0.0%)	0.72	8/18196 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	3	G	OP3-P	-10.27	1.48	1.61
2	D	1	G	OP3-P	-10.14	1.49	1.61
2	F	1	G	OP3-P	-10.12	1.49	1.61

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	35	ARG	NE-CZ-NH1	-8.62	115.99	120.30
2	F	53	G	O5'-P-OP2	-7.31	99.12	105.70
1	A	35	ARG	NE-CZ-NH2	7.26	123.93	120.30
2	D	53	G	O5'-P-OP2	-6.87	99.52	105.70
2	E	9	A	O5'-P-OP2	-6.77	99.61	105.70

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2473	0	2504	15	0
1	B	2447	0	2476	10	0
1	C	2477	0	2510	12	0
2	D	1581	0	799	9	0
2	E	1433	0	722	5	0
2	F	1519	0	766	16	0
3	A	31	0	19	1	0
3	B	31	0	19	0	0
3	C	31	0	19	0	0
4	B	1	0	0	0	0
4	D	8	0	0	0	0
4	E	8	0	0	0	0
4	F	8	0	0	0	0
5	D	1	0	0	0	0
5	E	1	0	0	0	0
5	F	1	0	0	0	0
6	A	122	0	0	1	0
6	B	89	0	0	1	0
6	C	148	0	0	2	0
6	D	95	0	0	1	0
6	E	67	0	0	0	0
6	F	93	0	0	2	0
All	All	12665	0	9834	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 56 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:207:GLN:OE1	2:D:29:G:O2'	2.11	0.69
1:A:35:ARG:HG2	2:D:14:A:O2'	1.96	0.66
2:F:38:A:N1	6:F:270:HOH:O	2.30	0.64
2:D:47:U:H2'	2:D:50:U:OP1	2.01	0.60
1:A:288:THR:OG1	6:A:617:HOH:O	2.17	0.59

There are no symmetry-related clashes.

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	314/321 (98%)	304 (97%)	10 (3%)	0	100	100
1	B	310/321 (97%)	303 (98%)	7 (2%)	0	100	100
1	C	314/321 (98%)	308 (98%)	6 (2%)	0	100	100
All	All	938/963 (97%)	915 (98%)	23 (2%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	261/267 (98%)	256 (98%)	5 (2%)	57	63
1	B	259/267 (97%)	255 (98%)	4 (2%)	65	71
1	C	262/267 (98%)	260 (99%)	2 (1%)	81	86
All	All	782/801 (98%)	771 (99%)	11 (1%)	69	73

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	192	ARG
1	B	35	ARG
1	B	294	VAL

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	146	SER
1	B	100	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	D	73/76 (96%)	12 (16%)	0
2	E	66/76 (86%)	11 (16%)	3 (4%)
2	F	70/76 (92%)	9 (12%)	2 (2%)
All	All	209/228 (91%)	32 (15%)	5 (2%)

5 of 32 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	D	3	G
2	D	4	C
2	D	16	U
2	D	19	G
2	D	21	A

All (5) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
2	E	44	G
2	E	45	U
2	E	47	U
2	F	47	U
2	F	48	C

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 31 ligands modelled in this entry, 28 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	FMN	A	401	-	31,33,33	1.42	4 (12%)	40,50,50	2.65	6 (15%)
3	FMN	B	402	-	31,33,33	1.52	6 (19%)	40,50,50	2.61	6 (15%)
3	FMN	C	401	-	31,33,33	1.45	6 (19%)	40,50,50	2.59	5 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FMN	A	401	-	-	1/18/18/18	0/3/3/3
3	FMN	B	402	-	-	1/18/18/18	0/3/3/3
3	FMN	C	401	-	-	1/18/18/18	0/3/3/3

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	402	FMN	C4A-C10	5.15	1.44	1.38
3	A	401	FMN	C4A-C10	4.84	1.43	1.38
3	C	401	FMN	C4A-C10	3.24	1.42	1.38
3	C	401	FMN	C4-N3	3.13	1.38	1.33
3	C	401	FMN	C9A-N10	3.10	1.42	1.38

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	401	FMN	C4-N3-C2	12.82	125.96	115.14
3	C	401	FMN	C4-N3-C2	12.69	125.86	115.14
3	B	402	FMN	C4-N3-C2	12.61	125.78	115.14
3	B	402	FMN	C4A-C4-N3	-7.22	113.55	123.43

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	401	FMN	C4A-C4-N3	-6.76	114.18	123.43

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	402	FMN	C4'-C5'-O5'-P
3	C	401	FMN	C4'-C5'-O5'-P
3	A	401	FMN	C4'-C5'-O5'-P

There are no ring outliers.


1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	401	FMN	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	311/321 (96%)	-0.21	0 100 100	25, 37, 55, 77	0
1	B	309/321 (96%)	-0.18	5 (1%) 72 75	29, 42, 66, 81	0
1	C	311/321 (96%)	-0.11	1 (0%) 94 94	24, 34, 55, 79	0
2	D	74/76 (97%)	-0.39	2 (2%) 54 60	33, 47, 120, 131	0
2	E	67/76 (88%)	-0.32	4 (5%) 21 27	42, 55, 190, 235	0
2	F	71/76 (93%)	-0.34	1 (1%) 75 78	34, 48, 108, 142	0
All	All	1143/1191 (95%)	-0.20	13 (1%) 80 84	24, 40, 77, 235	0

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	E	48	C	4.7
1	B	106	SER	4.5
2	E	46	G	4.1
1	B	312	ILE	2.9
1	B	105	GLY	2.7

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	MG	E	107	1/1	0.55	0.20	79,79,79,79	0
4	MG	E	102	1/1	0.59	0.13	64,64,64,64	0
4	MG	D	105	1/1	0.70	0.17	61,61,61,61	0
4	MG	F	108	1/1	0.78	0.24	63,63,63,63	0
4	MG	E	104	1/1	0.79	0.07	58,58,58,58	0
4	MG	D	104	1/1	0.80	0.12	58,58,58,58	0
4	MG	D	102	1/1	0.82	0.06	50,50,50,50	0
4	MG	F	106	1/1	0.82	0.07	59,59,59,59	0
4	MG	F	104	1/1	0.82	0.22	66,66,66,66	0
4	MG	D	103	1/1	0.85	0.18	66,66,66,66	0
4	MG	E	103	1/1	0.85	0.06	56,56,56,56	0
4	MG	F	102	1/1	0.85	0.21	59,59,59,59	0
4	MG	E	109	1/1	0.87	0.24	56,56,56,56	0
4	MG	D	109	1/1	0.87	0.12	53,53,53,53	0
4	MG	E	105	1/1	0.88	0.15	59,59,59,59	0
4	MG	F	103	1/1	0.89	0.06	53,53,53,53	0
4	MG	F	107	1/1	0.89	0.06	47,47,47,47	0
4	MG	D	107	1/1	0.91	0.23	39,39,39,39	0
4	MG	E	106	1/1	0.94	0.19	46,46,46,46	0
4	MG	B	401	1/1	0.94	0.05	69,69,69,69	0
4	MG	F	109	1/1	0.95	0.19	37,37,37,37	0
4	MG	D	106	1/1	0.95	0.05	48,48,48,48	0
4	MG	E	108	1/1	0.95	0.03	50,50,50,50	0
4	MG	D	108	1/1	0.95	0.14	56,56,56,56	0
3	FMN	B	402	31/31	0.98	0.11	31,35,37,38	0
5	MN	E	101	1/1	0.98	0.03	63,63,63,63	0
3	FMN	C	401	31/31	0.98	0.14	24,28,30,31	0
5	MN	F	101	1/1	0.98	0.03	56,56,56,56	0
3	FMN	A	401	31/31	0.98	0.12	26,29,32,33	0
4	MG	F	105	1/1	0.99	0.07	43,43,43,43	0
5	MN	D	101	1/1	1.00	0.04	50,50,50,50	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around FMN B 402:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around FMN C 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around FMN A 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.