



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 16, 2020 – 06:15 PM GMT

PDB ID : 6YCG  
Title : Structure the bromelain protease from Ananas comosus in complex with the TLCK inhibitor  
Authors : Azarkan, M.; Charlier, P.; Herman, R.; Delbrassine, F.; Sauvage, E.; M Rabet, N.; Calvo Esposito, R.; Kerff, F.  
Deposited on : 2020-03-18  
Resolution : 1.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.14.6  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.14.6


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1156 (1.46-1.46)
Clashscore	141614	1202 (1.46-1.46)
Ramachandran outliers	138981	1178 (1.46-1.46)
Sidechain outliers	138945	1178 (1.46-1.46)
RSRZ outliers	127900	1139 (1.46-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	219	<div> <div>2%</div> <div> <div></div> <div>93%</div> <div>6%</div> </div> </div>
1	B	219	<div> <div>3%</div> <div> <div></div> <div>90%</div> <div>8%</div> </div> </div>
2	C	6	<div> <div>17%</div> <div>83%</div> </div>
3	D	3	<div> <div>33%</div> <div>67%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	XYS	C	4	X	-	-	-

## 2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 4010 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FBSB.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	215	Total	C	N	O	S	0	15	0
			1705	1088	284	322	11			
1	B	215	Total	C	N	O	S	0	12	0
			1697	1079	287	320	11			

- Molecule 2 is an oligosaccharide called alpha-D-xylopyranose-(1-2)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-3)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	6	Total	C	N	O	0	0	0
			69	39	2	28			

- Molecule 3 is an oligosaccharide called alpha-L-fucopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	D	3	Total	C	N	O	0	0	0
			38	22	2	14			

- Molecule 4 is ISOPROPYL ALCOHOL (three-letter code: IPA) (formula: C<sub>3</sub>H<sub>8</sub>O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	3	1		
4	A	1	Total	C	O	0	0
			4	3	1		


- Molecule 5 is CITRIC ACID (three-letter code: CIT) (formula:  $C_6H_8O_7$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			13	6	7		

- Molecule 6 is N-[(1S)-5-amino-1-(chloroacetyl)pentyl]-4-methylbenzenesulfonamid

e (three-letter code: TCK) (formula:  $C_{14}H_{21}ClN_2O_3S$ ) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	B	1	Total	C	N	O	S	0	1
			40	28	4	6	2		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	233	Total	O	0	1
			233	233		
7	B	207	Total	O	0	4
			207	207		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: FBSB



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	91.24Å 137.23Å 82.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.62 – 1.45 45.62 – 1.45	Depositor EDS
% Data completeness (in resolution range)	99.9 (45.62-1.45) 99.9 (45.62-1.45)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.22 (at 1.45Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, $R_{free}$	0.115 , 0.155 0.131 , 0.166	Depositor DCC
$R_{free}$ test set	4590 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	14.1	Xtriage
Anisotropy	0.219	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 48.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	4010	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.25% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: YYS, IPA, NAG, BMA, FUC, CIT, MAN, TCK

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.73	0/1782	0.87	3/2410 (0.1%)
1	B	0.72	0/1760	0.88	5/2379 (0.2%)
All	All	0.73	0/3542	0.87	8/4789 (0.2%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	210[B]	PRO	CA-N-CD	-9.14	98.70	111.50
1	A	210[B]	PRO	N-CA-CB	9.04	114.15	103.30
1	A	210[B]	PRO	N-CD-CG	6.88	113.52	103.20
1	B	210[B]	PRO	N-CA-CB	6.65	111.28	103.30
1	B	115	ARG	NE-CZ-NH2	6.53	123.56	120.30
1	B	210[B]	PRO	CA-N-CD	-6.22	102.79	111.50
1	B	210[B]	PRO	N-CD-CG	5.62	111.63	103.20
1	B	135	ASP	CB-CG-OD2	-5.26	113.56	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1705	0	1674	13	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1697	0	1659	11	0
2	C	69	0	59	0	0
3	D	38	0	34	1	0
4	A	8	0	16	0	0
5	A	13	0	5	2	0
6	B	40	0	38	2	0
7	A	233	0	0	4	1
7	B	207	0	0	1	1
All	All	4010	0	3485	27	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:D:1:NAG:HN2	3:D:3:NAG:H83	1.34	0.93
1:A:61:TYR:HE2	1:A:67[B]:TRP:CH2	1.92	0.87
1:A:61:TYR:CE2	1:A:67[B]:TRP:CH2	2.65	0.85
1:A:47:GLU:CG	7:A:410:HOH:O	2.27	0.82
1:A:47:GLU:HG2	7:A:410:HOH:O	1.81	0.80
1:B:122[A]:MET:HE3	7:B:508:HOH:O	1.84	0.77
5:A:309:CIT:O4	7:A:401:HOH:O	2.11	0.68
1:A:115:ARG:HE	1:A:116:ASN:ND2	2.02	0.57
1:B:59:LYS:HB3	1:B:70[A]:ARG:HD2	1.87	0.57
1:B:2:VAL:HG21	1:B:122[A]:MET:HE3	1.88	0.55
1:A:61:TYR:CE2	1:A:67[B]:TRP:CZ3	2.99	0.50
1:A:61:TYR:CE2	1:A:67[B]:TRP:HH2	2.27	0.47
1:A:137:ASN:HD22	5:A:309:CIT:C6	2.29	0.46
1:A:61:TYR:OH	1:A:67[B]:TRP:HH2	1.99	0.46
1:B:130:ILE:HD12	1:B:210[A]:SER:OG	2.16	0.45
6:B:304[B]:TCK:HE3	6:B:304[B]:TCK:HB3	1.72	0.45
1:A:115:ARG:HH21	1:A:116:ASN:HD21	1.64	0.45
1:B:2:VAL:HG21	1:B:122[A]:MET:CE	2.47	0.44
1:B:130:ILE:HG22	1:B:212:TYR:HB3	2.00	0.44
1:B:122[A]:MET:HG3	1:B:165:TYR:CZ	2.53	0.44
1:B:157:ASN:HD21	6:B:304[B]:TCK:H1	1.83	0.43
1:B:55[A]:LEU:HD12	1:B:62:GLY:HA2	2.00	0.43
1:A:47:GLU:HG3	7:A:410:HOH:O	2.09	0.42
1:A:4:GLN:HB2	1:A:4:GLN:HE21	1.66	0.41
1:B:175:TRP:O	1:B:192:ARG:HA	2.20	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:25:ALA:HB1	1:B:55[B]:LEU:HD21	2.03	0.41
1:A:106:TYR:CE2	1:A:215:LEU:HD11	2.56	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:518:HOH:O	7:B:430:HOH:O[6_455]	2.17	0.03

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	222/219 (101%)	217 (98%)	5 (2%)	0	100	100
1	B	219/219 (100%)	216 (99%)	3 (1%)	0	100	100
All	All	441/438 (101%)	433 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	182/170 (107%)	179 (98%)	3 (2%)	62	31

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	179/170 (105%)	176 (98%)	3 (2%)	60	28
All	All	361/340 (106%)	355 (98%)	6 (2%)	59	28

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	87	TYR
1	A	172	LYS
1	A	216	GLU
1	B	4	GLN
1	B	87	TYR
1	B	112	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	4	GLN
1	A	116	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

9 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	C	1	1,2	14,14,15	0.99	1 (7%)	17,19,21	1.30	2 (11%)
2	NAG	C	2	2	14,14,15	0.81	0	17,19,21	1.01	0
2	BMA	C	3	2	11,11,12	1.59	2 (18%)	15,15,17	1.76	4 (26%)
2	XYS	C	4	2	9,9,10	1.27	2 (22%)	10,12,14	1.22	1 (10%)
2	MAN	C	5	2	11,11,12	0.78	0	15,15,17	1.81	5 (33%)
2	FUC	C	6	2	10,10,11	1.62	2 (20%)	14,14,16	2.65	4 (28%)
3	NAG	D	1	1,3	14,14,15	1.01	1 (7%)	17,19,21	2.12	6 (35%)
3	FUC	D	2	3	10,10,11	1.84	2 (20%)	14,14,16	2.52	5 (35%)
3	NAG	D	3	3	14,14,15	1.08	1 (7%)	17,19,21	1.82	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	2/6/23/26	0/1/1/1
2	BMA	C	3	2	-	0/2/19/22	0/1/1/1
2	XYS	C	4	2	1/1/3/4	-	0/1/1/1
2	MAN	C	5	2	-	2/2/19/22	0/1/1/1
2	FUC	C	6	2	-	-	0/1/1/1
3	NAG	D	1	1,3	-	2/6/23/26	0/1/1/1
3	FUC	D	2	3	-	-	0/1/1/1
3	NAG	D	3	3	-	2/6/23/26	0/1/1/1

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	2	FUC	O5-C1	-4.34	1.36	1.43
2	C	6	FUC	O5-C1	-3.76	1.37	1.43
2	C	3	BMA	O5-C1	-3.63	1.37	1.43
3	D	3	NAG	C8-C7	-3.24	1.43	1.50
3	D	1	NAG	C1-C2	2.74	1.56	1.52
2	C	3	BMA	C2-C3	2.71	1.56	1.52
2	C	1	NAG	C1-C2	2.48	1.56	1.52
2	C	6	FUC	C1-C2	-2.44	1.46	1.52
3	D	2	FUC	O5-C5	2.42	1.48	1.43
2	C	4	XYS	O5-C5	2.35	1.47	1.42
2	C	4	XYS	C1-C2	-2.23	1.47	1.52

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	6	FUC	C1-C2-C3	-8.46	99.26	109.67
3	D	1	NAG	C3-C4-C5	4.91	119.00	110.24
3	D	2	FUC	O5-C1-C2	4.87	118.29	110.77
3	D	2	FUC	C1-O5-C5	-4.78	101.95	112.78
3	D	3	NAG	C2-N2-C7	4.56	129.40	122.90
2	C	5	MAN	O5-C5-C6	4.28	113.92	107.20
3	D	2	FUC	O5-C5-C4	3.97	116.64	109.52
3	D	2	FUC	O2-C2-C1	3.57	116.46	109.15
3	D	3	NAG	O7-C7-C8	-3.36	115.82	122.06
3	D	1	NAG	O5-C1-C2	3.26	116.43	111.29
2	C	6	FUC	C1-O5-C5	-3.14	105.67	112.78
2	C	3	BMA	C1-C2-C3	-3.04	105.93	109.67
3	D	1	NAG	O3-C3-C4	-2.94	103.54	110.35
2	C	3	BMA	O6-C6-C5	-2.93	101.23	111.29
2	C	1	NAG	C4-C3-C2	-2.87	106.82	111.02
2	C	3	BMA	C1-O5-C5	2.86	116.07	112.19
3	D	1	NAG	O4-C4-C3	-2.85	103.75	110.35
2	C	3	BMA	C6-C5-C4	-2.84	106.35	113.00
3	D	2	FUC	C6-C5-C4	-2.80	107.91	113.07
3	D	1	NAG	C1-O5-C5	2.75	115.91	112.19
3	D	3	NAG	C8-C7-N2	2.54	120.40	116.10
2	C	6	FUC	O5-C1-C2	2.46	114.57	110.77
2	C	4	XYS	C5-O5-C1	2.36	115.16	111.52
2	C	5	MAN	C1-C2-C3	2.31	112.51	109.67
2	C	5	MAN	O2-C2-C3	-2.28	105.57	110.14
2	C	5	MAN	O3-C3-C2	-2.15	105.88	109.99
2	C	5	MAN	C6-C5-C4	-2.12	108.04	113.00
2	C	6	FUC	O4-C4-C3	-2.06	105.58	110.35
3	D	1	NAG	O4-C4-C5	-2.03	104.25	109.30
2	C	1	NAG	C2-N2-C7	-2.02	120.03	122.90

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	C	4	XYS	C1

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	5	MAN	O5-C5-C6-O6
3	D	1	NAG	O5-C5-C6-O6

*Continued on next page...*

*Continued from previous page...*





Mol	Chain	Res	Type	Atoms
3	D	3	NAG	C8-C7-N2-C2
3	D	3	NAG	O7-C7-N2-C2
2	C	2	NAG	C8-C7-N2-C2
2	C	2	NAG	O7-C7-N2-C2
2	C	5	MAN	C4-C5-C6-O6
3	D	1	NAG	C4-C5-C6-O6

There are no ring outliers.


2 monomers are involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	3	NAG	1	0
3	D	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

**Oligosaccharide Chain C****Bond lengths****Bond angles****Torsions****Rings**





## 5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	IPA	A	308	-	3,3,3	1.01	0	3,3,3	0.55	0
4	IPA	A	307	-	3,3,3	0.14	0	3,3,3	0.26	0
5	CIT	A	309	-	3,12,12	0.56	0	3,17,17	0.65	0
6	TCK	B	304[A]	1	20,20,21	2.70	3 (15%)	23,27,28	2.89	9 (39%)
6	TCK	B	304[B]	1	20,20,21	1.74	1 (5%)	23,27,28	2.34	8 (34%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	CIT	A	309	-	-	1/6/16/16	-
6	TCK	B	304[A]	1	-	4/19/20/22	0/1/1/1
6	TCK	B	304[B]	1	-	6/19/20/22	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	304[A]	TCK	C1-S	-10.46	1.60	1.76
6	B	304[B]	TCK	C1-S	-7.17	1.65	1.76
6	B	304[A]	TCK	O2S-S	3.75	1.47	1.43
6	B	304[A]	TCK	O1S-S	-2.93	1.40	1.43

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	304[A]	TCK	O2S-S-N	6.22	118.15	106.88
6	B	304[A]	TCK	O2S-S-O1S	-6.20	111.93	119.55
6	B	304[B]	TCK	C6-C1-S	5.54	125.79	119.77
6	B	304[A]	TCK	C2-C1-S	5.52	125.77	119.77
6	B	304[A]	TCK	C1-S-N	-5.48	100.19	107.78
6	B	304[B]	TCK	C3-C2-C1	4.99	124.61	119.45
6	B	304[B]	TCK	O-C-CM	-4.03	114.00	121.15
6	B	304[A]	TCK	O1S-S-C1	4.01	112.91	107.97
6	B	304[B]	TCK	C2-C1-S	-3.22	116.27	119.77
6	B	304[B]	TCK	O1S-S-C1	3.11	111.80	107.97
6	B	304[A]	TCK	O-C-CM	-2.86	116.08	121.15
6	B	304[A]	TCK	C3-C2-C1	2.65	122.19	119.45
6	B	304[B]	TCK	O2S-S-O1S	-2.63	116.32	119.55
6	B	304[B]	TCK	C1-S-N	-2.61	104.17	107.78
6	B	304[A]	TCK	C6-C1-S	-2.44	117.12	119.77
6	B	304[A]	TCK	C6-C1-C2	-2.42	117.07	120.44
6	B	304[B]	TCK	C6-C1-C2	-2.05	117.59	120.44

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	B	304[B]	TCK	N-CA-CB-CG

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
6	B	304[B]	TCK	C-CA-CB-CG
6	B	304[A]	TCK	CA-N-S-O2S
6	B	304[A]	TCK	CA-N-S-C1
6	B	304[B]	TCK	CE-CD-CG-CB
6	B	304[B]	TCK	CG-CD-CE-NZ
6	B	304[B]	TCK	CM-C-CA-CB
6	B	304[A]	TCK	O-C-CA-N
5	A	309	CIT	C1-C2-C3-C4
6	B	304[B]	TCK	O-C-CA-CB
6	B	304[A]	TCK	CG-CD-CE-NZ

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	309	CIT	2	0
6	B	304[B]	TCK	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	218/219 (99%)	-0.37	4 (1%) 68 69	9, 14, 30, 44	6 (2%)
1	B	218/219 (99%)	-0.10	7 (3%) 47 50	10, 15, 34, 55	6 (2%)
All	All	436/438 (99%)	-0.24	11 (2%) 60 60	9, 15, 32, 55	12 (2%)

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	171	GLY	5.1
1	B	169	SER	4.9
1	B	170	ASN	4.6
1	A	61	TYR	3.0
1	A	170	ASN	2.8
1	B	95	THR	2.7
1	A	64	LYS	2.1
1	B	90	LYS	2.0
1	A	142	TYR	2.0
1	B	99[A]	ASN	2.2
1	B	99[A]	ASN	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.


### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	MAN	C	5	11/12	0.89	0.35	38,41,52,64	0
3	NAG	D	3	14/15	0.91	0.16	26,37,54,56	0
3	FUC	D	2	10/11	0.92	0.11	21,25,31,32	0
3	NAG	D	1	14/15	0.95	0.10	17,21,25,29	0
2	BMA	C	3	11/12	0.95	0.09	17,20,24,35	0
2	XYS	C	4	9/10	0.96	0.08	18,20,29,30	0
2	NAG	C	1	14/15	0.96	0.07	14,16,22,28	0
2	NAG	C	2	14/15	0.97	0.07	13,18,27,50	0
2	FUC	C	6	10/11	0.97	0.06	17,26,34,37	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	IPA	A	308	4/4	0.61	0.17	27,36,37,37	0
5	CIT	A	309	13/13	0.85	0.16	23,33,40,40	0
4	IPA	A	307	4/4	0.91	0.09	24,28,28,29	0
6	TCK	B	304[A]	20/21	0.91	0.18	11,17,28,36	20
6	TCK	B	304[B]	20/21	0.91	0.18	14,19,35,38	20

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around TCK B 304 (A):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around TCK B 304 (B):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers ⓘ

There are no such residues in this entry.