



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 20, 2020 – 07:57 PM BST

PDB ID : 5YD6
Title : Crystal structure of PG-bound Nurr1-LBD
Authors : Sreekanth, R.; Yoon, H.S.
Deposited on : 2017-09-11
Resolution : 2.34 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2096 (2.36-2.32)
Clashscore	141614	2193 (2.36-2.32)
Ramachandran outliers	138981	2159 (2.36-2.32)
Sidechain outliers	138945	2160 (2.36-2.32)
RSRZ outliers	127900	2067 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	271	<div> <div>4%</div> <div> <div></div> <div>76%</div> <div>5%</div> <div>17%</div> </div> </div>
1	B	271	<div> <div>4%</div> <div> <div></div> <div>77%</div> <div>6%</div> <div>17%</div> </div> </div>
1	C	271	<div> <div>6%</div> <div> <div></div> <div>76%</div> <div>7%</div> <div>17%</div> </div> </div>
1	D	271	<div> <div>7%</div> <div> <div></div> <div>75%</div> <div>7%</div> <div>17%</div> </div> </div>

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 7624 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nuclear receptor subfamily 4 group A member 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	224	Total	C	N	O	S	0	0	0
			1790	1155	300	326	9			
1	B	225	Total	C	N	O	S	0	0	0
			1799	1158	303	329	9			
1	C	226	Total	C	N	O	S	0	0	0
			1805	1164	304	328	9			
1	D	226	Total	C	N	O	S	0	0	0
			1805	1164	304	328	9			

- Molecule 2 is ({Z})-7-[(1 {R},5 {S})-2-oxidanylidene-5-[({E},3 {S})-3-oxidanyloct-1-enyl]cyclopent-3-en-1-yl]hept-5-enoic acid (three-letter code: 8SU) (formula: C₂₀H₃₀O₄) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			24	20	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			24	20	4		
2	C	1	Total	C	O	0	0
			24	20	4		
2	D	1	Total	C	O	0	0
			24	20	4		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mg	0	0
			1	1		
3	A	1	Total	Mg	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	90	Total	O	0	0
			90	90		
4	B	90	Total	O	0	0
			90	90		
4	C	78	Total	O	0	0
			78	78		
4	D	69	Total	O	0	0
			69	69		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Nuclear receptor subfamily 4 group A member 2



- Molecule 1: Nuclear receptor subfamily 4 group A member 2




- Molecule 1: Nuclear receptor subfamily 4 group A member 2



- Molecule 1: Nuclear receptor subfamily 4 group A member 2





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	85.95Å 94.08Å 135.65Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.34 19.96 – 2.34	Depositor EDS
% Data completeness (in resolution range)	99.4 (20.00-2.34) 99.6 (19.96-2.34)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.76 (at 2.33Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.225 , 0.269 0.227 , 0.270	Depositor DCC
R_{free} test set	2343 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	44.9	Xtriage
Anisotropy	0.190	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 45.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7624	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 47.00 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.0571e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, 8SU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.53	0/1827	0.72	1/2472 (0.0%)
1	B	0.53	0/1835	0.73	2/2483 (0.1%)
1	C	0.54	1/1843 (0.1%)	0.71	1/2494 (0.0%)
1	D	0.62	2/1843 (0.1%)	0.77	3/2494 (0.1%)
All	All	0.56	3/7348 (0.0%)	0.73	7/9943 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	384	TYR	CD2-CE2	-9.39	1.25	1.39
1	C	495	ASN	CB-CG	6.45	1.65	1.51
1	D	384	TYR	CB-CG	-6.02	1.42	1.51

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	384	TYR	CD1-CE1-CZ	-9.89	110.90	119.80
1	D	384	TYR	CB-CG-CD1	-7.87	116.28	121.00
1	B	454	ARG	NE-CZ-NH2	-6.88	116.86	120.30
1	D	591	LEU	CA-CB-CG	5.83	128.71	115.30
1	B	454	ARG	NE-CZ-NH1	5.60	123.10	120.30
1	C	591	LEU	CA-CB-CG	5.59	128.16	115.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	454	ARG	NE-CZ-NH2	-5.59	117.51	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	384	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1790	0	1808	8	0
1	B	1799	0	1812	6	0
1	C	1805	0	1824	7	0
1	D	1805	0	1824	11	0
2	A	24	0	0	0	0
2	B	24	0	0	0	0
2	C	24	0	0	0	0
2	D	24	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	90	0	0	2	0
4	B	90	0	0	0	0
4	C	78	0	0	0	0
4	D	69	0	0	0	0
All	All	7624	0	7268	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:448:VAL:HG11	1:C:565:LEU:HD23	1.61	0.83

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:401:GLN:O	1:A:405:GLN:HG2	1.85	0.76
1:A:536:LYS:HG2	1:A:552:LEU:HD11	1.74	0.69
1:D:399:ASP:HB2	1:D:477:ARG:HH11	1.59	0.68
1:B:377:PRO:HD2	1:B:454:ARG:HD2	1.79	0.64
1:A:379:MET:HE1	4:A:715:HOH:O	2.00	0.62
1:D:591:LEU:O	1:D:595:THR:HB	2.08	0.53
1:C:440:GLU:HG2	1:C:591:LEU:HD22	1.91	0.53
1:C:379:MET:HE1	1:C:382:LEU:HD21	1.93	0.51
1:C:482:TRP:CD1	1:C:568:GLN:HB3	2.47	0.50
1:B:449:LEU:HD13	1:B:562:LEU:HD11	1.95	0.48
1:C:455:SER:HB3	1:C:461:LYS:O	2.12	0.48
1:B:426:GLY:O	1:B:429:ASP:HB2	2.12	0.48
1:D:536:LYS:HG2	1:D:552:LEU:HD11	1.95	0.47
1:A:537:ASP:HB3	1:C:541:PHE:HZ	1.81	0.46
1:B:414:MET:SD	1:B:587:ILE:HD12	2.56	0.45
1:D:456:ASN:HB3	1:D:461:LYS:HB2	1.98	0.45
1:B:411:THR:HG22	1:B:587:ILE:HD11	1.98	0.45
1:A:414:MET:SD	1:A:587:ILE:HD12	2.57	0.45
1:A:538:HIS:HD2	4:A:785:HOH:O	2.01	0.44
1:D:463:ILE:HG12	1:D:469:VAL:HG22	1.99	0.44
1:D:399:ASP:HB2	1:D:477:ARG:NH1	2.29	0.44
1:A:446:LEU:HD11	1:A:511:MET:HB2	1.99	0.44
1:A:377:PRO:HD2	1:A:454:ARG:HD2	1.99	0.43
1:D:448:VAL:HG11	1:D:565:LEU:HD13	2.01	0.42
1:B:482:TRP:CE2	1:B:486:ILE:HD11	2.55	0.42
1:D:489:PHE:CB	1:D:565:LEU:HD11	2.50	0.42
1:C:501:SER:HB2	1:C:538:HIS:CE1	2.55	0.42
1:D:360:PRO:N	1:D:361:PRO:HD3	2.35	0.41
1:D:483:ILE:HG13	1:D:483:ILE:H	1.74	0.41
1:D:449:LEU:HG	1:D:503:PHE:CE1	2.55	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	218/271 (80%)	216 (99%)	2 (1%)	0	100	100
1	B	219/271 (81%)	216 (99%)	3 (1%)	0	100	100
1	C	220/271 (81%)	219 (100%)	1 (0%)	0	100	100
1	D	220/271 (81%)	214 (97%)	6 (3%)	0	100	100
All	All	877/1084 (81%)	865 (99%)	12 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	200/242 (83%)	194 (97%)	6 (3%)	41	50
1	B	201/242 (83%)	195 (97%)	6 (3%)	41	50
1	C	202/242 (84%)	196 (97%)	6 (3%)	41	50
1	D	202/242 (84%)	196 (97%)	6 (3%)	41	50
All	All	805/968 (83%)	781 (97%)	24 (3%)	41	50

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	362	VAL
1	A	379	MET
1	A	462	LEU
1	A	491	SER
1	A	552	LEU
1	A	558	LYS
1	B	363	SER
1	B	366	SER
1	B	483	ILE
1	B	525	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	563	ARG
1	B	570	LEU
1	C	362	VAL
1	C	390	ASN
1	C	399	ASP
1	C	473	LEU
1	C	525	GLU
1	C	567	THR
1	D	390	ASN
1	D	418	ARG
1	D	458	VAL
1	D	476	VAL
1	D	483	ILE
1	D	595	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	372	HIS
1	A	435	GLN
1	A	492	ASN
1	A	494	GLN
1	A	495	ASN
1	B	372	HIS
1	B	435	GLN
1	B	492	ASN
1	B	494	GLN
1	B	495	ASN
1	B	497	ASN
1	B	538	HIS
1	C	372	HIS
1	C	494	GLN
1	C	497	ASN
1	C	538	HIS
1	D	494	GLN
1	D	538	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	8SU	D	601	1	20,24,24	0.77	1 (5%)	17,29,29	2.36	4 (23%)
2	8SU	C	601	1	20,24,24	0.68	1 (5%)	17,29,29	1.93	3 (17%)
2	8SU	B	601	1	20,24,24	0.75	1 (5%)	17,29,29	1.95	3 (17%)
2	8SU	A	601	1	20,24,24	0.78	1 (5%)	17,29,29	2.32	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	8SU	D	601	1	-	7/17/32/32	0/1/1/1
2	8SU	C	601	1	-	5/17/32/32	0/1/1/1
2	8SU	B	601	1	-	7/17/32/32	0/1/1/1
2	8SU	A	601	1	-	8/17/32/32	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	8SU	C8-C9	-2.78	1.49	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	601	8SU	C8-C9	-2.62	1.49	1.53
2	B	601	8SU	C8-C9	-2.51	1.49	1.53
2	C	601	8SU	C8-C9	-2.29	1.50	1.53

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	8SU	C11-C10-C9	-5.29	107.49	110.15
2	A	601	8SU	C15-C14-C13	-5.27	111.34	125.09
2	D	601	8SU	C11-C10-C9	-5.27	107.50	110.15
2	D	601	8SU	C8-C9-C10	4.98	112.17	107.56
2	B	601	8SU	C11-C10-C9	-4.97	107.66	110.15
2	D	601	8SU	O9-C9-C10	-4.82	123.57	127.08
2	C	601	8SU	C8-C9-C10	4.52	111.74	107.56
2	C	601	8SU	O9-C9-C10	-4.50	123.80	127.08
2	B	601	8SU	C8-C9-C10	4.07	111.32	107.56
2	A	601	8SU	O9-C9-C10	-3.96	124.20	127.08
2	C	601	8SU	C11-C10-C9	-3.77	108.26	110.15
2	A	601	8SU	C8-C9-C10	3.57	110.86	107.56
2	B	601	8SU	O9-C9-C10	-3.35	124.64	127.08
2	D	601	8SU	C15-C14-C13	-3.08	117.08	125.09

There are no chirality outliers.

All (27) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	601	8SU	C8-C12-C13-C14
2	A	601	8SU	C11-C12-C13-C14
2	A	601	8SU	C1-C2-C3-C4
2	B	601	8SU	C17-C18-C19-C20
2	C	601	8SU	C17-C18-C19-C20
2	D	601	8SU	C6-C7-C8-C12
2	B	601	8SU	C6-C7-C8-C12
2	A	601	8SU	C6-C7-C8-C12
2	A	601	8SU	C8-C12-C13-C14
2	A	601	8SU	C17-C18-C19-C20
2	C	601	8SU	C1-C2-C3-C4
2	A	601	8SU	C13-C14-C15-O15
2	D	601	8SU	C6-C7-C8-C9
2	B	601	8SU	C6-C7-C8-C9
2	A	601	8SU	C6-C7-C8-C9
2	C	601	8SU	C13-C14-C15-C16

Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	B	601	8SU	C13-C14-C15-C16
2	D	601	8SU	C11-C12-C13-C14
2	D	601	8SU	C2-C3-C4-C5
2	C	601	8SU	C13-C14-C15-O15
2	B	601	8SU	C13-C14-C15-O15
2	B	601	8SU	C16-C17-C18-C19
2	D	601	8SU	C14-C15-C16-C17
2	D	601	8SU	O15-C15-C16-C17
2	B	601	8SU	C2-C3-C4-C5
2	C	601	8SU	C3-C4-C5-C6
2	A	601	8SU	C13-C14-C15-C16


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	224/271 (82%)	0.16	10 (4%)	33 44	30, 47, 75, 101	0
1	B	225/271 (83%)	0.26	11 (4%)	29 40	33, 50, 75, 101	0
1	C	226/271 (83%)	0.38	15 (6%)	18 26	35, 55, 93, 121	0
1	D	226/271 (83%)	0.45	18 (7%)	12 18	34, 54, 91, 122	0
All	All	901/1084 (83%)	0.31	54 (5%)	21 30	30, 51, 85, 122	0

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	541	PHE	6.5
1	A	362	VAL	5.7
1	B	362	VAL	4.9
1	C	541	PHE	4.5
1	B	392	ASP	4.5
1	D	385	SER	4.1
1	C	361	PRO	4.1
1	D	516	HIS	4.0
1	C	458	VAL	3.6
1	D	389	ALA	3.5
1	C	540	THR	3.4
1	D	401	GLN	3.4
1	D	361	PRO	3.2
1	A	539	VAL	3.2
1	D	541	PHE	3.1
1	D	512	VAL	3.1
1	B	390	ASN	3.1
1	A	550	ASN	3.0
1	A	551	TYR	2.9
1	B	551	TYR	2.9
1	C	550	ASN	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	385	SER	2.8
1	A	458	VAL	2.8
1	D	477	ARG	2.8
1	C	462	LEU	2.8
1	D	381	SER	2.8
1	B	401	GLN	2.8
1	D	462	LEU	2.6
1	B	458	VAL	2.6
1	C	573	ILE	2.6
1	D	542	ASN	2.5
1	A	401	GLN	2.5
1	B	540	THR	2.5
1	D	458	VAL	2.4
1	C	481	GLU	2.4
1	D	551	TYR	2.3
1	B	543	ASN	2.3
1	C	543	ASN	2.3
1	A	516	HIS	2.3
1	C	549	PRO	2.3
1	D	501	SER	2.3
1	C	380	THR	2.2
1	D	473	LEU	2.2
1	C	517	GLY	2.2
1	A	392	ASP	2.2
1	A	451	LEU	2.1
1	B	522	LYS	2.1
1	D	564	THR	2.1
1	D	390	ASN	2.1
1	D	497	ASN	2.1
1	C	516	HIS	2.1
1	B	477	ARG	2.1
1	A	477	ARG	2.0
1	C	477	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no monosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	8SU	C	601	24/24	0.69	0.31	47,64,80,80	0
2	8SU	B	601	24/24	0.77	0.26	47,64,83,85	0
2	8SU	A	601	24/24	0.80	0.22	47,58,84,84	0
3	MG	A	602	1/1	0.81	0.31	58,58,58,58	0
2	8SU	D	601	24/24	0.83	0.24	33,55,75,76	0
3	MG	B	602	1/1	0.94	0.38	63,63,63,63	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 8SU C 601:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 8SU B 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 8SU A 601:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 8SU D 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.