



# Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 07:01 am BST

PDB ID : 1YFE  
Title : Crystal structure of apo fumarase C from Escherichia coli  
Authors : Weaver, T.  
Deposited on : 2004-12-31  
Resolution : 2.19 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.19 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	467	<div> <div>4%</div> <div> <div></div> <div>76%</div> <div>20%</div> <div>••</div> </div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3575 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fumarate hydratase class II.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	457	Total	C	N	O	S	0	0	0
			3463	2163	618	659	23			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	112	Total	O	0	0
			112	112		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Fumarate hydratase class II



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	121.57Å 127.96Å 62.12Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 2.19 22.05 – 2.07	Depositor EDS
% Data completeness (in resolution range)	92.3 (8.00-2.19) 85.6 (22.05-2.07)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	0.05	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.76 (at 2.08Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.203 , 0.210 0.196 , 0.185	Depositor DCC
$R_{free}$ test set	2590 reflections (9.37%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	21.8	Xtriage
Anisotropy	0.408	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 75.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.017 for -k,-h,-l	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3575	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.36% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.33	0/3521	0.59	0/4770

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3463	0	3482	68	0
2	A	112	0	0	0	0
All	All	3575	0	3482	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:227:LEU:HD12	1:A:246:VAL:HG11	1.45	0.98
1:A:112:ASN:HD21	1:A:126:ARG:HH21	1.15	0.94
1:A:343:ASN:HD21	1:A:371:GLN:HE21	1.26	0.83
1:A:436:LYS:HD2	1:A:447:GLU:HG3	1.66	0.77

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:112:ASN:ND2	1:A:126:ARG:HH21	1.84	0.76
1:A:397:ARG:HH21	1:A:401:LEU:HD21	1.52	0.75
1:A:116:GLU:HA	1:A:120:GLY:HA2	1.71	0.71
1:A:119:GLY:HA3	1:A:127:LYS:HE2	1.73	0.70
1:A:91:LEU:HD13	1:A:101:GLN:HB3	1.76	0.67
1:A:17:PRO:HB2	1:A:20:LYS:HG2	1.77	0.66
1:A:366:ILE:HG13	1:A:370:LEU:HD22	1.78	0.65
1:A:31:GLU:HG3	1:A:34:ARG:NH1	2.11	0.65
1:A:384:ASN:HA	1:A:388:ALA:HB3	1.80	0.62
1:A:413:ASN:N	1:A:413:ASN:HD22	2.00	0.60
1:A:41:PRO:HG3	1:A:370:LEU:HD23	1.83	0.60
1:A:413:ASN:HA	1:A:416:ILE:O	2.02	0.59
1:A:112:ASN:ND2	1:A:126:ARG:HD3	2.21	0.56
1:A:112:ASN:HD21	1:A:126:ARG:NH2	1.96	0.55
1:A:154:LEU:O	1:A:158:LYS:HG2	2.06	0.55
1:A:428:ALA:HB1	1:A:433:LEU:O	2.06	0.55
1:A:129:HIS:HB3	1:A:132:ASP:HB2	1.89	0.54
1:A:278:HIS:HE1	1:A:372:SER:OG	1.89	0.54
1:A:146:THR:O	1:A:150:VAL:HG23	2.08	0.54
1:A:5:ARG:HH11	1:A:122:ARG:HG2	1.74	0.53
1:A:436:LYS:HB2	1:A:450:PHE:CE2	2.44	0.53
1:A:155:ALA:O	1:A:159:GLN:HB3	2.09	0.52
1:A:132:ASP:O	1:A:136:LYS:HE3	2.09	0.52
1:A:41:PRO:CG	1:A:370:LEU:HD23	2.40	0.52
1:A:450:PHE:O	1:A:454:VAL:HG22	2.10	0.51
1:A:188:HIS:O	1:A:189:LEU:HB2	2.10	0.51
1:A:293:ASN:O	1:A:296:ARG:HG3	2.11	0.51
1:A:294:ASP:O	1:A:298:LEU:HB2	2.11	0.51
1:A:107:ASN:HA	1:A:134:VAL:HG11	1.93	0.50
1:A:397:ARG:NH2	1:A:401:LEU:HD21	2.25	0.50
1:A:278:HIS:HD2	1:A:344:ASP:OD1	1.93	0.50
1:A:337:CYS:O	1:A:341:MET:HG2	2.11	0.50
1:A:5:ARG:NH1	1:A:122:ARG:HG2	2.25	0.50
1:A:343:ASN:HD21	1:A:371:GLN:NE2	2.03	0.50
1:A:178:PHE:CZ	1:A:393:PRO:HD3	2.47	0.50
1:A:121:VAL:HB	1:A:126:ARG:HA	1.93	0.49
1:A:454:VAL:HG23	1:A:454:VAL:O	2.11	0.49
1:A:436:LYS:HD3	1:A:450:PHE:CD2	2.47	0.49
1:A:413:ASN:H	1:A:413:ASN:HD22	1.59	0.49
1:A:29:SER:HB3	1:A:101:GLN:HG2	1.95	0.48
1:A:320:ILE:C	1:A:322:PRO:HD3	2.34	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:31:GLU:HG3	1:A:34:ARG:HH11	1.79	0.46
1:A:17:PRO:HB2	1:A:20:LYS:CG	2.44	0.45
1:A:5:ARG:HH21	1:A:16:VAL:HG12	1.82	0.45
1:A:119:GLY:HA3	1:A:127:LYS:CE	2.44	0.45
1:A:306:ILE:HG22	1:A:308:GLU:HG3	2.00	0.44
1:A:431:GLU:HB3	1:A:433:LEU:HD13	1.98	0.44
1:A:248:ASP:O	1:A:252:VAL:HG23	2.18	0.44
1:A:376:LEU:O	1:A:380:MET:HG3	2.18	0.44
1:A:154:LEU:HD21	1:A:254:THR:HB	2.00	0.43
1:A:106:MET:O	1:A:110:LEU:HB2	2.18	0.42
1:A:130:PRO:O	1:A:134:VAL:HB	2.18	0.42
1:A:450:PHE:CE2	1:A:454:VAL:HG21	2.54	0.42
1:A:331:GLU:O	1:A:335:MET:HG3	2.19	0.42
1:A:385:LYS:HD2	1:A:386:HIS:CE1	2.54	0.42
1:A:144:PHE:HB3	1:A:145:PRO:HD3	2.01	0.42
1:A:232:VAL:HG23	1:A:234:THR:HG23	2.01	0.42
1:A:428:ALA:HA	1:A:433:LEU:HB2	2.02	0.42
1:A:89:PHE:HA	1:A:105:ASN:HD21	1.85	0.41
1:A:58:VAL:HG21	1:A:246:VAL:HA	2.02	0.41
1:A:278:HIS:HE1	1:A:372:SER:CB	2.34	0.41
1:A:146:THR:OG1	1:A:227:LEU:HB2	2.21	0.41
1:A:153:LEU:HG	1:A:157:ARG:HD3	2.03	0.41
1:A:446:SER:OG	1:A:449:GLU:HG3	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	455/467 (97%)	437 (96%)	17 (4%)	1 (0%)	47 55

All (1) Ramachandran outliers are listed below:



Mol	Chain	Res	Type
1	A	417	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	369/376 (98%)	348 (94%)	21 (6%)	20	24

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	101	GLN
1	A	110	LEU
1	A	121	VAL
1	A	149	HIS
1	A	156	LEU
1	A	167	LEU
1	A	197	LEU
1	A	213	LYS
1	A	221	HIS
1	A	227	LEU
1	A	282	LYS
1	A	289	MET
1	A	296	ARG
1	A	298	LEU
1	A	370	LEU
1	A	376	LEU
1	A	402	LEU
1	A	406	LEU
1	A	413	ASN
1	A	419	ASP
1	A	423	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	32	HIS
1	A	101	GLN
1	A	105	ASN
1	A	112	ASN
1	A	149	HIS
1	A	163	GLN
1	A	169	GLN
1	A	172	ASN
1	A	239	HIS
1	A	278	HIS
1	A	326	ASN
1	A	339	GLN
1	A	368	ASN
1	A	371	GLN
1	A	399	ASN
1	A	413	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	457/467 (97%)	-0.03	21 (4%) 32 31	7, 26, 73, 93	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	414	THR	4.1
1	A	415	HIS	3.6
1	A	319	SER	3.4
1	A	418	TYR	3.3
1	A	444	TYR	3.3
1	A	453	TRP	3.2
1	A	431	GLU	3.1
1	A	121	VAL	3.1
1	A	433	LEU	3.0
1	A	237	ASN	3.0
1	A	317	GLY	2.8
1	A	412	LEU	2.7
1	A	435	LEU	2.7
1	A	455	ARG	2.7
1	A	419	ASP	2.6
1	A	439	ALA	2.4
1	A	460	VAL	2.3
1	A	447	GLU	2.3
1	A	320	ILE	2.3
1	A	440	LEU	2.2
1	A	458	GLN	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.