



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 8, 2020 – 08:17 PM BST

PDB ID : 2YG1  
Title : APO STRUCTURE OF CELLOBIOHYDROLASE 1 (CEL7A) FROM HETEROBASIDION ANNOSUM  
Authors : Haddad-Momeni, M.; Hansson, H.; Mikkelsen, N.E.; Wang, X.; Svedberg, J.; Sandgren, M.; Stahlberg, J.  
Deposited on : 2011-04-11  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.13.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	440	<div> <div>2%</div> <div> <div></div> <div>91%</div> <div>8%</div> <div></div> </div> </div>
1	B	440	<div> <div>6%</div> <div> <div></div> <div>88%</div> <div>9%</div> <div></div> </div> </div>
2	C	2	<div> <div></div> <div> <div>50%</div> <div>50%</div> </div> </div>
2	D	2	<div> <div></div> <div>100%</div> </div>

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7481 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CELLULOSE 1,4-BETA-CELLOBIOSIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	440	Total	C	N	O	S	0	3	0
			3298	2020	559	692	27			
1	B	426	Total	C	N	O	S	0	1	0
			3201	1968	543	663	27			

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	D	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total	Mg	0	0
			2	2		
3	A	1	Total	Mg	0	0
			1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	531	Total	O	0	0
			531	531		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	392	Total	O	0	0
			392	392		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: CELLULOSE 1,4-BETA-CELLOBIOSIDASE




#### • Molecule 1: CELLULOSE 1,4-BETA-CELLOBIOSIDASE



#### • Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



#### • Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	60.51Å 84.10Å 75.88Å 90.00° 103.36° 90.00°	Depositor
Resolution (Å)	28.10 – 1.90 28.10 – 1.90	Depositor EDS
% Data completeness (in resolution range)	98.1 (28.10-1.90) 98.1 (28.10-1.90)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.39 (at 1.91Å)	Xtriage
Refinement program	REFMAC 5.5.0110	Depositor
R, $R_{free}$	0.197 , 0.253 0.198 , 0.256	Depositor DCC
$R_{free}$ test set	2859 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	17.1	Xtriage
Anisotropy	0.064	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 48.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7481	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 40.89 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.5574e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, PCA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.41	0/3365	0.57	0/4583
1	B	0.38	0/3259	0.53	0/4433
All	All	0.40	0/6624	0.55	0/9016

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3298	0	3056	36	0
1	B	3201	0	2969	25	0
2	C	28	0	25	0	0
2	D	28	0	25	0	0
3	A	1	0	0	0	0
3	B	2	0	0	0	0
4	A	531	0	0	2	0
4	B	392	0	0	0	0
All	All	7481	0	6075	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:258:LYS:HE2	4:A:2259:HOH:O	1.63	0.96
1:A:322[B]:THR:HG22	1:A:330:PHE:H	1.33	0.92
1:A:215:MET:SD	1:A:360:MET:CE	2.62	0.87
1:A:215:MET:HB2	1:A:360:MET:HE3	1.55	0.87
1:A:45:SER:HA	1:A:46:GLY:C	1.98	0.83
1:B:15:THR:HG23	1:B:27:THR:HG23	1.76	0.68
1:A:40:TRP:HE1	1:A:49:ASN:ND2	1.91	0.67
1:B:15:THR:HG23	1:B:27:THR:CG2	2.25	0.66
1:A:139:PRO:HD2	1:A:142:LEU:HD12	1.78	0.66
1:A:40:TRP:HE1	1:A:49:ASN:HD21	1.42	0.65
1:A:215:MET:CB	1:A:360:MET:HE3	2.28	0.64
1:A:125:LYS:HE2	1:A:365:ASP:OD1	1.99	0.62
1:B:246:GLY:O	1:B:258:LYS:HE3	1.99	0.62
1:A:45:SER:HA	1:A:47:THR:N	2.15	0.61
1:A:358:ASN:ND2	1:A:362:GLN:HE21	1.98	0.61
1:A:215:MET:SD	1:A:360:MET:HE2	2.42	0.60
1:B:143:ASN:HB2	1:B:382:MET:HE2	1.84	0.59
1:A:296:ASP:OD1	1:A:298:THR:HG22	2.07	0.55
1:B:51:TYR:HD1	1:B:56:TRP:HA	1.73	0.54
1:A:322[B]:THR:CG2	1:A:330:PHE:H	2.14	0.54
1:B:126:ASN:HB3	1:B:299:THR:HG22	1.89	0.54
1:A:215:MET:HA	1:A:360:MET:HE1	1.89	0.53
1:B:143:ASN:HD22	1:B:382:MET:HE1	1.74	0.52
1:A:229:PRO:HB3	1:A:360:MET:HE2	1.93	0.51
1:A:215:MET:SD	1:A:360:MET:HE1	2.50	0.50
1:A:17:SER:HB2	1:A:430:LYS:HG2	1.93	0.50
1:B:136:SER:HA	1:B:222:ILE:HD12	1.93	0.50
1:A:215:MET:SD	1:A:360:MET:HE3	2.51	0.49
1:A:298:THR:HG23	1:A:300:THR:H	1.78	0.49
1:A:358:ASN:HD22	1:A:362:GLN:HE21	1.59	0.49
1:A:229:PRO:HA	1:A:360:MET:HE1	1.95	0.48
1:B:220:ALA:HA	1:B:385:LEU:HD21	1.96	0.48
1:B:386:ASP:O	1:B:405:CYS:HB2	2.14	0.47
1:A:244:GLN:HE22	1:B:244:GLN:NE2	2.14	0.46
1:B:15:THR:CG2	1:B:27:THR:HG23	2.43	0.46
1:B:287:THR:HB	1:B:311:VAL:HB	1.98	0.45
1:B:51:TYR:CD1	1:B:56:TRP:HA	2.52	0.45
1:A:44:THR:O	1:A:45:SER:CB	2.65	0.44
1:B:175:ASP:HB2	1:B:214:GLU:OE1	2.18	0.44
1:B:15:THR:CG2	1:B:27:THR:CG2	2.95	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:11:HIS:HB3	1:A:32:VAL:O	2.18	0.44
1:A:257:ASP:OD2	1:A:260:GLY:HA2	2.18	0.43
1:A:35:ASP:OD2	1:A:37:ASN:HB2	2.18	0.43
1:B:308:ARG:O	1:B:319:ASN:HB2	2.19	0.43
1:B:419:ASN:N	1:B:420:PRO:HD3	2.34	0.42
1:B:161:PRO:O	1:B:164:LYS:HE3	2.18	0.42
1:B:176:THR:HG21	1:B:230:HIS:HB2	2.02	0.42
1:A:44:THR:O	1:A:45:SER:HB2	2.20	0.42
1:B:141:GLY:HA2	1:B:382:MET:HB2	2.01	0.42
1:B:179:PRO:HD2	1:B:210:SER:O	2.20	0.42
1:A:194:TRP:CZ2	1:A:204:GLY:HA3	2.55	0.41
1:A:329:GLN:HB2	1:A:329:GLN:HE21	1.58	0.41
1:A:45:SER:OG	1:A:47:THR:HB	2.21	0.41
1:A:47:THR:N	4:A:2087:HOH:O	2.53	0.41
1:A:42:HIS:NE2	1:A:74:ASP:OD2	2.54	0.40
1:B:176:THR:HG21	1:B:230:HIS:CB	2.51	0.40
1:A:136:SER:HA	1:A:222:ILE:HD12	2.04	0.40
1:A:215:MET:HB2	1:A:360:MET:CE	2.39	0.40
1:B:175:ASP:OD1	1:B:179:PRO:HD3	2.20	0.40
1:B:211:CYS:HB3	1:B:256:CYS:SG	2.61	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	441/440 (100%)	430 (98%)	9 (2%)	2 (0%)	29	18
1	B	419/440 (95%)	407 (97%)	11 (3%)	1 (0%)	47	38
All	All	860/880 (98%)	837 (97%)	20 (2%)	3 (0%)	34	31

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	45	SER
1	B	248	GLY
1	A	248	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	369/366 (101%)	360 (98%)	9 (2%)	49	43
1	B	356/366 (97%)	352 (99%)	4 (1%)	73	73
All	All	725/732 (99%)	712 (98%)	13 (2%)	59	55

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	37	ASN
1	A	47	THR
1	A	49	ASN
1	A	74	ASP
1	A	202	ASN
1	A	257	ASP
1	A	329	GLN
1	A	341	GLN
1	A	358	ASN
1	B	138	LEU
1	B	257	ASP
1	B	341	GLN
1	B	383	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	18	GLN
1	A	49	ASN
1	A	104	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	162	ASN
1	A	244	GLN
1	A	329	GLN
1	A	358	ASN
1	A	414	ASN
1	A	421	ASN
1	B	6	GLN
1	B	104	ASN
1	B	237	GLN
1	B	304	HIS
1	B	381	ASN
1	B	418	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	PCA	A	1	1	7,8,9	1.65	1 (14%)	9,10,12	2.05	5 (55%)
1	PCA	B	1	1	7,8,9	1.76	1 (14%)	9,10,12	2.08	5 (55%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PCA	A	1	1	-	0/0/11/13	0/1/1/1
1	PCA	B	1	1	-	0/0/11/13	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	1	PCA	CD-N	4.41	1.46	1.34
1	A	1	PCA	CD-N	4.21	1.45	1.34

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1	PCA	OE-CD-CG	-2.91	121.68	126.76
1	B	1	PCA	CA-N-CD	-2.88	103.72	113.58
1	A	1	PCA	CA-N-CD	-2.81	103.97	113.58
1	A	1	PCA	CB-CA-C	-2.78	108.87	112.70
1	A	1	PCA	CB-CA-N	2.62	110.81	103.30
1	B	1	PCA	CB-CA-C	-2.59	109.14	112.70
1	B	1	PCA	CB-CA-N	2.42	110.25	103.30
1	A	1	PCA	CG-CD-N	2.37	114.51	108.39
1	B	1	PCA	CG-CD-N	2.33	114.43	108.39
1	A	1	PCA	OE-CD-CG	-2.13	123.05	126.76

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAG	C	1	1,2	14,14,15	0.54	0	17,19,21	0.72	0
2	NAG	C	2	2	14,14,15	0.63	0	17,19,21	1.44	2 (11%)
2	NAG	D	1	1,2	14,14,15	0.53	0	17,19,21	0.99	1 (5%)
2	NAG	D	2	2	14,14,15	0.45	0	17,19,21	1.11	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	2/6/23/26	0/1/1/1
2	NAG	D	1	1,2	-	1/6/23/26	0/1/1/1
2	NAG	D	2	2	-	3/6/23/26	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	2	NAG	C4-C3-C2	4.22	117.21	111.02
2	C	2	NAG	C3-C4-C5	2.47	114.64	110.24
2	D	2	NAG	O5-C1-C2	-2.19	107.83	111.29
2	D	1	NAG	O5-C1-C2	-2.17	107.86	111.29

There are no chirality outliers.


All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	2	NAG	C4-C5-C6-O6
2	D	2	NAG	C8-C7-N2-C2
2	D	2	NAG	O7-C7-N2-C2
2	C	2	NAG	O5-C5-C6-O6
2	D	1	NAG	C4-C5-C6-O6
2	D	2	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	439/440 (99%)	-0.18	10 (2%) 60 63	7, 14, 26, 33	0
1	B	425/440 (96%)	0.34	25 (5%) 22 25	11, 24, 35, 41	0
All	All	864/880 (98%)	0.08	35 (4%) 37 40	7, 18, 32, 41	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	391	THR	4.5
1	A	46	GLY	3.9
1	A	198	SER	3.7
1	B	300	THR	3.5
1	A	300	THR	3.5
1	B	440	ALA	3.4
1	B	314	GLY	3.3
1	B	299	THR	3.2
1	B	248	GLY	3.2
1	B	51	TYR	3.0
1	B	315	LYS	3.0
1	A	248	GLY	2.8
1	A	249	ASP	2.8
1	B	249	ASP	2.7
1	B	408	THR	2.7
1	B	196	PRO	2.6
1	B	439	THR	2.6
1	B	195	THR	2.5
1	A	298	THR	2.5
1	B	392	THR	2.5
1	A	101	TYR	2.4
1	A	440	ALA	2.4
1	B	378	HIS	2.3
1	B	410	GLY	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	217	VAL	2.2
1	A	45	SER	2.2
1	A	299	THR	2.2
1	B	49	ASN	2.2
1	B	98[A]	ASN	2.1
1	B	379	GLU	2.1
1	B	56	TRP	2.1
1	B	44	THR	2.1
1	B	194	TRP	2.0
1	B	282	THR	2.0
1	B	389	TYR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	PCA	A	1	8/9	0.95	0.09	10,10,11,11	0
1	PCA	B	1	8/9	0.95	0.10	17,17,17,18	0

## 6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	C	2	14/15	0.79	0.29	35,37,38,39	0
2	NAG	D	2	14/15	0.82	0.30	39,41,42,42	0
2	NAG	D	1	14/15	0.89	0.12	31,33,35,37	0
2	NAG	C	1	14/15	0.90	0.10	25,28,30,32	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around Chain C:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	MG	B	444	1/1	0.98	0.14	21,21,21,21	0
3	MG	A	443	1/1	0.99	0.05	12,12,12,12	0
3	MG	B	443	1/1	0.99	0.04	16,16,16,16	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.