



Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 01:00 am BST

PDB ID : 5YHP
Title : Proline iminopeptidase from Psychrophilic yeast glaciozyma antarctica
Authors : Rodzli, N.A.; Kamaruddin, S.; Jonet, A.; Seman, W.M.K.W.; Tab, M.M.; Minor, N.; Jaafar, N.R.; Mahadi, N.M.; Murad, A.M.A.; Bakar, F.D.A.; Illias, R.M.D.
Deposited on : 2017-09-29
Resolution : 2.39 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	355	<div> <div>6%</div> <div> <div></div> <div>81%</div> <div>9%</div> <div>10%</div> </div> </div>
1	B	355	<div> <div>5%</div> <div> <div></div> <div>80%</div> <div>9%</div> <div>10%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	FLC	B	401	-	-	-	X

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 5177 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cold active proline iminopeptidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	318	Total	C	N	O	S	0	0	0
			2495	1601	425	463	6			
1	B	319	Total	C	N	O	S	0	0	0
			2503	1606	426	464	7			

- Molecule 2 is CITRATE ANION (three-letter code: FLC) (formula: $C_6H_5O_7$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			13	6	7		
2	A	1	Total	C	O	0	0
			13	6	7		
2	B	1	Total	C	O	0	0
			13	6	7		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	72	Total 72	O 72	0	0
3	B	68	Total 68	O 68	0	0

- Molecule 1: Cold active proline iminopeptidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2 ₁	Depositor
Cell constants a, b, c, α , β , γ	55.39 Å 82.27 Å 171.00 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	27.69 – 2.39 28.57 – 2.39	Depositor EDS
% Data completeness (in resolution range)	90.8 (27.69-2.39) 84.2 (28.57-2.39)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.61 (at 2.39 Å)	Xtriage
Refinement program	PHENIX 1.8.2_1309	Depositor
R, R_{free}	0.186 , 0.215 0.186 , 0.215	Depositor DCC
R_{free} test set	2000 reflections (6.95%)	wwPDB-VP
Wilson B-factor (Å ²)	58.8	Xtriage
Anisotropy	0.183	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 27.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5177	wwPDB-VP
Average B, all atoms (Å ²)	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.42% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.43	0/2565	0.73	6/3484 (0.2%)
1	B	0.44	0/2573	0.65	5/3494 (0.1%)
All	All	0.43	0/5138	0.69	11/6978 (0.2%)

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	23	ARG	NE-CZ-NH2	13.60	127.10	120.30
1	A	23	ARG	NE-CZ-NH1	-13.42	113.59	120.30
1	A	127	ARG	NE-CZ-NH2	13.04	126.82	120.30
1	A	127	ARG	NE-CZ-NH1	-11.37	114.62	120.30
1	B	127	ARG	NE-CZ-NH2	-10.33	115.13	120.30
1	B	127	ARG	NE-CZ-NH1	10.08	125.34	120.30
1	B	23	ARG	NE-CZ-NH2	-9.44	115.58	120.30
1	B	23	ARG	NE-CZ-NH1	8.85	124.72	120.30
1	A	127	ARG	CD-NE-CZ	6.02	132.03	123.60
1	A	23	ARG	CD-NE-CZ	5.86	131.81	123.60
1	B	127	ARG	CD-NE-CZ	5.41	131.18	123.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2495	0	2433	17	0
1	B	2503	0	2445	17	1
2	A	26	0	10	2	1
2	B	13	0	5	3	0
3	A	72	0	0	0	0
3	B	68	0	0	2	0
All	All	5177	0	4893	33	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:42:GLY:HA3	2:B:401:FLC:HG2	1.60	0.83
1:B:170:VAL:HA	1:B:173:ARG:HD3	1.62	0.80
2:B:401:FLC:OA1	2:B:401:FLC:HG1	1.92	0.69
1:A:170:VAL:HA	1:A:173:ARG:HD3	1.80	0.62
1:A:297:HIS:CD2	2:A:402:FLC:HG2	2.37	0.60
1:B:134:ARG:NH2	2:B:401:FLC:OG1	2.29	0.55
1:B:60:THR:HG21	1:B:316:ALA:HB1	1.88	0.54
1:A:275:THR:HA	1:B:158:PRO:HG3	1.92	0.52
1:B:35:LYS:NZ	1:B:318:LEU:O	2.39	0.51
1:A:60:THR:HG21	1:A:316:ALA:HB1	1.92	0.51
1:B:90:GLU:OE1	3:B:501:HOH:O	2.20	0.50
1:A:141:LYS:NZ	1:A:145:ASP:OD2	2.45	0.49
1:A:281:THR:HG21	1:A:289:PHE:CG	2.47	0.49
1:B:141:LYS:NZ	1:B:145:ASP:OD2	2.45	0.48
1:A:103:ALA:O	1:A:127:ARG:HD2	2.13	0.48
1:B:265:GLN:HG2	1:B:274:ILE:HG13	1.96	0.47
1:B:281:THR:HG21	1:B:289:PHE:CG	2.49	0.47
1:A:265:GLN:HG2	1:A:274:ILE:HG13	1.96	0.46
1:A:134:ARG:HH12	2:A:402:FLC:HG1	1.82	0.43
1:B:298:SER:OG	1:B:300:ILE:HG12	2.18	0.43
1:B:115:THR:HG21	1:B:139:LEU:HD13	2.01	0.43
1:B:173:ARG:HG3	1:B:176:MET:SD	2.58	0.43
1:B:21:LEU:HA	1:B:21:LEU:HD12	1.84	0.43
1:A:315:PHE:O	1:A:318:LEU:HB2	2.19	0.43
1:A:45:GLY:HA2	1:A:71:SER:HB3	2.00	0.43
1:A:298:SER:OG	1:A:300:ILE:HG12	2.19	0.42
1:A:282:LYS:NZ	1:B:162:GLU:OE2	2.31	0.42
1:A:173:ARG:HG3	1:A:176:MET:SD	2.60	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:15:LEU:HD12	1:B:15:LEU:HA	1.79	0.41
1:A:115:THR:HG21	1:A:139:LEU:HD13	2.02	0.41
1:A:21:LEU:HA	1:A:21:LEU:HD12	1.89	0.41
1:B:245:GLY:HA2	3:B:555:HOH:O	2.19	0.41
1:A:35:LYS:NZ	1:A:318:LEU:O	2.48	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:23:ARG:NH1	2:A:401:FLC:OG2[1_545]	2.10	0.10

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	316/355 (89%)	311 (98%)	5 (2%)	0	100	100
1	B	317/355 (89%)	311 (98%)	6 (2%)	0	100	100
All	All	633/710 (89%)	622 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	257/288 (89%)	252 (98%)	5 (2%)	57	75
1	B	258/288 (90%)	249 (96%)	9 (4%)	36	55
All	All	515/576 (89%)	501 (97%)	14 (3%)	44	65

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	SER
1	A	105	TRP
1	A	127	ARG
1	A	191	VAL
1	A	215	TYR
1	B	3	SER
1	B	4	LEU
1	B	17	ARG
1	B	23	ARG
1	B	101	VAL
1	B	105	TRP
1	B	127	ARG
1	B	215	TYR
1	B	247	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FLC	A	401	-	3,12,12	1.49	0	3,17,17	2.17	1 (33%)
2	FLC	B	401	-	3,12,12	1.98	1 (33%)	3,17,17	2.90	1 (33%)
2	FLC	A	402	-	3,12,12	1.74	1 (33%)	3,17,17	2.19	2 (66%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FLC	A	401	-	-	0/6/16/16	-
2	FLC	B	401	-	-	3/6/16/16	-
2	FLC	A	402	-	-	3/6/16/16	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	FLC	CA-CB	-2.87	1.50	1.54
2	A	402	FLC	CG-CB	-2.15	1.51	1.54

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	FLC	CB-CA-CAC	-4.54	107.71	114.98
2	A	401	FLC	CB-CG-CGC	-3.07	110.06	114.98
2	A	402	FLC	CB-CA-CAC	-2.94	110.27	114.98
2	A	402	FLC	CB-CG-CGC	-2.39	111.15	114.98

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	401	FLC	CBC-CB-CG-CGC
2	B	401	FLC	OHB-CB-CG-CGC
2	A	402	FLC	CAC-CA-CB-CBC
2	A	402	FLC	CAC-CA-CB-OHB
2	B	401	FLC	CA-CB-CG-CGC
2	A	402	FLC	CAC-CA-CB-CG

There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	FLC	0	1
2	B	401	FLC	3	0
2	A	402	FLC	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	318/355 (89%)	0.02	23 (7%) 15 14	50, 61, 81, 102	0
1	B	319/355 (89%)	-0.01	17 (5%) 26 25	49, 60, 76, 108	0
All	All	637/710 (89%)	0.00	40 (6%) 20 18	49, 61, 78, 108	0

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	40	LEU	5.5
1	A	110	GLY	4.6
1	A	39	PHE	4.4
1	B	111	SER	4.1
1	A	111	SER	3.8
1	A	113	GLY	3.6
1	A	112	TRP	3.5
1	A	114	SER	3.5
1	A	137	PHE	3.5
1	B	40	LEU	3.3
1	A	23	ARG	3.3
1	B	43	GLY	3.3
1	B	116	LEU	3.3
1	B	117	ALA	3.3
1	B	114	SER	3.2
1	B	112	TRP	3.2
1	B	115	THR	3.0
1	A	135	GLY	3.0
1	A	43	GLY	2.9
1	A	41	HIS	2.9
1	B	109	GLY	2.8
1	B	319	ALA	2.7
1	A	3	SER	2.7
1	A	42	GLY	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	42	GLY	2.7
1	A	271	VAL	2.6
1	B	137	PHE	2.5
1	A	116	LEU	2.5
1	A	107	VAL	2.5
1	A	272	CYS	2.5
1	B	272	CYS	2.5
1	B	110	GLY	2.4
1	A	109	GLY	2.4
1	B	136	ILE	2.3
1	B	38	VAL	2.2
1	A	117	ALA	2.2
1	A	136	ILE	2.2
1	A	44	PRO	2.2
1	B	133	LEU	2.0
1	A	115	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	FLC	B	401	13/13	0.72	0.41	90,90,90,90	0
2	FLC	A	402	13/13	0.85	0.39	98,98,98,98	0
2	FLC	A	401	13/13	0.94	0.16	70,70,70,70	0

6.5 Other polymers [i](#)

There are no such residues in this entry.