



Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 06:23 am BST

PDB ID : 4YOT
Title : Crystal structure of a trimeric exonuclease PhoExo I from *Pyrococcus horikoshii* OT3 at 2.15Å resolution
Authors : Miyazono, K.; Tsutsumi, K.; Tanokura, M.
Deposited on : 2015-03-12
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION




The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1479 (2.16-2.16)
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)
RSRZ outliers	127900	1456 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	229	
1	B	229	
1	C	229	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5484 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-5 exonuclease PhoExo I.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	225	Total	C	N	O	S	0	0	0
			1782	1151	296	332	3			
1	B	220	Total	C	N	O	S	0	0	0
			1738	1124	287	324	3			
1	C	218	Total	C	N	O	S	0	0	0
			1725	1116	284	322	3			

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mg	0	0
			1	1		
2	A	1	Total	Mg	0	0
			1	1		
2	C	1	Total	Mg	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	102	Total	O	0	0
			102	102		
3	B	80	Total	O	0	0
			80	80		
3	C	54	Total	O	0	0
			54	54		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: 3-5 exonuclease PhoExo I



• Molecule 1: 3-5 exonuclease PhoExo I



• Molecule 1: 3-5 exonuclease PhoExo I



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	94.61Å 94.82Å 99.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.57 – 2.15 19.57 – 2.15	Depositor EDS
% Data completeness (in resolution range)	99.8 (19.57-2.15) 99.8 (19.57-2.15)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.49 (at 2.15Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.182 , 0.224 0.182 , 0.224	Depositor DCC
R_{free} test set	2497 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	36.4	Xtriage
Anisotropy	0.859	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 57.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	0.039 for k,h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5484	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.29	0/1814	0.48	0/2449
1	B	0.29	0/1769	0.48	0/2387
1	C	0.28	0/1756	0.46	0/2371
All	All	0.29	0/5339	0.47	0/7207

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1782	0	1849	11	0
1	B	1738	0	1801	16	0
1	C	1725	0	1782	13	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
3	A	102	0	0	1	0
3	B	80	0	0	3	0
3	C	54	0	0	0	0
All	All	5484	0	5432	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:182:LYS:HE3	1:C:184:ILE:HD11	1.77	0.65
1:A:59:ARG:NH2	1:A:115:ASP:OD2	2.31	0.64
1:B:53:SER:OG	1:B:54:GLY:N	2.33	0.61
1:B:39:ARG:NH1	1:B:71:GLU:OE1	2.34	0.60
1:A:22:LEU:HB2	1:A:50:TYR:CZ	2.37	0.59
1:B:182:LYS:HG3	1:B:200:LYS:HD3	1.89	0.54
1:A:2:ARG:HG2	1:A:31:GLU:HG2	1.89	0.54
1:B:56:GLN:OE1	1:B:59:ARG:NH1	2.30	0.53
1:B:97:ASP:O	1:B:106:LYS:NZ	2.37	0.52
1:B:38:LYS:NZ	3:B:466:HOH:O	2.31	0.51
1:A:118:PRO:O	1:A:122:LYS:HG2	2.11	0.50
1:A:83:LEU:HG	1:A:99:LEU:HD13	1.94	0.49
1:B:62:ILE:HG21	1:B:116:LEU:HD22	1.96	0.47
1:A:62:ILE:HG21	1:A:116:LEU:HD22	1.96	0.47
1:C:160:VAL:O	1:C:164:GLY:N	2.41	0.47
1:C:32:LYS:HG3	1:C:33:PRO:HA	1.97	0.47
1:C:88:VAL:HA	1:C:91:LEU:HG	1.95	0.47
1:A:32:LYS:NZ	1:C:224:GLU:OE1	2.41	0.46
1:C:104:ARG:O	1:C:108:ILE:HG13	2.17	0.45
1:B:218:ARG:NH1	3:B:470:HOH:O	2.37	0.45
1:B:59:ARG:O	1:B:63:GLU:HG2	2.17	0.45
1:A:70:ARG:NH2	3:A:484:HOH:O	2.48	0.43
1:B:14:ASP:OD2	1:B:18:GLN:HB2	2.18	0.43
1:C:39:ARG:NH2	1:C:71:GLU:OE2	2.43	0.43
1:B:64:LEU:HD12	1:B:64:LEU:HA	1.88	0.42
1:C:106:LYS:O	1:C:110:LYS:HB2	2.18	0.42
1:A:97:ASP:O	1:A:106:LYS:NZ	2.47	0.42
1:C:4:VAL:HA	1:C:28:VAL:O	2.20	0.42
1:C:29:LEU:HD13	1:C:72:VAL:HG21	2.01	0.42
1:C:153:VAL:HG21	1:C:176:VAL:HG21	2.02	0.42
1:A:10:GLY:HA3	1:A:152:SER:OG	2.20	0.41
1:A:215:PRO:HA	1:B:139:VAL:HB	2.02	0.41
1:B:2:ARG:HD3	3:B:412:HOH:O	2.20	0.41
1:C:157:LEU:HD23	1:C:225:ILE:HD12	2.02	0.41
1:B:166:LEU:HA	1:B:166:LEU:HD12	1.87	0.40
1:B:167:LEU:HA	1:B:167:LEU:HD23	1.92	0.40
1:B:215:PRO:HA	1:C:139:VAL:HB	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	221/229 (96%)	217 (98%)	4 (2%)	0	100	100
1	B	214/229 (93%)	208 (97%)	6 (3%)	0	100	100
1	C	212/229 (93%)	205 (97%)	7 (3%)	0	100	100
All	All	647/687 (94%)	630 (97%)	17 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	193/196 (98%)	181 (94%)	12 (6%)	18	14
1	B	188/196 (96%)	183 (97%)	5 (3%)	44	46
1	C	186/196 (95%)	172 (92%)	14 (8%)	13	8
All	All	567/588 (96%)	536 (94%)	31 (6%)	21	17

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	GLN
1	A	55	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	58	ILE
1	A	83	LEU
1	A	93	GLU
1	A	104	ARG
1	A	109	TRP
1	A	113	SER
1	A	121	LYS
1	A	166	LEU
1	A	180	LYS
1	A	216	LEU
1	B	55	ARG
1	B	91	LEU
1	B	97	ASP
1	B	163	LYS
1	B	229	SER
1	C	15	GLU
1	C	56	GLN
1	C	73	SER
1	C	79	LEU
1	C	92	ASP
1	C	93	GLU
1	C	103	ASP
1	C	109	TRP
1	C	114	LYS
1	C	126	GLU
1	C	137	SER
1	C	161	LYS
1	C	163	LYS
1	C	205	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	205	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	225/229 (98%)	0.22	14 (6%) 20 27	21, 37, 79, 98	0
1	B	220/229 (96%)	0.49	25 (11%) 5 7	27, 46, 98, 119	0
1	C	218/229 (95%)	0.71	35 (16%) 1 2	28, 55, 110, 132	0
All	All	663/687 (96%)	0.47	74 (11%) 5 7	21, 46, 97, 132	0

All (74) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	95	THR	5.6
1	B	108	ILE	5.0
1	C	105	GLY	4.9
1	C	206	GLY	4.9
1	C	92	ASP	4.5
1	C	93	GLU	4.4
1	C	103	ASP	4.2
1	B	82	THR	4.2
1	C	107	GLU	3.9
1	B	97	ASP	3.9
1	B	163	LYS	3.9
1	C	95	THR	3.8
1	A	100	GLN	3.8
1	B	94	SER	3.6
1	C	108	ILE	3.6
1	B	96	ILE	3.5
1	B	15	GLU	3.5
1	C	153	VAL	3.5
1	C	104	ARG	3.3
1	B	53	SER	3.2
1	B	6	ALA	3.2
1	C	226	THR	3.2
1	C	97	ASP	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	162	GLU	3.2
1	C	168	VAL	3.1
1	C	180	LYS	3.0
1	C	127	THR	2.9
1	A	162	GLU	2.9
1	C	162	GLU	2.9
1	A	77	ILE	2.9
1	C	4	VAL	2.8
1	B	49	ASN	2.8
1	C	109	TRP	2.8
1	B	55	ARG	2.8
1	C	94	SER	2.7
1	C	5	ALA	2.7
1	C	84	GLY	2.7
1	C	110	LYS	2.7
1	C	164	GLY	2.7
1	C	131	ILE	2.7
1	C	163	LYS	2.7
1	B	150	ILE	2.6
1	A	161	LYS	2.6
1	C	140	PRO	2.6
1	C	85	GLY	2.6
1	B	83	LEU	2.6
1	C	88	VAL	2.5
1	C	56	GLN	2.5
1	A	163	LYS	2.5
1	A	5	ALA	2.3
1	B	54	GLY	2.3
1	A	131	ILE	2.3
1	C	124	TRP	2.3
1	B	202	GLU	2.3
1	A	229	SER	2.3
1	B	84	GLY	2.3
1	A	50	TYR	2.2
1	C	121	LYS	2.2
1	B	92	ASP	2.2
1	C	126	GLU	2.2
1	B	161	LYS	2.2
1	A	108	ILE	2.2
1	B	106	LYS	2.2
1	A	104	ARG	2.2
1	B	164	GLY	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	50	TYR	2.2
1	A	18	GLN	2.2
1	A	109	TRP	2.2
1	C	181	ASP	2.2
1	B	30	VAL	2.2
1	C	141	VAL	2.1
1	A	168	VAL	2.1
1	B	93	GLU	2.1
1	B	18	GLN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	MG	B	301	1/1	0.92	0.09	56,56,56,56	0
2	MG	C	301	1/1	0.95	0.08	64,64,64,64	0
2	MG	A	301	1/1	0.96	0.02	43,43,43,43	0

6.5 Other polymers [i](#)

There are no such residues in this entry.