



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 13, 2020 – 10:52 PM BST

PDB ID : 5YP3
Title : Crystal structure of dipeptidyl peptidase IV (DPP IV) with Ile-Pro from *Pseudomonas mexicana*
Authors : Roppongi, S.; Suzuki, Y.; Tateoka, C.; Fuimoto, M.; Morisawa, S.; Iizuka, I.; Nakamura, A.; Honma, N.; Shida, Y.; Ogasawara, W.; Tanaka, N.; Sakamoto, Y.; Nonaka, T.
Deposited on : 2017-11-01
Resolution : 2.44 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.14.4.dev1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.14.4.dev1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.44 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1564 (2.46-2.42)
Clashscore	141614	1631 (2.46-2.42)
Ramachandran outliers	138981	1617 (2.46-2.42)
Sidechain outliers	138945	1617 (2.46-2.42)
RSRZ outliers	127900	1547 (2.46-2.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	745	<div> <div>%</div> <div> <div></div> <div>78%</div> <div>17%</div> <div>• •</div> </div> </div>
1	B	745	<div> <div>%</div> <div> <div></div> <div>82%</div> <div>14%</div> <div>• •</div> </div> </div>
1	C	745	<div> <div>2%</div> <div> <div></div> <div>86%</div> <div>10%</div> <div>• •</div> </div> </div>
1	D	745	<div> <div></div> <div> <div></div> <div>81%</div> <div>14%</div> <div>• •</div> </div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 23158 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Dipeptidyl aminopeptidase 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	724	Total	C	N	O	S	0	0	0
			5660	3580	994	1076	10			
1	B	724	Total	C	N	O	S	0	0	0
			5660	3580	994	1076	10			
1	C	724	Total	C	N	O	S	0	0	0
			5660	3580	994	1076	10			
1	D	724	Total	C	N	O	S	0	0	0
			5660	3580	994	1076	10			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	12	ILE	MET	see sequence details	UNP Q6F3I7
B	12	ILE	MET	see sequence details	UNP Q6F3I7
C	12	ILE	MET	see sequence details	UNP Q6F3I7
D	12	ILE	MET	see sequence details	UNP Q6F3I7

- Molecule 2 is ISOLEUCINE (three-letter code: ILE) (formula: C₆H₁₃NO₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			8	6	1	1		
2	B	1	Total	C	N	O	0	0
			8	6	1	1		
2	C	1	Total	C	N	O	0	0
			8	6	1	1		
2	D	1	Total	C	N	O	0	0
			8	6	1	1		

- Molecule 3 is PROLINE (three-letter code: PRO) (formula: C₅H₉NO₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			7	5	1	1		
3	B	1	Total	C	N	O	0	0
			7	5	1	1		
3	C	1	Total	C	N	O	0	0
			7	5	1	1		
3	D	1	Total	C	N	O	0	0
			7	5	1	1		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	D	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	143	Total	O	0	0
			143	143		
5	B	93	Total	O	0	0
			93	93		
5	C	99	Total	O	0	0
			99	99		
5	D	117	Total	O	0	0
			117	117		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Dipeptidyl aminopeptidase 4





- Molecule 1: Dipeptidyl aminopeptidase 4







• Molecule 1: Dipeptidyl aminopeptidase 4



• Molecule 1: Dipeptidyl aminopeptidase 4



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	119.88Å 120.12Å 262.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 2.44 60.06 – 2.44	Depositor EDS
% Data completeness (in resolution range)	97.4 (40.00-2.44) 97.4 (60.06-2.44)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.22 (at 2.45Å)	Xtriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.238 , 0.286 0.243 , 0.286	Depositor DCC
R_{free} test set	6835 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	39.3	Xtriage
Anisotropy	0.342	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 41.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.38$, $\langle L^2 \rangle = 0.21$	Xtriage
Estimated twinning fraction	0.004 for k,h,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	23158	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 39.35 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.2151e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.77	0/5799	0.95	10/7887 (0.1%)
1	B	0.70	0/5799	0.90	8/7887 (0.1%)
1	C	0.68	0/5799	0.87	7/7887 (0.1%)
1	D	0.75	0/5799	0.96	13/7887 (0.2%)
All	All	0.72	0/23196	0.92	38/31548 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	D	0	2
All	All	0	4

There are no bond length outliers.

The worst 5 of 38 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	104	ARG	NE-CZ-NH2	-14.85	112.87	120.30
1	D	104	ARG	NE-CZ-NH1	10.91	125.76	120.30
1	B	104	ARG	NE-CZ-NH2	9.40	125.00	120.30
1	B	104	ARG	NE-CZ-NH1	-8.78	115.91	120.30
1	D	309	ARG	NE-CZ-NH1	7.36	123.98	120.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	192	ASP	Peptide
1	B	638	PRO	Peptide
1	D	164	LYS	Peptide
1	D	638	PRO	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5660	0	5521	70	0
1	B	5660	0	5521	63	0
1	C	5660	0	5521	35	0
1	D	5660	0	5521	63	0
2	A	8	0	10	1	0
2	B	8	0	10	1	0
2	C	8	0	10	1	0
2	D	8	0	10	0	0
3	A	7	0	7	0	0
3	B	7	0	7	0	0
3	C	7	0	7	0	0
3	D	7	0	7	0	0
4	D	6	0	8	0	0
5	A	143	0	0	3	0
5	B	93	0	0	3	0
5	C	99	0	0	0	0
5	D	117	0	0	1	0
All	All	23158	0	22160	226	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 226 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:262:ASN:ND2	1:D:647:THR:HG21	1.59	1.16
1:D:262:ASN:HD21	1:D:647:THR:HG21	0.96	1.11
1:B:262:ASN:HD21	1:B:647:THR:HG21	1.00	1.09

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:262:ASN:ND2	1:B:647:THR:HG21	1.75	1.01
1:D:262:ASN:HD21	1:D:647:THR:CG2	1.79	0.96

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	722/745 (97%)	669 (93%)	49 (7%)	4 (1%)	25	29
1	B	722/745 (97%)	661 (92%)	58 (8%)	3 (0%)	34	41
1	C	722/745 (97%)	664 (92%)	55 (8%)	3 (0%)	34	41
1	D	722/745 (97%)	666 (92%)	53 (7%)	3 (0%)	34	41
All	All	2888/2980 (97%)	2660 (92%)	215 (7%)	13 (0%)	29	34

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	542	ARG
1	A	222	ASP
1	C	543	SER
1	A	181	LEU
1	A	369	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	587/602 (98%)	546 (93%)	41 (7%)	15	18
1	B	587/602 (98%)	555 (94%)	32 (6%)	21	28
1	C	587/602 (98%)	562 (96%)	25 (4%)	29	38
1	D	587/602 (98%)	552 (94%)	35 (6%)	19	25
All	All	2348/2408 (98%)	2215 (94%)	133 (6%)	21	26

5 of 133 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	416	LEU
1	C	88	LEU
1	D	577	LEU
1	B	445	SER
1	B	563	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 41 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	548	ASN
1	C	45	GLN
1	D	487	HIS
1	B	549	GLN
1	B	553	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PRO	D	802	1,2	5,7,8	0.54	0	7,8,10	1.34	1 (14%)
3	PRO	A	802	1,2	5,7,8	0.61	0	7,8,10	1.95	2 (28%)
3	PRO	B	802	1,2	5,7,8	0.78	0	7,8,10	1.29	1 (14%)
4	GOL	D	803	-	5,5,5	1.01	0	5,5,5	1.27	0
2	ILE	B	801	3	6,7,8	0.66	0	5,8,10	1.75	2 (40%)
2	ILE	D	801	3	6,7,8	0.85	0	5,8,10	2.23	2 (40%)
3	PRO	C	802	1,2	5,7,8	0.50	0	7,8,10	2.64	3 (42%)
2	ILE	A	801	3	6,7,8	0.66	0	5,8,10	2.10	1 (20%)
2	ILE	C	801	3	6,7,8	0.73	0	5,8,10	2.57	2 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PRO	D	802	1,2	-	0/0/9/11	0/1/1/1
3	PRO	A	802	1,2	-	0/0/9/11	0/1/1/1
3	PRO	B	802	1,2	-	0/0/9/11	0/1/1/1
4	GOL	D	803	-	-	4/4/4/4	-
2	ILE	B	801	3	-	2/7/8/10	-
2	ILE	D	801	3	-	2/7/8/10	-
3	PRO	C	802	1,2	-	0/0/9/11	0/1/1/1
2	ILE	A	801	3	-	3/7/8/10	-
2	ILE	C	801	3	-	1/7/8/10	-

There are no bond length outliers.

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	C	801	ILE	O-C-CA	-4.54	112.88	124.78
3	C	802	PRO	CB-CA-C	-4.50	106.51	112.70
2	A	801	ILE	O-C-CA	-4.17	113.84	124.78
3	C	802	PRO	O-C-CA	-4.08	114.08	124.78
3	A	802	PRO	CB-CA-C	-3.85	107.40	112.70

There are no chirality outliers.

5 of 12 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	801	ILE	O-C-CA-CB
2	A	801	ILE	CG2-CB-CG1-CD1
2	A	801	ILE	CA-CB-CG1-CD1
4	D	803	GOL	O1-C1-C2-C3
4	D	803	GOL	C1-C2-C3-O3

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	801	ILE	1	0
2	A	801	ILE	1	0
2	C	801	ILE	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	724/745 (97%)	-0.01	11 (1%) 73 71	21, 42, 68, 105	0
1	B	724/745 (97%)	0.06	7 (0%) 82 81	24, 45, 68, 106	0
1	C	724/745 (97%)	0.09	13 (1%) 68 64	22, 44, 73, 94	0
1	D	724/745 (97%)	-0.09	3 (0%) 92 92	18, 39, 61, 99	0
All	All	2896/2980 (97%)	0.01	34 (1%) 79 77	18, 42, 68, 106	0

The worst 5 of 34 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	241	TYR	4.4
1	C	245	PRO	4.3
1	A	241	TYR	4.2
1	A	245	PRO	4.2
1	A	142	GLY	3.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	GOL	D	803	6/6	0.79	0.27	42,53,57,58	0
3	PRO	A	802	7/8	0.92	0.20	23,24,24,24	0
3	PRO	C	802	7/8	0.93	0.15	27,28,29,29	0
2	ILE	D	801	8/9	0.94	0.18	21,24,25,27	0
3	PRO	B	802	7/8	0.95	0.13	24,25,28,30	0
2	ILE	B	801	8/9	0.95	0.20	23,25,25,27	0
2	ILE	C	801	8/9	0.95	0.15	24,26,26,27	0
2	ILE	A	801	8/9	0.96	0.17	24,24,26,26	0
3	PRO	D	802	7/8	0.97	0.10	24,24,26,26	0

6.5 Other polymers

There are no such residues in this entry.