



Full wwPDB X-ray Structure Validation Report ⓘ

May 24, 2020 – 10:49 pm BST

PDB ID : 5YQ2
Title : Crystal structure of E.coli aminopeptidase N in complex with Puromycin aminonucleoside
Authors : Marapaka, A.K.; Ganji, R.J.; Reddi, R.; Addlagatta, A.
Deposited on : 2017-11-04
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	891	<div> <div>2%</div> <div> <div></div> <div>85%</div> <div>11%</div> <div>..</div> </div> </div>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7870 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aminopeptidase N.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	866	7016	4445	1204	1338	29	0	17	0


There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	initiating methionine	UNP P04825
A	-19	GLY	-	expression tag	UNP P04825
A	-18	SER	-	expression tag	UNP P04825
A	-17	SER	-	expression tag	UNP P04825
A	-16	HIS	-	expression tag	UNP P04825
A	-15	HIS	-	expression tag	UNP P04825
A	-14	HIS	-	expression tag	UNP P04825
A	-13	HIS	-	expression tag	UNP P04825
A	-12	HIS	-	expression tag	UNP P04825
A	-11	HIS	-	expression tag	UNP P04825
A	-10	SER	-	expression tag	UNP P04825
A	-9	SER	-	expression tag	UNP P04825
A	-8	GLY	-	expression tag	UNP P04825
A	-7	LEU	-	expression tag	UNP P04825
A	-6	VAL	-	expression tag	UNP P04825
A	-5	PRO	-	expression tag	UNP P04825
A	-4	ARG	-	expression tag	UNP P04825
A	-3	GLY	-	expression tag	UNP P04825
A	-2	SER	-	expression tag	UNP P04825
A	-1	HIS	-	expression tag	UNP P04825
A	0	MET	-	expression tag	UNP P04825

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is (2R,3R,4S,5S)-4-AMINO-2-[6-(DIMETHYLAMINO)-9H-PURIN-9-YL]-5-(HYDROXYMETHYL)TETRAHYDRO-3-FURANOL (three-letter code: GMC) (formula: $C_{12}H_{18}N_6O_3$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			21	12	6	3		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is MALONATE ION (three-letter code: MLI) (formula: $C_3H_2O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			7	3	4		
5	A	1	Total	C	O	0	0
			7	3	4		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	806	Total	O	0	0
			806	806		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Aminopeptidase N



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	120.22Å 120.22Å 170.12Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 1.60 29.61 – 1.60	Depositor EDS
% Data completeness (in resolution range)	99.7 (50.00-1.60) 99.7 (29.61-1.60)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.51 (at 1.60Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.157 , 0.179 0.171 , 0.189	Depositor DCC
R_{free} test set	9525 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å ²)	22.8	Xtriage
Anisotropy	0.003	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 44.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	0.006 for -h,-k,l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7870	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.01% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GMC, GOL, ZN, MLI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.40	27/7222 (0.4%)	1.49	98/9805 (1.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (27) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	260[A]	MET	C-O	10.51	1.43	1.23
1	A	260[B]	MET	C-O	10.51	1.43	1.23
1	A	261	GLY	N-CA	-10.20	1.30	1.46
1	A	543[A]	GLU	CD-OE1	9.50	1.36	1.25
1	A	543[B]	GLU	CD-OE1	9.50	1.36	1.25
1	A	734[A]	GLU	N-CA	-7.63	1.31	1.46
1	A	734[B]	GLU	N-CA	-7.63	1.31	1.46
1	A	525	TYR	CG-CD2	7.28	1.48	1.39
1	A	525	TYR	CG-CD1	7.20	1.48	1.39
1	A	107	GLU	CD-OE1	6.73	1.33	1.25
1	A	207	ARG	CZ-NH1	6.67	1.41	1.33
1	A	69	GLU	CD-OE1	-6.42	1.18	1.25
1	A	564	ARG	CD-NE	-6.38	1.35	1.46
1	A	525	TYR	CB-CG	6.33	1.61	1.51
1	A	585	GLN	CG-CD	6.29	1.65	1.51
1	A	299	TYR	CE1-CZ	-6.17	1.30	1.38
1	A	615	GLU	CD-OE2	6.16	1.32	1.25
1	A	204[A]	ARG	CZ-NH2	6.01	1.40	1.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	204[B]	ARG	CZ-NH2	6.01	1.40	1.33
1	A	564	ARG	CZ-NH1	5.97	1.40	1.33
1	A	525	TYR	CE1-CZ	5.75	1.46	1.38
1	A	275	TYR	CZ-OH	5.61	1.47	1.37
1	A	97	GLU	CD-OE1	-5.40	1.19	1.25
1	A	69	GLU	CD-OE2	-5.30	1.19	1.25
1	A	168	GLU	C-O	5.19	1.33	1.23
1	A	546	TRP	CA-CB	-5.17	1.42	1.53
1	A	490	TYR	CE1-CZ	-5.14	1.31	1.38

All (98) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	556	ARG	NE-CZ-NH1	21.57	131.09	120.30
1	A	556	ARG	NE-CZ-NH2	-18.35	111.12	120.30
1	A	349	ARG	NE-CZ-NH1	-17.26	111.67	120.30
1	A	204[A]	ARG	NE-CZ-NH2	14.12	127.36	120.30
1	A	204[B]	ARG	NE-CZ-NH2	14.12	127.36	120.30
1	A	624	GLU	OE1-CD-OE2	12.72	138.57	123.30
1	A	837	ASP	CB-CG-OD1	12.27	129.34	118.30
1	A	432	ASP	CB-CG-OD1	11.95	129.05	118.30
1	A	126	ARG	NE-CZ-NH2	11.75	126.17	120.30
1	A	293	ARG	NE-CZ-NH1	11.53	126.06	120.30
1	A	248	ASP	CB-CG-OD1	-11.22	108.20	118.30
1	A	307	ARG	NE-CZ-NH2	-10.97	114.82	120.30
1	A	207	ARG	NE-CZ-NH1	10.55	125.58	120.30
1	A	437	ARG	NE-CZ-NH1	10.44	125.52	120.30
1	A	556	ARG	CD-NE-CZ	10.24	137.94	123.60
1	A	161	ARG	NE-CZ-NH2	-10.15	115.23	120.30
1	A	564	ARG	NE-CZ-NH2	-10.10	115.25	120.30
1	A	566	ASP	CB-CG-OD2	-10.06	109.24	118.30
1	A	615	GLU	OE1-CD-OE2	9.45	134.64	123.30
1	A	438	ARG	NE-CZ-NH1	8.79	124.69	120.30
1	A	825	ARG	NE-CZ-NH1	8.67	124.63	120.30
1	A	10	ARG	NE-CZ-NH1	8.44	124.52	120.30
1	A	349	ARG	NE-CZ-NH2	8.34	124.47	120.30
1	A	830	LEU	CB-CG-CD1	8.32	125.14	111.00
1	A	453	ASP	CB-CG-OD1	-8.24	110.88	118.30
1	A	365	ARG	NE-CZ-NH1	8.13	124.36	120.30
1	A	256	ASP	CB-CG-OD1	7.96	125.46	118.30
1	A	367	ASP	CB-CG-OD1	7.95	125.46	118.30
1	A	845	ARG	NE-CZ-NH2	-7.88	116.36	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	240	ARG	NE-CZ-NH2	-7.85	116.38	120.30
1	A	220	LEU	CB-CG-CD1	7.68	124.06	111.00
1	A	216	ASP	CB-CG-OD1	7.56	125.11	118.30
1	A	353	PHE	CB-CG-CD1	7.50	126.05	120.80
1	A	18	TYR	CB-CG-CD2	-7.45	116.53	121.00
1	A	410	ARG	NE-CZ-NH2	-7.35	116.63	120.30
1	A	556	ARG	CG-CD-NE	-7.30	96.48	111.80
1	A	199	ASP	CB-CG-OD2	-7.22	111.80	118.30
1	A	258	PHE	CB-CG-CD1	-7.16	115.79	120.80
1	A	438	ARG	NE-CZ-NH2	-7.12	116.74	120.30
1	A	114	ASP	CB-CG-OD2	-6.96	112.03	118.30
1	A	327	ASP	CB-CG-OD1	6.95	124.56	118.30
1	A	633	ASP	CB-CG-OD1	6.94	124.55	118.30
1	A	260[A]	MET	CA-CB-CG	6.94	125.09	113.30
1	A	260[B]	MET	CA-CB-CG	6.94	125.09	113.30
1	A	293	ARG	NE-CZ-NH2	-6.93	116.84	120.30
1	A	307	ARG	NE-CZ-NH1	6.78	123.69	120.30
1	A	474	ASP	CB-CG-OD1	-6.77	112.20	118.30
1	A	491	ASP	CB-CG-OD1	6.77	124.39	118.30
1	A	548	ASP	CB-CG-OD1	6.76	124.38	118.30
1	A	845	ARG	NE-CZ-NH1	6.71	123.66	120.30
1	A	712	ASP	CB-CG-OD1	6.64	124.28	118.30
1	A	695	LEU	CB-CG-CD1	6.61	122.23	111.00
1	A	575	TYR	CB-CG-CD2	6.51	124.91	121.00
1	A	522	ASP	CB-CG-OD1	6.43	124.08	118.30
1	A	171	ARG	NE-CZ-NH1	6.40	123.50	120.30
1	A	491	ASP	CB-CG-OD2	-6.32	112.61	118.30
1	A	154	PHE	CB-CG-CD2	-6.30	116.39	120.80
1	A	260[A]	MET	CB-CG-SD	6.25	131.16	112.40
1	A	260[B]	MET	CB-CG-SD	6.25	131.16	112.40
1	A	736	ASP	CB-CG-OD1	6.25	123.92	118.30
1	A	133	ASP	CB-CG-OD1	-6.21	112.71	118.30
1	A	214	TYR	CB-CG-CD2	-6.11	117.33	121.00
1	A	346	ARG	NE-CZ-NH1	6.08	123.34	120.30
1	A	728	ARG	NE-CZ-NH2	-6.07	117.26	120.30
1	A	256	ASP	CB-CG-OD2	-6.05	112.85	118.30
1	A	543[A]	GLU	CG-CD-OE1	5.94	130.18	118.30
1	A	543[B]	GLU	CG-CD-OE1	5.94	130.18	118.30
1	A	337	ARG	NE-CZ-NH1	5.93	123.27	120.30
1	A	522	ASP	CB-CG-OD2	-5.90	112.99	118.30
1	A	10	ARG	NE-CZ-NH2	-5.90	117.35	120.30
1	A	678	ARG	NE-CZ-NH2	-5.89	117.35	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	609	ASP	CB-CG-OD1	5.89	123.60	118.30
1	A	204[A]	ARG	NE-CZ-NH1	-5.88	117.36	120.30
1	A	204[B]	ARG	NE-CZ-NH1	-5.88	117.36	120.30
1	A	772	ARG	NE-CZ-NH2	5.82	123.21	120.30
1	A	806	LEU	CB-CG-CD2	5.80	120.87	111.00
1	A	772	ARG	NE-CZ-NH1	-5.78	117.41	120.30
1	A	14	ARG	NE-CZ-NH1	5.78	123.19	120.30
1	A	654	ASP	CB-CG-OD2	-5.76	113.11	118.30
1	A	536	PHE	CB-CG-CD1	5.75	124.83	120.80
1	A	553	PHE	CB-CG-CD1	-5.71	116.81	120.80
1	A	148	ASP	CB-CG-OD1	-5.65	113.21	118.30
1	A	389	ARG	NE-CZ-NH1	-5.64	117.48	120.30
1	A	419	ASP	CB-CG-OD2	-5.56	113.30	118.30
1	A	624	GLU	CG-CD-OE2	-5.47	107.35	118.30
1	A	440	TYR	CZ-CE2-CD2	5.40	124.66	119.80
1	A	783	ARG	NE-CZ-NH2	-5.39	117.60	120.30
1	A	686	ARG	NE-CZ-NH2	-5.38	117.61	120.30
1	A	126	ARG	NE-CZ-NH1	-5.38	117.61	120.30
1	A	825	ARG	NE-CZ-NH2	-5.27	117.66	120.30
1	A	207	ARG	NE-CZ-NH2	-5.21	117.69	120.30
1	A	356	ASP	O-C-N	5.18	130.99	122.70
1	A	837	ASP	OD1-CG-OD2	-5.18	113.46	123.30
1	A	90	PHE	CB-CG-CD2	5.13	124.39	120.80
1	A	825	ARG	CG-CD-NE	5.10	122.50	111.80
1	A	830	LEU	CA-CB-CG	5.05	126.92	115.30
1	A	774	PHE	CB-CG-CD1	5.01	124.31	120.80
1	A	729	ASP	CB-CG-OD1	5.00	122.80	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	556	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7016	0	6874	26	0
2	A	1	0	0	0	0
3	A	21	0	18	2	0
4	A	12	0	16	0	0
5	A	14	0	4	0	0
6	A	806	0	0	8	0
All	All	7870	0	6912	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:279:ARG:NH1	1:A:281[A]:ASP:OD2	2.02	0.92
1:A:279:ARG:NH1	1:A:281[A]:ASP:CG	2.45	0.69
1:A:207:ARG:NH2	6:A:1002:HOH:O	2.24	0.68
1:A:279:ARG:NH1	1:A:281[A]:ASP:OD1	2.26	0.68
1:A:488[B]:GLU:OE2	1:A:559:ARG:NH2	2.28	0.66
1:A:207:ARG:NH1	6:A:1002:HOH:O	2.27	0.65
3:A:902:GMC:N7	3:A:902:GMC:H101	2.10	0.64
1:A:28[A]:ASP:OD2	1:A:171:ARG:NH2	2.38	0.57
1:A:260[B]:MET:HE3	6:A:1651:HOH:O	2.08	0.53
1:A:862:TYR:O	1:A:866:THR:HG23	2.12	0.50
1:A:734[A]:GLU:HG3	6:A:1194:HOH:O	2.13	0.49
1:A:28[A]:ASP:CG	1:A:171:ARG:NH2	2.68	0.47
1:A:207:ARG:CZ	6:A:1002:HOH:O	2.58	0.46
1:A:608:ILE:CG2	6:A:1168:HOH:O	2.64	0.46
1:A:714:LEU:HD22	1:A:748:LYS:HD3	1.96	0.45
1:A:75:LYS:HE3	1:A:77:GLU:OE2	2.18	0.44
1:A:605:ASP:O	1:A:608:ILE:HG22	2.17	0.44
1:A:495:LYS:HA	1:A:495:LYS:HE3	2.00	0.43
1:A:279:ARG:HG3	1:A:281[A]:ASP:OD1	2.19	0.43
1:A:492:ASN:HB3	1:A:526:PHE:CE2	2.53	0.42
1:A:220:LEU:HB2	6:A:1296:HOH:O	2.19	0.42
1:A:28[A]:ASP:OD1	1:A:171:ARG:NH2	2.49	0.42
1:A:93:LYS:HE3	6:A:1704:HOH:O	2.19	0.41
1:A:30:ASP:HB3	1:A:33:LYS:O	2.19	0.41
1:A:825:ARG:HH21	3:A:902:GMC:C2	2.33	0.41
1:A:265:ASN:HB2	1:A:268:LEU:O	2.21	0.41
1:A:28[A]:ASP:CG	1:A:171:ARG:HH21	2.24	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	881/891 (99%)	868 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	758/763 (99%)	742 (98%)	16 (2%)	53	29

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	164	GLN
1	A	172	HIS
1	A	220	LEU
1	A	260[A]	MET
1	A	260[B]	MET
1	A	327	ASP
1	A	356	ASP
1	A	477	GLU
1	A	493	GLU
1	A	495	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	517	GLN
1	A	695	LEU
1	A	699	LEU
1	A	825	ARG
1	A	830	LEU
1	A	860	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	665	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GMC	A	902	-	18,23,23	2.09	7 (38%)	16,34,34	3.35	5 (31%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	MLI	A	906	-	0,6,6	0.00	-	0,7,7	0.00	-
4	GOL	A	903	-	5,5,5	0.35	0	5,5,5	0.62	0
4	GOL	A	904	-	5,5,5	1.14	0	5,5,5	1.01	0
5	MLI	A	905	-	0,6,6	0.00	-	0,7,7	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GMC	A	902	-	-	2/6/26/26	0/3/3/3
5	MLI	A	906	-	-	0/0/4/4	-
4	GOL	A	903	-	-	0/4/4/4	-
4	GOL	A	904	-	-	0/4/4/4	-
5	MLI	A	905	-	-	0/0/4/4	-

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	902	GMC	O4'-C4'	-5.14	1.33	1.45
3	A	902	GMC	C6-N1	3.21	1.37	1.33
3	A	902	GMC	C2-N3	2.99	1.36	1.32
3	A	902	GMC	C2'-C3'	-2.80	1.49	1.53
3	A	902	GMC	C2'-C1'	2.67	1.57	1.53
3	A	902	GMC	C3'-N3'	2.34	1.50	1.47
3	A	902	GMC	O2'-C2'	-2.20	1.37	1.43

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	902	GMC	C5'-C4'-C3'	-7.72	105.06	115.67
3	A	902	GMC	O4'-C4'-C3'	7.68	115.16	104.15
3	A	902	GMC	N1-C6-N6	5.38	122.72	117.06
3	A	902	GMC	O2'-C2'-C1'	-2.81	100.48	110.85
3	A	902	GMC	C9-N6-C10	2.49	124.14	116.12

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	902	GMC	C5-C6-N6-C9
3	A	902	GMC	C5-C6-N6-C10

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	902	GMC	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	866/891 (97%)	-0.28	16 (1%) 68 67	16, 22, 36, 62	1 (0%)

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	504	HIS	4.9
1	A	169	ASN	4.8
1	A	168	GLU	3.8
1	A	694	HIS	3.5
1	A	606	GLU	3.4
1	A	503	GLY	3.3
1	A	855	GLU	3.0
1	A	260[A]	MET	3.0
1	A	249	ILE	2.9
1	A	5	PRO	2.9
1	A	525	TYR	2.8
1	A	502	GLY	2.5
1	A	493	GLU	2.3
1	A	856	ASN	2.3
1	A	45	HIS	2.3
1	A	492	ASN	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	GOL	A	903	6/6	0.59	0.30	57,64,65,66	0
3	GMC	A	902	21/21	0.81	0.15	22,32,45,51	21
5	MLI	A	906	7/7	0.82	0.24	35,52,60,62	0
4	GOL	A	904	6/6	0.86	0.11	26,33,35,36	0
5	MLI	A	905	7/7	0.91	0.10	21,25,27,27	7
2	ZN	A	901	1/1	1.00	0.03	16,16,16,16	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.