



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2021 – 01:17 PM GMT

PDB ID : 6YQ9
Title : Taka-amylase in complex with alpha-glucosyl epi-cyclophellitol epoxide inhibitor
Authors : Armstrong, Z.; Chen, Y.; Artola, M.; Overkleeft, H.; Davies, G.
Deposited on : 2020-04-16
Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.16
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.16


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1483 (1.56-1.56)
Clashscore	141614	1529 (1.56-1.56)
Ramachandran outliers	138981	1498 (1.56-1.56)
Sidechain outliers	138945	1495 (1.56-1.56)
RSRZ outliers	127900	1465 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	499	
1	BBB	499	

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 15539 atoms, of which 7152 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alpha-amylase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	AAA	476	Total	C	H	N	O	S	220	10	0
			7305	2375	3547	609	756	18			
1	BBB	476	Total	C	H	N	O	S	217	6	0
			7229	2355	3507	600	749	18			

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	AAA	1	Total	C	H	N	O	3	0
			28	8	14	1	5		
2	BBB	1	Total	C	H	N	O	3	0
			28	8	14	1	5		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	AAA	1	Total	C	H	O	1	0
			10	2	6	2		
3	BBB	1	Total	C	H	O	1	0
			10	2	6	2		
3	BBB	1	Total	C	H	O	1	0
			10	2	6	2		
3	BBB	1	Total	C	H	O	1	0
			10	2	6	2		

- Molecule 4 is (1R,2R,3S,5R,6S)-2,3,5-trihydroxy-6-(hydroxymethyl)cyclohexyl alpha-D-glucopyranoside (three-letter code: 5QP) (formula: C₁₃H₂₄O₁₀) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	AAA	1	Total	C	H	O	7	0
			46	13	23	10		
4	BBB	1	Total	C	H	O	7	0
			46	13	23	10		

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	BBB	1	Total	Ca	0	0
			1	1		
5	AAA	1	Total	Ca	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	AAA	416	Total	O	0	0
			416	416		
6	BBB	399	Total	O	0	0
			399	399		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alpha-amylase



- Molecule 1: Alpha-amylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	65.21Å 103.10Å 75.37Å 90.00° 103.63° 90.00°	Depositor
Resolution (Å)	73.36 – 1.55 73.25 – 1.55	Depositor EDS
% Data completeness (in resolution range)	99.9 (73.36-1.55) 99.9 (73.25-1.55)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.73 (at 1.55Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.146 , 0.174 0.160 , 0.182	Depositor DCC
R_{free} test set	6862 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	15.6	Xtriage
Anisotropy	0.775	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 40.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	15539	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.34% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: 5QP, CA, NAG, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	AAA	0.76	0/3858	0.92	4/5267 (0.1%)
1	BBB	0.74	0/3822	0.90	2/5221 (0.0%)
All	All	0.75	0/7680	0.91	6/10488 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	AAA	397[A]	ARG	NE-CZ-NH2	7.01	123.80	120.30
1	AAA	397[B]	ARG	NE-CZ-NH2	7.01	123.80	120.30
1	AAA	397[A]	ARG	NE-CZ-NH1	-5.77	117.42	120.30
1	AAA	397[B]	ARG	NE-CZ-NH1	-5.77	117.42	120.30
1	BBB	397	ARG	NE-CZ-NH1	-5.48	117.56	120.30
1	BBB	397	ARG	NE-CZ-NH2	5.05	122.82	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	3758	3547	3533	14	0
1	BBB	3722	3507	3495	19	0
2	AAA	14	14	13	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	BBB	14	14	13	0	0
3	AAA	4	6	6	0	0
3	BBB	12	18	18	1	0
4	AAA	23	23	0	5	0
4	BBB	23	23	0	7	0
5	AAA	1	0	0	0	0
5	BBB	1	0	0	0	0
6	AAA	416	0	0	3	0
6	BBB	399	0	0	5	0
All	All	8387	7152	7078	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:BBB:206[A]:ASP:OD1	4:BBB:504:5QP:CAN	1.95	1.14
1:AAA:206[A]:ASP:OD1	1:AAA:207:THR:OG1	1.72	1.08
4:AAA:503:5QP:CAR	6:AAA:601:HOH:O	2.09	1.01
4:BBB:504:5QP:CAR	6:BBB:601:HOH:O	2.14	0.94
4:AAA:503:5QP:OAB	6:AAA:601:HOH:O	1.90	0.88
4:AAA:503:5QP:CAK	6:AAA:601:HOH:O	2.37	0.73
1:BBB:206[A]:ASP:OD2	1:BBB:207:THR:OG1	2.07	0.72
3:BBB:505:EDO:H12	6:BBB:690:HOH:O	1.91	0.70
1:BBB:264:SER:HB2	6:BBB:940:HOH:O	1.92	0.68
4:BBB:504:5QP:OAB	6:BBB:601:HOH:O	2.16	0.61
1:AAA:397[B]:ARG:HH21	1:AAA:427:GLY:HA3	1.67	0.59
1:BBB:206[A]:ASP:OD2	4:BBB:504:5QP:OAG	2.24	0.56
1:AAA:206[A]:ASP:CG	1:AAA:207:THR:OG1	2.44	0.55
1:AAA:397[A]:ARG:HD3	1:AAA:397[A]:ARG:C	2.26	0.55
4:BBB:504:5QP:CAK	6:BBB:601:HOH:O	2.57	0.53
1:BBB:206[B]:ASP:OD1	1:BBB:230:GLU:OE1	2.29	0.51
1:BBB:361:LEU:C	1:BBB:361:LEU:HD13	2.32	0.50
1:BBB:270:ASP:OD1	1:BBB:389:LYS:HD2	2.12	0.49
1:AAA:319[B]:ILE:O	1:AAA:369:ARG:HD2	2.13	0.49
1:AAA:206[A]:ASP:OD1	4:AAA:503:5QP:OAG	2.33	0.47
1:BBB:119:VAL:HG22	1:BBB:206[A]:ASP:HB3	1.97	0.46
1:BBB:206[A]:ASP:OD2	4:BBB:504:5QP:CAR	2.64	0.46
1:BBB:319[B]:ILE:O	1:BBB:369:ARG:HD2	2.17	0.45
1:AAA:206[A]:ASP:OD1	4:AAA:503:5QP:CAR	2.64	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:302:ALA:HA	1:AAA:305:THR:O	2.16	0.45
1:AAA:119:VAL:HG22	1:AAA:206[A]:ASP:HB3	1.99	0.44
1:AAA:397[B]:ARG:NH2	1:AAA:427:GLY:HA3	2.32	0.44
1:BBB:105:SER:O	1:BBB:109:GLU:HG3	2.19	0.43
1:BBB:204:ARG:NH1	1:BBB:206[B]:ASP:HB2	2.34	0.43
1:BBB:119:VAL:HG22	1:BBB:206[A]:ASP:CB	2.49	0.42
1:AAA:37:TYR:CE1	1:AAA:79:TYR:HA	2.55	0.42
1:AAA:296:HIS:N	1:AAA:296:HIS:ND1	2.65	0.41
1:BBB:352:TYR:N	1:BBB:353:PRO:CD	2.83	0.41
1:AAA:115:MET:HA	1:AAA:202:GLY:O	2.21	0.41
1:BBB:204:ARG:HH12	1:BBB:206[B]:ASP:CG	2.24	0.41
1:BBB:206[A]:ASP:CG	4:BBB:504:5QP:CAN	2.72	0.41
1:BBB:302:ALA:HA	1:BBB:305:THR:O	2.20	0.41
1:BBB:176:LEU:HD13	1:BBB:187:TRP:HE1	1.86	0.40
1:BBB:37:TYR:CE1	1:BBB:79:TYR:HA	2.57	0.40
1:AAA:129:GLY:HA3	1:AAA:149:PHE:CZ	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	484/499 (97%)	475 (98%)	9 (2%)	0	100	100
1	BBB	480/499 (96%)	468 (98%)	12 (2%)	0	100	100
All	All	964/998 (97%)	943 (98%)	21 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	407/414 (98%)	401 (98%)	6 (2%)	65	37
1	BBB	402/414 (97%)	395 (98%)	7 (2%)	60	32
All	All	809/828 (98%)	796 (98%)	13 (2%)	65	35

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	61	TRP
1	AAA	207	THR
1	AAA	220	ASN
1	AAA	397[A]	ARG
1	AAA	397[B]	ARG
1	AAA	443	VAL
1	BBB	61	TRP
1	BBB	207	THR
1	BBB	220	ASN
1	BBB	232	LEU
1	BBB	264	SER
1	BBB	293	VAL
1	BBB	436	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	EDO	BBB	505	-	3,3,3	0.50	0	2,2,2	0.13	0
3	EDO	AAA	502	-	3,3,3	0.39	0	2,2,2	0.70	0
2	NAG	BBB	501	1	14,14,15	1.08	1 (7%)	17,19,21	2.27	9 (52%)
4	5QP	BBB	504	1	24,24,24	1.39	4 (16%)	32,35,35	1.24	4 (12%)
3	EDO	BBB	503	-	3,3,3	0.50	0	2,2,2	0.89	0
4	5QP	AAA	503	1	24,24,24	1.27	4 (16%)	32,35,35	1.39	5 (15%)
3	EDO	BBB	502	-	3,3,3	0.32	0	2,2,2	0.04	0
2	NAG	AAA	501	1	14,14,15	0.92	0	17,19,21	1.50	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	BBB	505	-	-	1/1/1/1	-
3	EDO	AAA	502	-	-	0/1/1/1	-
2	NAG	BBB	501	1	-	0/6/23/26	0/1/1/1
4	5QP	BBB	504	1	-	0/8/48/48	0/2/2/2
3	EDO	BBB	503	-	-	1/1/1/1	-
4	5QP	AAA	503	1	-	0/8/48/48	0/2/2/2
3	EDO	BBB	502	-	-	0/1/1/1	-
2	NAG	AAA	501	1	-	0/6/23/26	0/1/1/1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	AAA	503	5QP	CAN-CAO	3.29	1.58	1.52
4	BBB	504	5QP	CAO-CAT	3.23	1.57	1.52
4	BBB	504	5QP	CAN-CAR	-2.47	1.48	1.52
4	BBB	504	5QP	O4-C4	2.39	1.48	1.43
4	AAA	503	5QP	CAO-CAT	2.34	1.56	1.52
2	BBB	501	NAG	O5-C5	2.29	1.48	1.43
4	AAA	503	5QP	CAN-CAR	-2.24	1.48	1.52
4	BBB	504	5QP	CAV-CAR	-2.22	1.48	1.53
4	AAA	503	5QP	CAV-CAX	-2.03	1.49	1.52

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	BBB	501	NAG	C6-C5-C4	-4.61	102.21	113.00
4	AAA	503	5QP	CAK-CAV-CAR	3.48	116.16	111.99
2	AAA	501	NAG	C1-O5-C5	-3.41	107.57	112.19
2	BBB	501	NAG	C1-O5-C5	3.40	116.80	112.19
4	BBB	504	5QP	C6-C5-C4	3.36	120.87	113.00
2	BBB	501	NAG	O5-C1-C2	-3.00	106.55	111.29
2	BBB	501	NAG	O3-C3-C4	-2.80	103.88	110.35
2	BBB	501	NAG	C8-C7-N2	-2.72	111.50	116.10
4	AAA	503	5QP	C6-C5-C4	2.71	119.35	113.00
2	BBB	501	NAG	O4-C4-C5	2.70	116.00	109.30
4	AAA	503	5QP	C4-C3-C2	2.64	115.44	110.82
4	BBB	504	5QP	C1-O1-CAX	-2.54	111.68	117.96
2	BBB	501	NAG	O7-C7-N2	2.42	126.40	121.95
2	BBB	501	NAG	O5-C5-C4	2.31	116.45	110.83
4	BBB	504	5QP	O4-C4-C3	2.21	115.46	110.35
2	AAA	501	NAG	O3-C3-C4	2.16	115.33	110.35
4	AAA	503	5QP	O1-CAX-CAV	2.13	114.91	108.63
4	BBB	504	5QP	C4-C3-C2	2.11	114.50	110.82
2	BBB	501	NAG	C3-C4-C5	-2.09	106.50	110.24
4	AAA	503	5QP	CAN-CAR-CAV	2.04	118.11	112.65

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	BBB	505	EDO	O1-C1-C2-O2
3	BBB	503	EDO	O1-C1-C2-O2


There are no ring outliers.

3 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	BBB	505	EDO	1	0
4	BBB	504	5QP	7	0
4	AAA	503	5QP	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	AAA	476/499 (95%)	-0.58	3 (0%)	89 92	11, 15, 26, 53	1 (0%)
1	BBB	476/499 (95%)	-0.54	2 (0%)	92 94	11, 18, 31, 62	3 (0%)
All	All	952/998 (95%)	-0.56	5 (0%)	91 93	11, 16, 30, 62	4 (0%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	BBB	476	SER	3.0
1	AAA	476	SER	2.9
1	AAA	128	ALA	2.7
1	BBB	470	ALA	2.2
1	AAA	447	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	EDO	BBB	505	4/4	0.81	0.18	29,36,39,39	1
2	NAG	BBB	501	14/15	0.89	0.14	25,33,45,51	3
4	5QP	BBB	504	23/23	0.89	0.13	12,17,20,21	46
2	NAG	AAA	501	14/15	0.90	0.10	24,29,33,34	3
3	EDO	BBB	502	4/4	0.91	0.09	22,24,26,26	1
4	5QP	AAA	503	23/23	0.92	0.12	11,15,18,18	46
3	EDO	BBB	503	4/4	0.97	0.07	18,21,22,22	1
3	EDO	AAA	502	4/4	0.97	0.10	20,25,27,27	1
5	CA	BBB	506	1/1	1.00	0.08	14,14,14,14	0
5	CA	AAA	504	1/1	1.00	0.07	12,12,12,12	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 5QP BBB 504:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 5QP AAA 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.