



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 10, 2022 – 11:21 PM EST

PDB ID : 1YTV
Title : Maltose-binding protein fusion to a C-terminal fragment of the V1a vasopressin receptor
Authors : Adikesavan, N.V.; Mahmood, S.S.; Stanley, S.; Xu, Z.; Wu, N.; Thibonnier, M.; Shoham, M.
Deposited on : 2005-02-11
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.31.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.31.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	366	<div> <div>2%</div> <div>86%</div> <div>14%</div> </div>
1	B	366	<div> <div>3%</div> <div>80%</div> <div>19%</div> </div>
2	M	84	<div> <div>2%</div> <div>7%</div> <div>93%</div> </div>
2	N	84	<div> <div>4%</div> <div>6%</div> <div>94%</div> </div>
3	C	2	<div> <div>100%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	D	2	 100%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6445 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Maltose-binding periplasmic protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	366	Total	C	N	O	S	0	0	0
			2842	1831	461	544	6			
1	B	366	Total	C	N	O	S	0	0	0
			2842	1831	461	544	6			

- Molecule 2 is a protein called Vasopressin V1a receptor.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	M	6	Total	C	N	O	0	0	0
			42	21	9	12			
2	N	5	Total	C	N	O	0	0	0
			34	17	7	10			

There are 50 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
M	381	ASN	-	linker	UNP P37288
M	382	SER	-	linker	UNP P37288
M	383	SER	-	linker	UNP P37288
M	384	SER	-	linker	UNP P37288
M	385	ASN	-	linker	UNP P37288
M	386	ASN	-	linker	UNP P37288
M	387	ASN	-	linker	UNP P37288
M	388	ASN	-	linker	UNP P37288
M	389	ASN	-	linker	UNP P37288
M	390	ASN	-	linker	UNP P37288
M	391	ASN	-	linker	UNP P37288
M	392	ASN	-	linker	UNP P37288
M	393	ASN	-	linker	UNP P37288
M	394	ASN	-	linker	UNP P37288
M	395	LEU	-	linker	UNP P37288
M	396	GLY	-	linker	UNP P37288

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
M	397	ILE	-	linker	UNP P37288
M	398	GLU	-	linker	UNP P37288
M	399	GLU	-	linker	UNP P37288
M	400	ASN	-	linker	UNP P37288
M	401	LEU	-	linker	UNP P37288
M	402	TYR	-	linker	UNP P37288
M	403	PHE	-	linker	UNP P37288
M	404	GLN	-	linker	UNP P37288
M	405	GLY	-	linker	UNP P37288
N	381	ASN	-	linker	UNP P37288
N	382	SER	-	linker	UNP P37288
N	383	SER	-	linker	UNP P37288
N	384	SER	-	linker	UNP P37288
N	385	ASN	-	linker	UNP P37288
N	386	ASN	-	linker	UNP P37288
N	387	ASN	-	linker	UNP P37288
N	388	ASN	-	linker	UNP P37288
N	389	ASN	-	linker	UNP P37288
N	390	ASN	-	linker	UNP P37288
N	391	ASN	-	linker	UNP P37288
N	392	ASN	-	linker	UNP P37288
N	393	ASN	-	linker	UNP P37288
N	394	ASN	-	linker	UNP P37288
N	395	LEU	-	linker	UNP P37288
N	396	GLY	-	linker	UNP P37288
N	397	ILE	-	linker	UNP P37288
N	398	GLU	-	linker	UNP P37288
N	399	GLU	-	linker	UNP P37288
N	400	ASN	-	linker	UNP P37288
N	401	LEU	-	linker	UNP P37288
N	402	TYR	-	linker	UNP P37288
N	403	PHE	-	linker	UNP P37288
N	404	GLN	-	linker	UNP P37288
N	405	GLY	-	linker	UNP P37288

- Molecule 3 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
3	C	2	Total	C	O	0	0	0
			22	12	10			
3	D	2	Total	C	O	0	0	0
			22	12	10			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	337	Total	O	0	0
			337	337		
4	M	6	Total	O	0	0
			6	6		
4	B	296	Total	O	0	0
			296	296		
4	N	2	Total	O	0	0
			2	2		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Maltose-binding periplasmic protein



• Molecule 1: Maltose-binding periplasmic protein




• Molecule 2: Vasopressin V1a receptor




• Molecule 2: Vasopressin V1a receptor





- Molecule 3: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain C:  100%

GLC1
GLC2

- Molecule 3: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain D:  100%

GLC1
GLC2

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	51.10Å 66.56Å 115.72Å 90.00° 95.99° 90.00°	Depositor
Resolution (Å)	34.43 – 1.80 34.43 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.5 (34.43-1.80) 99.6 (34.43-1.80)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.03 (at 1.81Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.199 , 0.225 0.208 , 0.232	Depositor DCC
R_{free} test set	3609 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	18.7	Xtriage
Anisotropy	0.532	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 56.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6445	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.32	0/2911	0.54	0/3952
1	B	0.32	0/2911	0.53	0/3952
2	M	0.30	0/41	0.45	0/54
2	N	0.44	0/33	0.74	0/43
All	All	0.32	0/5896	0.54	0/8001

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2842	0	2812	39	0
1	B	2842	0	2812	59	0
2	M	42	0	32	0	0
2	N	34	0	26	0	0
3	C	22	0	19	0	0
3	D	22	0	19	0	0
4	A	337	0	0	8	0
4	B	296	0	0	10	0
4	M	6	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	N	2	0	0	0	0
All	All	6445	0	5720	95	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (95) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:93:ILE:HG22	1:A:291:LYS:HE3	1.40	1.02
1:B:16:ILE:H	1:B:16:ILE:HD12	1.35	0.89
1:A:314:GLY:HA3	4:A:892:HOH:O	1.79	0.81
1:A:136:LEU:HD21	1:A:140:PRO:HD3	1.68	0.76
1:A:345:PRO:HG2	1:A:350:MET:SD	2.26	0.76
1:B:93:ILE:HG22	1:B:95:PRO:HD3	1.69	0.75
1:B:82:GLY:HA3	1:B:346:ASN:O	1.88	0.73
1:B:19:GLY:H	1:B:286:ASN:HD21	1.36	0.72
1:B:345:PRO:HG2	1:B:350:MET:SD	2.29	0.71
1:B:136:LEU:HD21	1:B:140:PRO:HD3	1.71	0.71
1:A:19:GLY:H	1:A:286:ASN:HD21	1.39	0.70
1:A:192:ILE:HG23	1:A:349:GLN:HG2	1.73	0.70
1:B:343:ILE:HB	4:B:791:HOH:O	1.90	0.69
1:B:314:GLY:HA3	4:B:868:HOH:O	1.93	0.68
1:A:192:ILE:HD11	1:A:347:ILE:HD12	1.75	0.68
1:B:19:GLY:H	1:B:286:ASN:ND2	1.91	0.68
1:A:292:GLU:OE1	1:B:98:ALA:HA	1.96	0.65
1:A:292:GLU:OE1	1:B:101:ASP:HB3	1.97	0.63
1:A:308:ASN:ND2	1:A:313:LEU:H	1.98	0.62
1:B:349:GLN:NE2	1:B:349:GLN:H	1.97	0.62
1:B:191:ASP:OD1	1:B:193:LYS:HG2	2.00	0.61
1:A:82:GLY:HA3	1:A:346:ASN:O	2.02	0.60
1:B:349:GLN:H	1:B:349:GLN:HE21	1.52	0.57
1:B:147:PRO:HG2	4:B:609:HOH:O	2.04	0.56
1:A:192:ILE:CD1	1:A:347:ILE:HD12	2.35	0.56
1:B:16:ILE:H	1:B:16:ILE:CD1	2.11	0.56
1:A:147:PRO:HG2	4:A:686:HOH:O	2.05	0.56
1:A:19:GLY:H	1:A:286:ASN:ND2	2.04	0.55
4:A:979:HOH:O	1:B:323:GLU:HG2	2.07	0.55
1:B:146:ILE:N	1:B:147:PRO:HD2	2.22	0.55
1:A:280:ILE:CD1	1:A:291:LYS:HD3	2.38	0.54
1:B:60:LYS:HE2	4:B:972:HOH:O	2.08	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:120:TYR:CE1	1:B:295:GLU:HG3	2.43	0.54
1:B:192:ILE:HD12	4:B:754:HOH:O	2.08	0.54
1:B:366:SER:OG	1:B:368:ARG:HG3	2.08	0.53
1:A:61:PHE:HB3	1:A:62:PRO:HD3	1.90	0.53
1:B:58:GLU:HG2	1:B:59:GLU:HG3	1.91	0.52
1:A:291:LYS:HE2	4:A:1056:HOH:O	2.09	0.52
1:A:190:TYR:CZ	1:A:345:PRO:HB3	2.44	0.52
1:B:210:VAL:HG12	1:B:214:LYS:HE3	1.91	0.52
1:A:146:ILE:N	1:A:147:PRO:HD2	2.26	0.51
1:B:363:ASN:HB3	1:B:369:GLN:HB2	1.93	0.49
1:A:195:VAL:HG12	1:A:197:VAL:H	1.78	0.49
1:A:292:GLU:OE1	1:B:101:ASP:CB	2.61	0.48
1:A:308:ASN:HD22	1:A:313:LEU:H	1.58	0.48
1:A:253:LYS:HE3	4:A:1136:HOH:O	2.12	0.48
1:A:268:PRO:HB3	4:A:984:HOH:O	2.13	0.48
1:A:223:ASP:OD1	1:A:226:ILE:HG12	2.13	0.48
1:A:219:ASN:HB2	4:A:842:HOH:O	2.12	0.48
1:B:19:GLY:N	1:B:286:ASN:HD21	2.06	0.48
1:B:232:ASN:HD22	1:B:232:ASN:N	2.11	0.48
1:A:103:LEU:HD13	1:A:121:PRO:HG2	1.96	0.47
1:B:335:MET:O	1:B:339:GLN:HG3	2.13	0.47
1:A:190:TYR:CE2	1:A:345:PRO:HB3	2.48	0.47
1:A:287:LYS:O	1:A:291:LYS:HG2	2.15	0.47
1:A:198:ASP:O	1:A:203:LYS:HE3	2.15	0.47
1:B:103:LEU:CD1	1:B:121:PRO:HG2	2.45	0.47
1:B:57:LEU:C	1:B:57:LEU:HD12	2.35	0.47
1:B:40:LYS:HD3	1:B:43:LYS:HE2	1.96	0.46
1:B:136:LEU:HD22	1:B:149:LEU:HD21	1.97	0.46
1:B:103:LEU:HD13	1:B:121:PRO:HG2	1.96	0.46
1:B:332:ALA:O	1:B:336:GLU:HG3	2.15	0.46
1:B:243:PRO:HA	1:B:246:TRP:CE2	2.51	0.46
1:A:62:PRO:HG3	1:A:84:TYR:CE2	2.50	0.46
1:B:272:PHE:CG	1:B:344:MET:HG2	2.51	0.46
1:A:335:MET:O	1:A:339:GLN:HG3	2.15	0.45
1:B:136:LEU:CD2	1:B:140:PRO:HD3	2.41	0.45
1:B:210:VAL:CG1	1:B:214:LYS:HE3	2.47	0.44
1:B:131:TYR:CE2	1:B:139:PRO:HD3	2.53	0.44
1:A:234:GLY:HA2	4:A:520:HOH:O	2.17	0.44
1:B:139:PRO:HA	1:B:140:PRO:HD3	1.89	0.44
1:B:291:LYS:O	1:B:295:GLU:HB2	2.18	0.43
1:B:321:TYR:CE2	1:B:325:LEU:HD11	2.53	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:328:ASP:HB3	1:B:331:ILE:HG12	2.00	0.43
1:B:46:GLY:HA2	4:B:798:HOH:O	2.19	0.43
1:A:288:GLU:O	1:A:292:GLU:HG3	2.18	0.43
1:B:150:ASP:HA	1:B:160:ALA:HB2	2.01	0.43
1:B:61:PHE:N	1:B:62:PRO:HD2	2.33	0.43
1:A:104:TYR:HA	1:A:105:PRO:HD3	1.94	0.42
1:B:342:GLU:HG2	4:B:650:HOH:O	2.20	0.42
1:A:57:LEU:HD12	1:A:57:LEU:C	2.40	0.42
1:B:190:TYR:CZ	1:B:345:PRO:HB3	2.55	0.42
1:B:268:PRO:HB3	4:B:803:HOH:O	2.20	0.42
1:A:217:HIS:O	1:A:218:MET:HG3	2.20	0.42
1:B:143:TRP:NE1	1:B:262:PRO:HG2	2.35	0.42
1:B:368:ARG:HG2	1:B:368:ARG:HH11	1.85	0.42
1:A:232:ASN:N	1:A:232:ASN:HD22	2.18	0.41
1:B:103:LEU:HD23	1:B:318:LEU:HA	2.03	0.41
1:B:190:TYR:CE2	1:B:345:PRO:HB3	2.55	0.41
1:B:234:GLY:HA2	4:B:571:HOH:O	2.20	0.41
1:B:302:GLU:HG3	4:B:643:HOH:O	2.20	0.41
1:B:217:HIS:O	1:B:218:MET:HG3	2.21	0.41
1:A:243:PRO:HA	1:A:246:TRP:CE2	2.57	0.40
1:B:223:ASP:OD1	1:B:226:ILE:HG12	2.20	0.40
1:B:103:LEU:CD2	1:B:318:LEU:HA	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	364/366 (100%)	360 (99%)	4 (1%)	0	100	100
1	B	364/366 (100%)	357 (98%)	7 (2%)	0	100	100
2	M	4/84 (5%)	2 (50%)	2 (50%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	N	3/84 (4%)	3 (100%)	0	0	100	100
All	All	735/900 (82%)	722 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	293/293 (100%)	290 (99%)	3 (1%)	76	71
1	B	293/293 (100%)	287 (98%)	6 (2%)	55	44
2	M	6/80 (8%)	6 (100%)	0	100	100
2	N	5/80 (6%)	5 (100%)	0	100	100
All	All	597/746 (80%)	588 (98%)	9 (2%)	65	56

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	174	LEU
1	A	272	PHE
1	A	323	GLU
1	B	174	LEU
1	B	192	ILE
1	B	216	LYS
1	B	272	PHE
1	B	323	GLU
1	B	349	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	32	ASN
1	A	63	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	114	ASN
1	A	232	ASN
1	A	248	ASN
1	A	286	ASN
1	A	296	ASN
1	A	308	ASN
1	A	339	GLN
1	A	349	GLN
1	B	32	ASN
1	B	232	ASN
1	B	248	ASN
1	B	286	ASN
1	B	296	ASN
1	B	339	GLN
1	B	349	GLN
2	N	381	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GLC	C	1	3	11,11,12	1.25	1 (9%)	15,15,17	2.03	4 (26%)
3	GLC	C	2	3	11,11,12	0.92	0	15,15,17	1.48	2 (13%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GLC	D	1	3	11,11,12	1.26	1 (9%)	15,15,17	2.03	4 (26%)
3	GLC	D	2	3	11,11,12	1.04	0	15,15,17	1.25	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GLC	C	1	3	-	0/2/19/22	0/1/1/1
3	GLC	C	2	3	-	0/2/19/22	0/1/1/1
3	GLC	D	1	3	-	0/2/19/22	0/1/1/1
3	GLC	D	2	3	-	0/2/19/22	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	1	GLC	O5-C5	2.39	1.48	1.43
3	D	1	GLC	O5-C5	2.35	1.48	1.43

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	2	GLC	C1-O5-C5	3.96	117.56	112.19
3	C	1	GLC	C2-C3-C4	-3.88	104.19	110.89
3	D	1	GLC	O2-C2-C3	-3.79	102.54	110.14
3	D	1	GLC	C2-C3-C4	-3.73	104.44	110.89
3	C	1	GLC	O2-C2-C3	-3.71	102.71	110.14
3	D	1	GLC	C1-C2-C3	3.37	113.81	109.67
3	C	1	GLC	C1-C2-C3	3.22	113.62	109.67
3	D	2	GLC	C1-O5-C5	3.21	116.54	112.19
3	D	1	GLC	O6-C6-C5	-2.96	101.12	111.29
3	C	2	GLC	C2-C3-C4	-2.86	105.94	110.89
3	C	1	GLC	O6-C6-C5	-2.67	102.14	111.29
3	D	2	GLC	C2-C3-C4	-2.44	106.68	110.89


There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	366/366 (100%)	0.05	9 (2%) 57 52	11, 19, 35, 50	0
1	B	366/366 (100%)	0.10	11 (3%) 50 44	11, 19, 36, 56	0
2	M	6/84 (7%)	2.53	2 (33%) 0 0	22, 36, 63, 68	0
2	N	5/84 (5%)	2.79	3 (60%) 0 0	26, 27, 50, 54	0
All	All	743/900 (82%)	0.12	25 (3%) 45 39	11, 19, 36, 68	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	15	LYS	10.7
1	B	16	ILE	9.4
2	N	385	ASN	7.6
2	M	386	ASN	6.4
1	B	68	GLY	6.4
2	M	385	ASN	5.7
1	A	66	ALA	4.0
1	B	69	ASP	3.6
2	N	384	SER	3.3
1	A	187	ASN	2.9
1	A	15	LYS	2.8
1	B	187	ASN	2.7
1	A	67	THR	2.7
1	A	186	GLU	2.7
1	B	186	GLU	2.6
1	A	219	ASN	2.5
1	A	216	LYS	2.4
1	B	193	LYS	2.3
1	A	59	GLU	2.2
1	A	188	GLY	2.2
1	B	327	LYS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	152	GLU	2.1
1	B	149	LEU	2.1
2	N	383	SER	2.1
1	B	188	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	GLC	D	1	11/12	0.91	0.11	12,14,17,17	0
3	GLC	C	1	11/12	0.92	0.12	12,15,18,22	0
3	GLC	C	2	11/12	0.97	0.07	10,11,13,14	0
3	GLC	D	2	11/12	0.98	0.08	8,10,13,14	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.