



Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 10:31 am BST

PDB ID : 1YTW
Title : YERSINIA PTPASE COMPLEXED WITH TUNGSTATE
Authors : Fauman, E.B.; Schubert, H.L.; Saper, M.A.
Deposited on : 1996-05-01
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	306	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2328 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called YERSINIA PROTEIN TYROSINE PHOSPHATASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	283	2179	1331	404	427	17	0	1	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	235	ARG	CYS	CONFLICT	UNP P15273

- Molecule 2 is TUNGSTATE(VI)ION (three-letter code: WO4) (formula: O₄W).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	W		
2	A	1	5	4	1	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	139	Total	O	0	0
			139	139		

4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	56.30 Å 49.80 Å 100.60 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	7.00 – 2.40	Depositor
% Data completeness (in resolution range)	98.0 (7.00-2.40)	Depositor
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.179 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2328	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: WO4, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.89	2/2206 (0.1%)	0.99	3/2980 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	221	CYS	CB-SG	6.83	1.93	1.82
1	A	314	GLU	CB-CG	-5.66	1.41	1.52

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	228	ARG	NE-CZ-NH1	7.08	123.84	120.30
1	A	228	ARG	NE-CZ-NH2	-6.83	116.89	120.30
1	A	222	GLY	N-CA-C	6.66	129.75	113.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2179	0	2180	42	0
2	A	5	0	0	1	0
3	A	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	139	0	0	1	0
All	All	2328	0	2180	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:429:VAL:HA	1:A:432:MET:HE3	1.51	0.93
1:A:251:VAL:HG12	1:A:427:LEU:HD21	1.54	0.90
1:A:189:TYR:O	1:A:194:ARG:HD2	1.91	0.71
1:A:357:GLN:OE1	1:A:447:LYS:HD3	1.92	0.69
1:A:228:ARG:NH2	1:A:297:GLY:O	2.28	0.64
1:A:363:GLU:HA	1:A:366:LYS:HG2	1.80	0.64
1:A:251:VAL:HB	1:A:418[B]:CYS:SG	2.40	0.62
1:A:337:ARG:HG3	1:A:337:ARG:HH11	1.65	0.60
1:A:404:ARG:HD3	4:A:664:HOH:O	2.00	0.60
1:A:380:ARG:HG3	1:A:397:LEU:HD22	1.84	0.57
1:A:250:GLN:HA	1:A:254:THR:O	2.05	0.56
1:A:455:ILE:HG23	1:A:465:LEU:HD21	1.87	0.56
1:A:205:ARG:HD3	1:A:442:GLY:O	2.06	0.55
1:A:251:VAL:CG1	1:A:432:MET:HA	2.36	0.55
1:A:251:VAL:HG11	1:A:432:MET:HG3	1.88	0.55
1:A:363:GLU:OE2	1:A:366:LYS:HD2	2.06	0.55
1:A:337:ARG:NH1	1:A:337:ARG:HG3	2.23	0.54
1:A:337:ARG:HG2	1:A:343:THR:HG23	1.90	0.53
1:A:251:VAL:HG11	1:A:432:MET:HA	1.93	0.51
1:A:251:VAL:HA	1:A:435:GLN:OE1	2.11	0.50
1:A:292:ALA:O	1:A:294:GLN:NE2	2.45	0.49
1:A:355:PRO:O	1:A:358:THR:OG1	2.30	0.47
1:A:313:VAL:CG1	1:A:334:LEU:HD22	2.45	0.47
1:A:428:SER:OG	1:A:467:ASN:ND2	2.48	0.46
1:A:192:GLU:O	1:A:195:ALA:HB3	2.17	0.45
1:A:404:ARG:HB3	2:A:1:WO4:O2	2.17	0.45
1:A:236:ARG:O	1:A:236:ARG:HD3	2.16	0.45
1:A:318:THR:C	1:A:319:GLN:HG3	2.36	0.45
1:A:288:SER:OG	1:A:303:ARG:NH2	2.49	0.43
1:A:187:SER:HA	1:A:188:PRO:HD3	1.82	0.42
1:A:380:ARG:NH2	1:A:397:LEU:O	2.52	0.42
1:A:318:THR:O	1:A:319:GLN:HG3	2.19	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:279:THR:HA	1:A:280:PRO:HD2	1.96	0.41
1:A:236:ARG:HH11	1:A:239:ALA:HB3	1.86	0.41
1:A:388:SER:HB3	1:A:391:VAL:HG23	2.01	0.41
1:A:251:VAL:HG11	1:A:432:MET:CG	2.49	0.41
1:A:205:ARG:HD2	1:A:443:ILE:HD13	2.02	0.41
1:A:228:ARG:NH1	1:A:261:TYR:O	2.47	0.41
1:A:366:LYS:HG3	1:A:367:ALA:N	2.36	0.41
1:A:259:CYS:O	1:A:402:HIS:HB2	2.21	0.40
1:A:211:ALA:O	1:A:214:ASP:HB2	2.20	0.40
1:A:300:ASP:OD1	1:A:303:ARG:HD2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	282/306 (92%)	266 (94%)	15 (5%)	1 (0%)	34	48

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	445	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	240/255 (94%)	228 (95%)	12 (5%)	24	40

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	205	ARG
1	A	230	ARG
1	A	236	ARG
1	A	304	GLN
1	A	310	SER
1	A	325	ASP
1	A	362	SER
1	A	377	GLU
1	A	380	ARG
1	A	397	LEU
1	A	420	ASN
1	A	448	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	253	ASN
1	A	264	GLN
1	A	294	GLN
1	A	304	GLN
1	A	319	GLN
1	A	467	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	A	3	-	4,4,4	0.62	0	6,6,6	0.34	0
2	WO4	A	1	-	2,4,4	2.26	1 (50%)	-		

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1	WO4	W-O2	2.52	1.80	1.74

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1	WO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.