



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 30, 2021 – 05:11 am GMT

PDB ID : 6YTR  
Title : Structure of recombinant human beta-glucocerebrosidase in complex with cyclophellitol aziridine inhibitor  
Authors : Rowland, R.J.; Davies, G.J.  
Deposited on : 2020-04-24  
Resolution : 1.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4 (270009), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.23.2  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0267  
CCP4 : 7.1.010 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	497	 91% 9%
1	BBB	497	 91% 9%
2	AdA	4	 100%
3	BdB	3	 100%
4	BuB	2	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	SO4	AAA	521	-	-	X	-
8	EDO	BBB	2727	-	-	X	-

## 2 Entry composition [i](#)

There are 10 unique types of molecules in this entry. The entry contains 17369 atoms, of which 8197 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lysosomal acid glucosylceramidase.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	AAA	497	Total	C	H	N	O	S	208	8	0
			7906	2567	3915	684	724	16			
1	BBB	497	Total	C	H	N	O	S	205	3	0
			7808	2537	3868	672	715	16			

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	AdA	4	Total	C	H	N	O		11	0	0
			98	28	48	2	20				

- Molecule 3 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	BdB	3	Total	C	H	N	O		8	0	0
			76	22	37	2	15				

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	BuB	2	Total	C	H	N	O	5	0	0
			55	16	27	2	10			

- Molecule 5 is (1 {R},2 {S},3 {S},4 {S},5 {R},6 {R})-5-azanyl-6-(hydroxymethyl)cyclohexane-1,2,3,4-tetrol (three-letter code: PO8) (formula: C<sub>7</sub>H<sub>16</sub>NO<sub>5</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	AAA	1	Total	C	H	N	O	3	0
			27	7	15	1	4		
5	BBB	1	Total	C	H	N	O	3	0
			27	7	15	1	4		

- Molecule 6 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	AAA	1	Total	O	S	0	0
			5	4	1		
6	AAA	1	Total	O	S	0	0
			5	4	1		
6	AAA	1	Total	O	S	0	0
			5	4	1		
6	AAA	1	Total	O	S	0	0
			5	4	1		
6	BBB	1	Total	O	S	0	0
			5	4	1		
6	BBB	1	Total	O	S	0	0
			5	4	1		
6	BBB	1	Total	O	S	0	0
			5	4	1		

- Molecule 7 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	AAA	1	Total	C	H	N	O	3	0
			28	8	14	1	5		

- Molecule 8 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	AAA	1	Total	C	H	O	1	0
			10	2	6	2		
8	AAA	1	Total	C	H	O	1	0
			10	2	6	2		
8	AAA	1	Total	C	H	O	1	0
			10	2	6	2		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	AAA	1	Total 10	C 2	H 6	O 2	1	0
8	AAA	1	Total 10	C 2	H 6	O 2	1	0
8	AAA	1	Total 10	C 2	H 6	O 2	1	0
8	AAA	1	Total 10	C 2	H 6	O 2	1	0
8	AAA	1	Total 10	C 2	H 6	O 2	1	0
8	AAA	1	Total 10	C 2	H 6	O 2	1	0
8	AAA	1	Total 10	C 2	H 6	O 2	1	0
8	AAA	1	Total 10	C 2	H 6	O 2	1	0
8	AAA	1	Total 10	C 2	H 6	O 2	1	0
8	AAA	1	Total 10	C 2	H 6	O 2	1	0
8	AAA	1	Total 10	C 2	H 6	O 2	1	0
8	AAA	1	Total 10	C 2	H 6	O 2	1	0
8	AAA	1	Total 10	C 2	H 6	O 2	1	0
8	AAA	1	Total 10	C 2	H 6	O 2	1	0
8	AAA	1	Total 10	C 2	H 6	O 2	1	0
8	AAA	1	Total 10	C 2	H 6	O 2	1	0
8	AAA	1	Total 10	C 2	H 6	O 2	1	0
8	AAA	1	Total 10	C 2	H 6	O 2	1	0
8	AAA	1	Total 10	C 2	H 6	O 2	1	0
8	BBB	1	Total 10	C 2	H 6	O 2	1	0
8	BBB	1	Total 10	C 2	H 6	O 2	1	0
8	BBB	1	Total 10	C 2	H 6	O 2	1	0
8	BBB	1	Total 10	C 2	H 6	O 2	1	0
8	BBB	1	Total 20	C 4	H 12	O 4	2	1

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	BBB	1	Total	C	H	O	1	0
			10	2	6	2		
8	BBB	1	Total	C	H	O	1	0
			10	2	6	2		
8	BBB	1	Total	C	H	O	1	0
			10	2	6	2		
8	BBB	1	Total	C	H	O	1	0
			10	2	6	2		
8	BBB	1	Total	C	H	O	1	0
			10	2	6	2		
8	BBB	1	Total	C	H	O	1	0
			10	2	6	2		
8	BBB	1	Total	C	H	O	1	0
			10	2	6	2		
8	BBB	1	Total	C	H	O	1	0
			10	2	6	2		
8	BBB	1	Total	C	H	O	1	0
			10	2	6	2		
8	BBB	1	Total	C	H	O	1	0
			10	2	6	2		
8	BBB	1	Total	C	H	O	1	0
			10	2	6	2		
8	BBB	1	Total	C	H	O	1	0
			10	2	6	2		
8	BBB	1	Total	C	H	O	1	0
			10	2	6	2		
8	BBB	1	Total	C	H	O	1	0
			10	2	6	2		
8	BBB	1	Total	C	H	O	1	0
			10	2	6	2		
8	BBB	1	Total	C	H	O	1	0
			10	2	6	2		
8	BBB	1	Total	C	H	O	1	0
			10	2	6	2		

- Molecule 9 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	AAA	1	Total	Na	0	0
			1	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	BBB	1	Total	Na	0	0
			1	1		


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	AAA	468	Total	O	0	0
			468	468		
10	BBB	409	Total	O	0	0
			409	409		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Lysosomal acid glucosylceramidase



- Molecule 1: Lysosomal acid glucosylceramidase



- Molecule 2: alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain BuB:

100%

3AG1  
3AG2

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	53.16Å 156.78Å 68.23Å 90.00° 102.12° 90.00°	Depositor
Resolution (Å)	66.80 – 1.70 66.71 – 1.70	Depositor EDS
% Data completeness (in resolution range)	99.9 (66.80-1.70) 99.9 (66.71-1.70)	Depositor EDS
$R_{merge}$	0.16	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.75 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, $R_{free}$	0.159 , 0.203 0.170 , 0.213	Depositor DCC
$R_{free}$ test set	5927 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	23.0	Xtriage
Anisotropy	0.050	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	(Not available) , (Not available)	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	17369	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.88% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO8, BMA, NAG, MAN, SO4, EDO, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	AAA	0.73	1/4111 (0.0%)	0.82	0/5607
1	BBB	0.74	1/4059 (0.0%)	0.81	1/5538 (0.0%)
All	All	0.74	2/8170 (0.0%)	0.81	1/11145 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	BBB	340	GLU	CD-OE2	14.93	1.42	1.25
1	AAA	340	GLU	CD-OE2	13.77	1.40	1.25

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	BBB	285	ARG	NE-CZ-NH1	5.06	122.83	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	3991	3915	3882	32	0
1	BBB	3940	3868	3839	32	1
2	AdA	50	48	43	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	BdB	39	37	34	0	0
4	BuB	28	27	25	0	0
5	AAA	12	15	0	0	0
5	BBB	12	15	0	0	0
6	AAA	20	0	0	3	0
6	BBB	15	0	0	0	0
7	AAA	14	14	13	0	0
8	AAA	76	114	114	11	0
8	BBB	96	144	144	13	0
9	AAA	1	0	0	0	0
9	BBB	1	0	0	0	0
10	AAA	468	0	0	10	0
10	BBB	409	0	0	4	0
All	All	9172	8197	8094	67	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 67 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:BBB:349:GLU:HA	8:BBB:2714:EDO:H12	1.63	0.79
1:BBB:56:ILE:HG21	1:BBB:477:VAL:HG12	1.69	0.74
1:BBB:237:SER:OG	8:BBB:2726:EDO:H11	1.88	0.73
1:AAA:165:LEU:HD22	1:AAA:172:VAL:HB	1.71	0.73
1:BBB:211:ARG:NH2	10:BBB:2801:HOH:O	2.20	0.71

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:BBB:63:THR:H	1:BBB:200:GLN:HE21[1_554]	1.17	0.43

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	503/497 (101%)	486 (97%)	14 (3%)	3 (1%)	25	11
1	BBB	498/497 (100%)	475 (95%)	21 (4%)	2 (0%)	34	18
All	All	1001/994 (101%)	961 (96%)	35 (4%)	5 (0%)	29	13

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	AAA	233	GLU
1	BBB	233	GLU
1	AAA	281	LEU
1	AAA	381	TRP
1	BBB	281	LEU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	428/424 (101%)	422 (99%)	6 (1%)	67	53
1	BBB	423/424 (100%)	419 (99%)	4 (1%)	78	70
All	All	851/848 (100%)	841 (99%)	10 (1%)	71	59

5 of 10 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	BBB	335	MET
1	BBB	420	LEU
1	BBB	483	ILE
1	AAA	420	LEU
1	AAA	470	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are



no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

9 monosaccharides are modelled in this entry.

There are no bond length outliers.

There are no bond angle outliers.



There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

Of 55 ligands modelled in this entry, 2 are monoatomic - leaving 53 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is

within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	AAA	497/497 (100%)	-0.56	0 <a href="#">100</a> <a href="#">100</a>	15, 22, 34, 48	1 (0%)
1	BBB	497/497 (100%)	-0.57	4 (0%) <a href="#">86</a> <a href="#">88</a>	18, 25, 39, 59	5 (1%)
All	All	994/994 (100%)	-0.56	4 (0%) <a href="#">92</a> <a href="#">93</a>	15, 24, 38, 59	6 (0%)

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	BBB	31	PHE	7.9
1	BBB	30	THR	3.8
1	BBB	1	ALA	2.8
1	BBB	29	PRO	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	BMA	BdB	3	11/12	0.67	0.23	55,66,71,72	4
4	NAG	BuB	1	14/15	0.70	0.18	62,70,75,86	2
4	NAG	BuB	2	14/15	0.70	0.20	69,77,80,82	3
2	MAN	AdA	4	11/12	0.78	0.15	62,72,79,81	4
2	BMA	AdA	3	11/12	0.82	0.10	58,63,65,67	3

*Continued on next page...*




*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NAG	AdA	2	14/15	0.93	0.07	31,36,42,48	2
3	NAG	BdB	2	14/15	0.96	0.08	30,32,35,40	2
2	NAG	AdA	1	14/15	0.96	0.07	25,29,31,33	2
3	NAG	BdB	1	14/15	0.97	0.08	28,31,40,44	2

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




#### Electron density around Chain AdA:

2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around Chain BdB:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
8	EDO	BBB	2723	4/4	0.57	0.17	60,65,68,69	1
8	EDO	AAA	511	4/4	0.66	0.25	48,55,59,59	1
7	NAG	AAA	504	14/15	0.71	0.16	57,64,70,74	3
8	EDO	BBB	2725	4/4	0.71	0.13	56,58,61,61	1
8	EDO	AAA	524	4/4	0.72	0.10	50,57,59,59	1
8	EDO	BBB	2715	4/4	0.75	0.16	59,62,62,64	1
8	EDO	AAA	525	4/4	0.76	0.20	52,56,59,59	1
8	EDO	BBB	2727	4/4	0.77	0.25	20,54,55,56	1
8	EDO	BBB	2713	4/4	0.79	0.12	49,52,62,63	1
8	EDO	AAA	507	4/4	0.81	0.16	56,60,61,61	1
8	EDO	BBB	2701	4/4	0.83	0.18	57,58,60,61	1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
8	EDO	BBB	2716	4/4	0.85	0.10	55,56,58,58	1
6	SO4	BBB	2721	5/5	0.85	0.22	55,69,80,82	0
8	EDO	BBB	2714	4/4	0.86	0.16	47,48,50,52	1
8	EDO	BBB	2710	4/4	0.86	0.14	45,48,53,54	1
8	EDO	AAA	513	4/4	0.86	0.13	53,55,59,62	1
8	EDO	AAA	523	4/4	0.87	0.17	50,53,58,59	1
8	EDO	BBB	2724	4/4	0.88	0.15	28,35,38,39	1
6	SO4	AAA	521	5/5	0.89	0.22	38,44,49,51	5
8	EDO	AAA	514	4/4	0.89	0.29	40,44,51,53	1
8	EDO	AAA	518	4/4	0.89	0.16	36,45,53,53	1
8	EDO	BBB	2719	4/4	0.89	0.21	56,56,58,58	1
8	EDO	AAA	515	4/4	0.90	0.11	32,36,41,41	1
8	EDO	BBB	2706	4/4	0.90	0.18	29,35,38,38	1
8	EDO	BBB	2707	4/4	0.90	0.14	47,53,55,56	1
8	EDO	BBB	2708[A]	4/4	0.90	0.19	24,26,29,30	10
8	EDO	BBB	2708[B]	4/4	0.90	0.19	25,27,30,31	10
8	EDO	AAA	512	4/4	0.90	0.16	35,42,48,48	1
8	EDO	BBB	2712	4/4	0.90	0.19	35,48,50,51	1
8	EDO	AAA	519	4/4	0.90	0.07	45,49,53,55	1
8	EDO	BBB	2728	4/4	0.90	0.21	20,46,49,53	1
8	EDO	AAA	508	4/4	0.91	0.14	45,51,52,52	1
8	EDO	BBB	2718	4/4	0.91	0.14	44,54,65,68	1
8	EDO	AAA	506	4/4	0.91	0.12	22,26,27,27	1
6	SO4	AAA	510	5/5	0.91	0.15	74,75,82,88	0
8	EDO	AAA	520	4/4	0.92	0.12	34,46,55,56	1
8	EDO	AAA	517	4/4	0.92	0.11	36,39,48,49	1
8	EDO	AAA	516	4/4	0.93	0.15	42,47,49,50	1
8	EDO	BBB	2726	4/4	0.93	0.12	20,43,47,48	1
8	EDO	BBB	2717	4/4	0.93	0.20	41,47,48,48	1
8	EDO	AAA	505	4/4	0.93	0.13	47,48,49,50	1
8	EDO	AAA	509	4/4	0.94	0.10	36,43,51,51	1
8	EDO	BBB	2705	4/4	0.94	0.18	47,51,52,54	1
8	EDO	AAA	526	4/4	0.94	0.10	20,46,47,52	1
8	EDO	BBB	2711	4/4	0.95	0.10	48,49,53,54	1
5	PO8	BBB	2702	12/13	0.96	0.07	20,24,27,28	3
5	PO8	AAA	501	12/13	0.96	0.07	18,21,26,26	3
8	EDO	BBB	2720	4/4	0.96	0.17	41,48,55,57	1
8	EDO	BBB	2709	4/4	0.97	0.24	32,34,38,39	1
9	NA	BBB	2722	1/1	0.97	0.09	34,34,34,34	0
6	SO4	BBB	2703	5/5	0.99	0.07	27,31,36,36	0
6	SO4	BBB	2704	5/5	0.99	0.06	40,41,44,44	0
6	SO4	AAA	502	5/5	0.99	0.07	26,27,30,31	0

*Continued on next page...*




*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
9	NA	AAA	522	1/1	0.99	0.06	34,34,34,34	0
6	SO4	AAA	503	5/5	0.99	0.05	33,34,35,35	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around PO8 BBB 2702:**

2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around PO8 AAA 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.