



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 2, 2021 – 01:02 pm GMT

PDB ID : 6YVP  
Title : Human histidine triad nucleotide-binding protein 2 (hHINT2) complexed with dGMP and refined to 2.77 Å  
Authors : Dolot, R.D.; Krakowiak, A.; Nawrot, B.C.  
Deposited on : 2020-04-28  
Resolution : 2.77 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4 (270009), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.23.2  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0267  
CCP4 : 7.1.010 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4107 (2.80-2.76)
Clashscore	141614	4575 (2.80-2.76)
Ramachandran outliers	138981	4487 (2.80-2.76)
Sidechain outliers	138945	4489 (2.80-2.76)
RSRZ outliers	127900	4027 (2.80-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	163	
1	BBB	163	

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 1929 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histidine triad nucleotide-binding protein 2, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	AAA	112	Total	C	N	O	S	0	0	0
			878	565	155	157	1			
1	BBB	112	Total	C	N	O	S	0	1	0
			887	571	157	158	1			

- Molecule 2 is 2'-DEOXYGUANOSINE-5'-MONOPHOSPHATE (three-letter code: DGP) (formula:  $C_{10}H_{14}N_5O_7P$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	AAA	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
2	BBB	1	Total	C	N	O	P	0	0
			23	10	5	7	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	AAA	57	Total 57	O 57	0	0
3	BBB	61	Total 61	O 61	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Histidine triad nucleotide-binding protein 2, mitochondrial



- Molecule 1: Histidine triad nucleotide-binding protein 2, mitochondrial



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	71.65Å 36.53Å 86.89Å 90.00° 103.03° 90.00°	Depositor
Resolution (Å)	42.32 – 2.77 42.32 – 2.77	Depositor EDS
% Data completeness (in resolution range)	98.6 (42.32-2.77) 98.7 (42.32-2.77)	Depositor EDS
$R_{merge}$	0.20	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.13 (at 2.77Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, $R_{free}$	0.208 , 0.308 0.214 , 0.313	Depositor DCC
$R_{free}$ test set	565 reflections (9.92%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	40.0	Xtriage
Anisotropy	0.036	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	(Not available) , (Not available)	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	1929	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 15.45% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DGP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	AAA	0.64	0/899	0.78	0/1221
1	BBB	0.63	0/908	0.82	0/1232
All	All	0.63	0/1807	0.80	0/2453

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	878	0	895	18	0
1	BBB	887	0	907	17	0
2	AAA	23	0	12	0	0
2	BBB	23	0	12	3	0
3	AAA	57	0	0	2	0
3	BBB	61	0	0	0	0
All	All	1929	0	1826	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:BBB:144:SER:OG	2:BBB:201:DGP:OP1	1.73	1.06
1:AAA:52:ALA:HB3	1:AAA:53:PRO:HD2	1.47	0.95
1:AAA:52:ALA:HB3	1:AAA:53:PRO:CD	2.12	0.80
1:AAA:52:ALA:CB	1:AAA:53:PRO:CD	2.64	0.76
1:AAA:93:PRO:HD2	1:AAA:148:LEU:O	1.93	0.68
1:AAA:77:VAL:HG11	1:AAA:121:THR:HG21	1.84	0.60
1:BBB:93:PRO:HD2	1:BBB:148:LEU:O	2.02	0.60
1:BBB:57:SER:HB2	1:BBB:58:ARG:HH21	1.68	0.57
1:BBB:127:LEU:CD2	1:BBB:154:GLY:HA3	2.36	0.56
1:BBB:54:THR:H	1:BBB:57:SER:HG	1.53	0.56
1:BBB:127:LEU:HD22	1:BBB:154:GLY:HA3	1.89	0.55
1:BBB:58:ARG:HB3	1:BBB:64:LEU:HB2	1.87	0.55
1:AAA:52:ALA:HB1	3:AAA:335:HOH:O	2.06	0.55
1:AAA:99:ARG:HB3	1:AAA:137:ASP:OD1	2.07	0.54
1:BBB:87:VAL:HG11	1:BBB:127:LEU:HD11	1.90	0.54
1:AAA:55:ILE:HD11	1:BBB:54:THR:O	2.07	0.53
1:AAA:67:ASP:HB3	1:AAA:79:ARG:HG3	1.90	0.53
1:AAA:64:LEU:HD22	1:BBB:58:ARG:HG3	1.92	0.50
1:AAA:68:ILE:HD13	1:AAA:78:PHE:HB3	1.93	0.49
1:AAA:78:PHE:HE2	1:AAA:92:ILE:HD13	1.79	0.46
1:BBB:81:VAL:HG13	2:BBB:201:DGP:C2	2.46	0.45
1:BBB:77:VAL:HG11	1:BBB:121:THR:HG21	1.99	0.45
1:AAA:68:ILE:CD1	1:AAA:78:PHE:HB3	2.46	0.44
1:BBB:122:ALA:O	1:BBB:127:LEU:HD12	2.17	0.44
1:BBB:98:PRO:HA	1:BBB:146:TYR:O	2.18	0.44
1:BBB:58:ARG:HB3	1:BBB:64:LEU:CB	2.49	0.43
1:BBB:96:PRO:HA	1:BBB:147:HIS:CE1	2.55	0.42
1:AAA:81:VAL:C	1:AAA:83:PRO:HD3	2.40	0.42
1:AAA:160:TRP:HA	1:AAA:161:PRO:C	2.40	0.42
1:BBB:73:GLN:OE1	1:BBB:73:GLN:HA	2.20	0.42
1:AAA:62:LYS:NZ	3:AAA:305:HOH:O	2.54	0.41
1:AAA:93:PRO:CD	1:AAA:148:LEU:O	2.68	0.40
1:AAA:58:ARG:NH2	2:BBB:201:DGP:N7	2.70	0.40

There are no symmetry-related clashes.



## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	110/163 (68%)	103 (94%)	7 (6%)	0	100	100
1	BBB	111/163 (68%)	103 (93%)	8 (7%)	0	100	100
All	All	221/326 (68%)	206 (93%)	15 (7%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	94/120 (78%)	85 (90%)	9 (10%)	8	22
1	BBB	95/120 (79%)	90 (95%)	5 (5%)	22	51
All	All	189/240 (79%)	175 (93%)	14 (7%)	13	34

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	62	LYS
1	AAA	81	VAL
1	AAA	92	ILE
1	AAA	106	GLU
1	AAA	109	GLN
1	AAA	119	LYS
1	AAA	123	LYS
1	AAA	144	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	AAA	158	LEU
1	BBB	57	SER
1	BBB	99	ARG
1	BBB	119	LYS
1	BBB	129	ASP
1	BBB	144	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be

highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	AAA	112/163 (68%)	0.15	1 (0%) 84 82	27, 41, 55, 67	0
1	BBB	112/163 (68%)	-0.06	2 (1%) 68 65	24, 35, 44, 53	0
All	All	224/326 (68%)	0.04	3 (1%) 77 75	24, 37, 52, 67	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	163	GLY	3.3
1	BBB	52	ALA	2.4
1	BBB	163	GLY	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	DGP	AAA	201	23/23	0.88	0.21	43,50,66,66	0
2	DGP	BBB	201	23/23	0.93	0.16	41,45,46,46	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around DGP AAA 201:**

2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around DGP BBB 201:**

$2mF_o - DF_c$  (at 0.7 rmsd) in gray  
 $mF_o - DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.