



Full wwPDB X-ray Structure Validation Report ⓘ

May 17, 2022 – 01:14 pm BST

PDB ID : 6YXL
Title : Crystal structure of ACPA F3
Authors : Ge, C.; Holmdahl, R.
Deposited on : 2020-05-03
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.28.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.28.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	HHH	221	<div> <div>11%</div> <div>85%</div> <div>9%</div> <div>• •</div> </div>
2	LLL	213	<div> <div>12%</div> <div>90%</div> <div>10%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	SO4	HHH	302	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6666 atoms, of which 3204 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ACPA F3 Fab fragment - heavy chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	HHH	214	Total	C	H	N	O	S	94	0	0
			3190	1013	1578	278	314	7			

- Molecule 2 is a protein called ACPA F3 Fab fragment - light chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	LLL	213	Total	C	H	N	O	S	102	0	0
			3268	1032	1618	287	327	4			

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	HHH	1	Total	C	H	O	2	0
			14	3	8	3		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	HHH	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	HHH	110	Total	O	0	0
			110	110		
5	LLL	79	Total	O	0	0
			79	79		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ACPA F3 Fab fragment - heavy chain



- Molecule 2: ACPA F3 Fab fragment - light chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	130.89Å 130.89Å 61.03Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	65.53 – 2.10 65.44 – 2.10	Depositor EDS
% Data completeness (in resolution range)	100.0 (65.53-2.10) 100.0 (65.44-2.10)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.02 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.8.0258, REFMAC 5.8.0258	Depositor
R, R_{free}	0.199 , 0.231 0.208 , 0.238	Depositor DCC
R_{free} test set	1551 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å ²)	42.0	Xtriage
Anisotropy	0.039	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6666	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	HHH	0.88	3/1650 (0.2%)	0.99	3/2252 (0.1%)
2	LLL	0.72	0/1685	0.93	0/2289
All	All	0.80	3/3335 (0.1%)	0.96	3/4541 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	HHH	0	4

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	HHH	55	HIS	C-N	12.71	1.63	1.34
1	HHH	29	VAL	C-N	11.78	1.61	1.34
1	HHH	127	SER	C-N	8.30	1.53	1.34

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	HHH	29	VAL	O-C-N	-8.61	108.92	122.70
1	HHH	27	PHE	CA-C-O	-6.19	107.09	120.10
1	HHH	55	HIS	O-C-N	-5.47	113.95	122.70

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	HHH	127	SER	Mainchain
1	HHH	29	VAL	Mainchain
1	HHH	55	HIS	Mainchain
1	HHH	69	LEU	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	HHH	1612	1578	1572	17	2
2	LLL	1650	1618	1616	11	1
3	HHH	6	8	8	0	0
4	HHH	5	0	0	0	2
5	HHH	110	0	0	1	0
5	LLL	79	0	0	1	0
All	All	3462	3204	3196	28	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:HHH:68:LYS:C	1:HHH:69:LEU:HA	1.67	1.15
1:HHH:69:LEU:HD12	1:HHH:69:LEU:N	1.90	0.86
1:HHH:68:LYS:C	1:HHH:69:LEU:CA	2.50	0.79
1:HHH:229:PRO:C	5:HHH:406:HOH:O	2.19	0.79
1:HHH:70:GLN:O	1:HHH:71:ASP:OD2	2.01	0.79
2:LLL:192:VAL:HG22	2:LLL:211:ASN:ND2	1.98	0.79
1:HHH:39:ARG:NH2	1:HHH:69:LEU:HD21	2.20	0.57
1:HHH:51:TRP:CE2	1:HHH:64:ASP:HB2	2.42	0.55
2:LLL:192:VAL:HG22	2:LLL:211:ASN:HD22	1.72	0.54
1:HHH:69:LEU:N	1:HHH:69:LEU:CD1	2.60	0.53
1:HHH:217:LYS:HB2	1:HHH:218:PRO:HD3	1.91	0.53
2:LLL:101:GLN:HB2	5:LLL:375:HOH:O	2.08	0.52
1:HHH:31:THR:O	1:HHH:55:HIS:HB2	2.12	0.49
1:HHH:215:ASN:HD22	1:HHH:216:HIS:N	2.11	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:HHH:1:GLN:OE1	1:HHH:1:GLN:N	2.37	0.48
2:LLL:192:VAL:HG22	2:LLL:211:ASN:HD21	1.75	0.48
2:LLL:137:LEU:HD12	2:LLL:137:LEU:N	2.32	0.45
2:LLL:184:LYS:O	2:LLL:188:GLU:HG2	2.18	0.44
2:LLL:187:TYR:HA	2:LLL:193:TYR:OH	2.18	0.43
2:LLL:121:PRO:HD3	2:LLL:133:VAL:HG22	2.00	0.43
1:HHH:152:ALA:O	1:HHH:199:THR:HA	2.19	0.43
1:HHH:201:PRO:O	1:HHH:204:SER:HB2	2.20	0.42
1:HHH:161:TYR:CE1	1:HHH:166:VAL:HG13	2.55	0.42
2:LLL:162:GLU:HA	2:LLL:177:SER:O	2.20	0.41
2:LLL:33:TYR:HB2	2:LLL:93:VAL:HB	2.02	0.41
2:LLL:141:TYR:CG	2:LLL:142:PRO:HA	2.56	0.41
1:HHH:215:ASN:HD22	1:HHH:215:ASN:C	2.25	0.40
1:HHH:70:GLN:O	1:HHH:71:ASP:CG	2.59	0.40

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:HHH:302:SO4:O3	4:HHH:302:SO4:O4[7_555]	1.36	0.84
1:HHH:116:TYR:OH	1:HHH:116:TYR:OH[8_555]	1.62	0.58
2:LLL:82:GLU:HB3	2:LLL:82:GLU:HB3[8_555]	1.23	0.37
1:HHH:19:ARG:HH11	1:HHH:19:ARG:HH11[7_556]	1.29	0.31
4:HHH:302:SO4:O1	4:HHH:302:SO4:O1[7_555]	2.08	0.12

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	HHH	206/221 (93%)	200 (97%)	5 (2%)	1 (0%)	29	26
2	LLL	211/213 (99%)	206 (98%)	5 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	417/434 (96%)	406 (97%)	10 (2%)	1 (0%)	47	49

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	HHH	71	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	HHH	180/186 (97%)	177 (98%)	3 (2%)	60	67
2	LLL	185/185 (100%)	181 (98%)	4 (2%)	52	57
All	All	365/371 (98%)	358 (98%)	7 (2%)	57	63

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	HHH	64	ASP
1	HHH	165	PRO
1	HHH	215	ASN
2	LLL	71	ASP
2	LLL	106	GLU
2	LLL	123	ASP
2	LLL	191	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SO4	HHH	302	-	4,4,4	0.40	0	6,6,6	0.23	0
3	GOL	HHH	301	-	5,5,5	0.45	0	5,5,5	0.73	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	HHH	301	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	HHH	301	GOL	O1-C1-C2-C3
3	HHH	301	GOL	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	HHH	302	SO4	0	2

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	HHH	4

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	HHH	27:PHE	C	29:VAL	N	3.79
1	HHH	68:LYS	C	69:LEU	N	3.74
1	HHH	55:HIS	C	60:ASN	N	1.63
1	HHH	29:VAL	C	31:THR	N	1.61

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	HHH	214/221 (96%)	1.13	25 (11%) 4 6	28, 40, 69, 82	0
2	LLL	213/213 (100%)	1.12	26 (12%) 4 5	34, 47, 62, 74	0
All	All	427/434 (98%)	1.12	51 (11%) 4 5	28, 44, 68, 82	0

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	HHH	1	GLN	5.2
2	LLL	81	PRO	4.2
1	HHH	207	THR	4.1
1	HHH	69	LEU	3.7
2	LLL	186	ASP	3.5
2	LLL	155	LEU	3.2
1	HHH	204	SER	3.1
1	HHH	141	PRO	2.9
2	LLL	193	TYR	2.7
1	HHH	29	VAL	2.7
2	LLL	24	ARG	2.7
1	HHH	116	TYR	2.7
1	HHH	133	GLY	2.6
2	LLL	150	LYS	2.6
1	HHH	70	GLN	2.6
1	HHH	214	VAL	2.6
2	LLL	21	LEU	2.5
2	LLL	148	GLN	2.5
2	LLL	127	LYS	2.5
1	HHH	209	THR	2.5
2	LLL	195	CYS	2.4
1	HHH	27	PHE	2.4
2	LLL	157	SER	2.4
2	LLL	187	TYR	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	LLL	25	ALA	2.4
1	HHH	71	ASP	2.4
1	HHH	140	ALA	2.3
1	HHH	206	GLY	2.3
2	LLL	182	LEU	2.3
2	LLL	128	SER	2.3
1	HHH	138	PRO	2.3
2	LLL	101	GLN	2.3
1	HHH	22	CYS	2.3
1	HHH	165	PRO	2.2
1	HHH	208	GLN	2.2
2	LLL	153	ASN	2.2
2	LLL	61	ASP	2.2
2	LLL	208	LYS	2.2
2	LLL	149	TRP	2.2
2	LLL	107	ILE	2.1
2	LLL	159	ASN	2.1
1	HHH	205	LEU	2.1
2	LLL	75	THR	2.1
1	HHH	199	THR	2.1
2	LLL	181	THR	2.1
1	HHH	37	TRP	2.1
1	HHH	44	GLN	2.1
2	LLL	154	ALA	2.1
1	HHH	34	PHE	2.0
1	HHH	150	GLY	2.0
2	LLL	160	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	GOL	HHH	301	6/6	0.62	0.29	30,56,61,62	2
4	SO4	HHH	302	5/5	0.87	0.39	38,44,82,113	2

6.5 Other polymers [i](#)

There are no such residues in this entry.