



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 8, 2020 – 09:24 AM BST

PDB ID : 4YZQ  
Title : C. bescii Family 3 pectate lyase double mutant K108A/Q111N in complex with trigalacturonic acid  
Authors : Alahuhta, P.M.; Lunin, V.V.  
Deposited on : 2015-03-25  
Resolution : 1.48 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.13.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.13.1


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4690 (1.50-1.46)
Clashscore	141614	4955 (1.50-1.46)
Ramachandran outliers	138981	4846 (1.50-1.46)
Sidechain outliers	138945	4844 (1.50-1.46)
RSRZ outliers	127900	4614 (1.50-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	204	<div> <div></div> <div>85% 11% . .</div> </div>
1	B	204	<div> <div></div> <div>90% 6% .</div> </div>
2	C	3	<div> <div></div> <div>100%</div> </div>
2	D	3	<div> <div></div> <div>100%</div> </div>

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 3846 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pectate lyase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	198	Total	C	N	O	S	0	13	0
			1598	998	275	318	7			
1	B	195	Total	C	N	O	S	0	8	0
			1535	962	261	306	6			

There are 26 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-10	MET	-	expression tag	UNP B9MKT4
A	-9	ALA	-	expression tag	UNP B9MKT4
A	-8	HIS	-	expression tag	UNP B9MKT4
A	-7	HIS	-	expression tag	UNP B9MKT4
A	-6	HIS	-	expression tag	UNP B9MKT4
A	-5	HIS	-	expression tag	UNP B9MKT4
A	-4	HIS	-	expression tag	UNP B9MKT4
A	-3	HIS	-	expression tag	UNP B9MKT4
A	-2	VAL	-	expression tag	UNP B9MKT4
A	-1	GLY	-	expression tag	UNP B9MKT4
A	0	THR	-	expression tag	UNP B9MKT4
A	108	ALA	LYS	engineered mutation	UNP B9MKT4
A	111	ASN	GLN	engineered mutation	UNP B9MKT4
B	-10	MET	-	expression tag	UNP B9MKT4
B	-9	ALA	-	expression tag	UNP B9MKT4
B	-8	HIS	-	expression tag	UNP B9MKT4
B	-7	HIS	-	expression tag	UNP B9MKT4
B	-6	HIS	-	expression tag	UNP B9MKT4
B	-5	HIS	-	expression tag	UNP B9MKT4
B	-4	HIS	-	expression tag	UNP B9MKT4
B	-3	HIS	-	expression tag	UNP B9MKT4
B	-2	VAL	-	expression tag	UNP B9MKT4
B	-1	GLY	-	expression tag	UNP B9MKT4
B	0	THR	-	expression tag	UNP B9MKT4
B	108	ALA	LYS	engineered mutation	UNP B9MKT4

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	111	ASN	GLN	engineered mutation	UNP B9MKT4

- Molecule 2 is an oligosaccharide called alpha-D-galactopyranuronic acid-(1-4)-alpha-D-galactopyranuronic acid-(1-4)-alpha-D-galactopyranuronic acid.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	C	3	Total	C	O	0	3	0
			37	18	19			
2	D	3	Total	C	O	0	3	0
			37	18	19			

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	4	Total	Ca	0	0
			4	4		
3	A	4	Total	Ca	0	0
			4	4		

- Molecule 4 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			8	6	2		
4	A	1	Total	C	O	0	0
			8	6	2		
4	A	1	Total	C	O	0	0
			8	6	2		
4	B	1	Total	C	O	0	0
			8	6	2		
4	B	1	Total	C	O	0	0
			8	6	2		

- Molecule 5 is beta-D-galactopyranuronic acid (three-letter code: GTR) (formula: C<sub>6</sub>H<sub>10</sub>O<sub>7</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	1
			13	6	7		
5	B	1	Total	C	O	0	1
			13	6	7		

- Molecule 6 is alpha-D-galactopyranuronic acid (three-letter code: ADA) (formula: C<sub>6</sub>H<sub>10</sub>O<sub>7</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	1
			12	6	6		
6	A	1	Total	C	O	0	1
			12	6	6		
6	B	1	Total	C	O	0	1
			12	6	6		
6	B	1	Total	C	O	0	1
			12	6	6		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	257	Total	O	0	9
			266	266		
7	B	246	Total	O	0	5
			251	251		



- Molecule 1: Pectate lyase




Chain B:  90% 6% .



Chain C:  100%



Chain D:  100%



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	137.73Å 36.21Å 99.47Å 90.00° 132.39° 90.00°	Depositor
Resolution (Å)	49.63 – 1.48 49.63 – 1.48	Depositor EDS
% Data completeness (in resolution range)	96.7 (49.63-1.48) 96.7 (49.63-1.48)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.76 (at 1.48Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, $R_{free}$	0.162 , 0.199 0.172 , 0.209	Depositor DCC
$R_{free}$ test set	2790 reflections (4.73%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	11.9	Xtriage
Anisotropy	0.188	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 59.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.015 for h,-k,-h-l	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	3846	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.26% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: MPD, GTR, CA, ADA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.96	1/1623 (0.1%)	0.95	3/2199 (0.1%)
1	B	0.98	1/1558 (0.1%)	0.92	1/2108 (0.0%)
All	All	0.97	2/3181 (0.1%)	0.93	4/4307 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	104	GLU	CD-OE2	-5.27	1.19	1.25
1	B	184	PHE	CG-CD1	5.21	1.46	1.38

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	21	ASP	CB-CG-OD1	8.65	126.09	118.30
1	A	10	ASP	CB-CG-OD1	7.93	125.43	118.30
1	B	21	ASP	CB-CG-OD1	6.77	124.39	118.30
1	A	49	LYS	CD-CE-NZ	5.50	124.34	111.70

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1598	0	1572	20	0
1	B	1535	0	1524	7	0
2	C	37	0	12	0	0
2	D	37	0	7	0	0
3	A	4	0	0	0	0
3	B	4	0	0	0	0
4	A	24	0	42	0	0
4	B	16	0	28	0	0
5	A	13	0	3	0	0
5	B	13	0	2	3	0
6	A	24	0	8	0	0
6	B	24	0	13	1	0
7	A	266	0	0	15	0
7	B	251	0	0	3	0
All	All	3846	0	3211	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 30 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:210[B]:GTR:C2	5:B:210[B]:GTR:C3	1.74	1.59
1:A:152[B]:ASN:ND2	7:A:302:HOH:O	1.80	1.11
1:A:122[B]:ASN:ND2	7:A:303:HOH:O	1.89	1.06
1:B:192[A]:GLN:OE1	7:B:302:HOH:O	1.86	0.93
1:A:30[A]:GLN:OE1	7:A:304:HOH:O	1.94	0.83

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	209/204 (102%)	200 (96%)	9 (4%)	0	100	100
1	B	201/204 (98%)	189 (94%)	12 (6%)	0	100	100
All	All	410/408 (100%)	389 (95%)	21 (5%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	180/172 (105%)	177 (98%)	3 (2%)	60	31
1	B	172/172 (100%)	171 (99%)	1 (1%)	86	72
All	All	352/344 (102%)	348 (99%)	4 (1%)	71	50

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	64	ASP
1	A	111	ASN
1	A	155	LYS
1	B	111	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	111	ASN
1	A	176	ASN
1	B	187	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	ADA	C	1[A]	2	10,13,13	1.02	0	15,19,19	1.36	2 (13%)
2	ADA	C	2[A]	3,2	9,12,13	1.36	1 (11%)	12,17,19	1.27	2 (16%)
2	ADA	C	3[A]	3,2	9,12,13	1.13	0	12,17,19	1.34	2 (16%)
2	ADA	D	1[A]	2	10,13,13	1.94	3 (30%)	15,19,19	1.77	3 (20%)
2	ADA	D	2[A]	3,2	9,12,13	3.89	4 (44%)	12,17,19	2.88	5 (41%)
2	ADA	D	3[A]	3,2	9,12,13	2.99	6 (66%)	12,17,19	1.56	1 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADA	C	1[A]	2	-	0/0/24/24	0/1/1/1
2	ADA	C	2[A]	3,2	-	0/0/21/24	0/1/1/1
2	ADA	C	3[A]	3,2	-	0/0/21/24	0/1/1/1
2	ADA	D	1[A]	2	-	0/0/24/24	0/1/1/1
2	ADA	D	2[A]	3,2	-	0/0/21/24	0/1/1/1
2	ADA	D	3[A]	3,2	-	0/0/21/24	0/1/1/1

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	2[A]	ADA	C2-C3	7.35	1.63	1.52
2	D	2[A]	ADA	O5-C5	7.28	1.51	1.43

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	3[A]	ADA	O3-C3	4.49	1.53	1.43
2	D	1[A]	ADA	O1-C1	4.26	1.53	1.39
2	D	3[A]	ADA	O5-C5	4.24	1.48	1.43

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	2[A]	ADA	O4-C4-C5	6.99	123.40	110.05
2	D	2[A]	ADA	C3-C4-C5	-5.22	98.30	109.02
2	D	3[A]	ADA	C1-C2-C3	-3.82	104.97	109.67
2	D	1[A]	ADA	C6-C5-C4	-3.30	104.79	113.04
2	D	1[A]	ADA	O2-C2-C3	-3.26	102.80	110.35


There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

Of 19 ligands modelled in this entry, 8 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	ADA	B	211[B]	3,5,6	9,12,13	4.59	7 (77%)	12,17,19	3.02	5 (41%)
4	MPD	A	206	-	7,7,7	0.52	0	9,10,10	0.34	0
4	MPD	B	206	-	7,7,7	0.49	0	9,10,10	1.01	0
6	ADA	A	213[B]	3,6	9,12,13	0.94	0	12,17,19	1.28	2 (16%)
6	ADA	B	212[B]	3,6	9,12,13	7.75	6 (66%)	12,17,19	1.37	2 (16%)
4	MPD	A	205	-	7,7,7	0.46	0	9,10,10	0.82	0
5	GTR	A	211[B]	6	10,13,13	1.28	1 (10%)	15,19,19	1.28	1 (6%)
5	GTR	B	210[B]	6	10,13,13	5.35	9 (90%)	15,19,19	1.92	4 (26%)
6	ADA	A	212[B]	3,5,6	9,12,13	1.12	0	12,17,19	1.36	3 (25%)
4	MPD	A	207	-	7,7,7	0.19	0	9,10,10	0.56	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	MPD	B	205	-	7,7,7	0.50	0	9,10,10	1.02	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	ADA	B	211[B]	3,5,6	-	0/0/21/24	0/1/1/1
4	MPD	A	206	-	-	2/5/5/5	-
4	MPD	B	206	-	-	1/5/5/5	-
6	ADA	A	213[B]	3,6	-	0/0/21/24	0/1/1/1
6	ADA	B	212[B]	3,6	-	0/0/21/24	0/1/1/1
4	MPD	A	205	-	-	1/5/5/5	-
5	GTR	A	211[B]	6	-	0/0/24/24	0/1/1/1
5	GTR	B	210[B]	6	-	0/0/24/24	0/1/1/1
6	ADA	A	212[B]	3,5,6	-	0/0/21/24	0/1/1/1
4	MPD	A	207	-	-	4/5/5/5	-
4	MPD	B	205	-	-	2/5/5/5	-

The worst 5 of 23 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	212[B]	ADA	O5-C5	11.43	1.55	1.43
6	B	212[B]	ADA	O5-C1	10.66	1.60	1.43
6	B	212[B]	ADA	C2-C3	10.57	1.68	1.52
5	B	210[B]	GTR	O5-C5	10.12	1.56	1.44
5	B	210[B]	GTR	C3-C2	8.86	1.74	1.52

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	211[B]	ADA	O5-C1-C2	7.14	121.80	110.77
5	B	210[B]	GTR	C1-O5-C5	4.77	119.44	112.31
6	B	211[B]	ADA	C3-C4-C5	4.50	118.26	109.02
6	B	211[B]	ADA	O4-C4-C3	-4.20	100.64	110.35
5	B	210[B]	GTR	O5-C5-C4	3.10	114.01	108.91

There are no chirality outliers.

5 of 10 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	207	MPD	O2-C2-C3-C4
4	B	206	MPD	O2-C2-C3-C4
4	B	205	MPD	C2-C3-C4-O4
4	A	207	MPD	C1-C2-C3-C4
4	A	207	MPD	CM-C2-C3-C4

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	211[B]	ADA	1	0
5	B	210[B]	GTR	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	198/204 (97%)	-0.59	3 (1%) 73 77	8, 12, 24, 49	3 (1%)
1	B	195/204 (95%)	-0.55	0 100 100	8, 14, 27, 39	0
All	All	393/408 (96%)	-0.57	3 (0%) 86 88	8, 13, 26, 49	3 (0%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	-2	VAL	3.2
1	A	-3	HIS	3.0
1	A	-4	HIS	2.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	ADA	D	1[A]	13/13	0.96	0.07	13,15,18,21	13
2	ADA	C	2[A]	12/13	0.97	0.06	12,15,17,17	12
2	ADA	C	1[A]	13/13	0.97	0.07	17,21,24,25	13
2	ADA	C	3[A]	12/13	0.97	0.06	12,14,17,18	12
2	ADA	D	3[A]	12/13	0.98	0.06	9,11,13,13	12
2	ADA	D	2[A]	12/13	0.98	0.06	11,11,12,12	12

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around Chain C:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around Chain D:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	MPD	B	205	8/8	0.69	0.19	32,38,42,42	0
4	MPD	A	206	8/8	0.88	0.13	30,34,36,41	0
4	MPD	B	206	8/8	0.88	0.12	29,38,40,42	8
4	MPD	A	205	8/8	0.92	0.10	31,33,36,39	0
4	MPD	A	207	8/8	0.92	0.10	23,28,28,30	8
5	GTR	A	211[B]	13/13	0.97	0.07	17,20,22,22	13
6	ADA	A	212[B]	12/13	0.97	0.06	14,16,19,19	12
5	GTR	B	210[B]	13/13	0.97	0.07	14,15,19,20	13
6	ADA	A	213[B]	12/13	0.97	0.06	14,16,19,20	12
6	ADA	B	211[B]	12/13	0.98	0.06	13,14,14,14	12
6	ADA	B	212[B]	12/13	0.98	0.06	13,15,17,17	12
3	CA	A	202	1/1	0.99	0.03	14,14,14,14	0
3	CA	A	201	1/1	1.00	0.03	15,15,15,15	1
3	CA	B	201	1/1	1.00	0.03	12,12,12,12	1
3	CA	A	204	1/1	1.00	0.02	11,11,11,11	0
3	CA	B	202	1/1	1.00	0.04	11,11,11,11	0
3	CA	A	203	1/1	1.00	0.03	12,12,12,12	0
3	CA	B	204	1/1	1.00	0.04	9,9,9,9	1
3	CA	B	203	1/1	1.00	0.04	12,12,12,12	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.