



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 18, 2020 – 01:47 am BST

PDB ID : 5Z16  
Title : A novel dimeric isocitrate dehydrogenase from *Acinetobacter baumannii*  
Authors : Song, P.; Huang, S.P.; Wang, P.; Zhu, G.P.  
Deposited on : 2017-12-25  
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	751	<div> <div>5%</div> <div>58%</div> <div>33%</div> <div>5% ..</div> </div>
1	B	751	<div> <div>74%</div> <div>23%</div> <div>..</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 11788 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Isocitrate dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	735	Total	C	N	O	S	0	0	0
			5666	3571	969	1091	35			
1	B	739	Total	C	N	O	S	0	0	0
			5741	3626	978	1102	35			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	HIS	-	expression tag	UNP V5VCI0
A	-4	HIS	-	expression tag	UNP V5VCI0
A	-3	HIS	-	expression tag	UNP V5VCI0
A	-2	HIS	-	expression tag	UNP V5VCI0
A	-1	HIS	-	expression tag	UNP V5VCI0
A	0	HIS	-	expression tag	UNP V5VCI0
A	629	THR	ALA	conflict	UNP V5VCI0
B	-5	HIS	-	expression tag	UNP V5VCI0
B	-4	HIS	-	expression tag	UNP V5VCI0
B	-3	HIS	-	expression tag	UNP V5VCI0
B	-2	HIS	-	expression tag	UNP V5VCI0
B	-1	HIS	-	expression tag	UNP V5VCI0
B	0	HIS	-	expression tag	UNP V5VCI0
B	629	THR	ALA	conflict	UNP V5VCI0

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mg	0	0
			1	1		
3	A	1	Total	Mg	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	201	Total	O	0	0
			201	201		
4	B	168	Total	O	0	0
			168	168		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Isocitrate dehydrogenase



H663	F664	Y665	L674	K684	F687	A693	N697	L705	A706	Q707	V708	K711	I715	Y718	V721	V726	V729	K730	R731	T742	V743	Q744	ALA																			
D540	T541	N547	R550	D551	E561	S570	E582	S588	V593	L596	V597	H601	L602	R603	E609	L613	V614	V615	E619	K623	N626	T629	K634	L636	D637	T640	L643	K648	R652	R653	T654	L657	D658	N659	R660	G661	S662					
L414	M415	A416	P432	I436	A437	L442	T450	Q451	E455	R460	M461	C462	D466	A467	P468	T469	R470	D471	W472	R479	S483	L491	D492	P493	H497	E498	N499	E500	K504	L509	T515	S523	R526	R529	Y530	T531	R534	V535	V536			
G285	K286	N294	V295	N296	A300	G301	L302	Y303	E304	K305	I306	T311	R314	I317	R328	S336	N342	M355	I359	R360	A361	M365	W366	G367	A368	D369	Y373	E381	S382	I387	Y388	Q389	E390	K396	N400	F401	D402	P403	K404	T408	V409	P410
T123	I124	R127	N137	P138	R141	E142	G143	N144	S145	R148	A149	P150	V153	S171	H174	V175	F183	S189	M190	T191	T204	N205	S206	I223	I224	D225	S226	M227	Y238	E247	I250	L254	K257	M260	M261	I267	F281	E282	K283	H284		

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 42 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	137.16 Å   137.16 Å   238.13 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	48.54 – 3.00 48.54 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.54-3.00) 100.0 (48.54-3.00)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.64 (at 3.01 Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, $R_{free}$	0.205   ,   0.276 0.209   ,   0.277	Depositor DCC
$R_{free}$ test set	2267 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	53.4	Xtriage
Anisotropy	0.008	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 44.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	11788	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.83% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.53	0/5773	0.82	17/7810 (0.2%)
1	B	0.48	0/5855	0.66	1/7923 (0.0%)
All	All	0.51	0/11628	0.74	18/15733 (0.1%)

There are no bond length outliers.

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	722	ASP	C-N-CD	-12.32	93.48	120.60
1	A	102	LEU	CA-CB-CG	9.56	137.28	115.30
1	A	683	LEU	CA-CB-CG	-8.43	95.91	115.30
1	A	689	PRO	N-CA-C	-7.31	93.08	112.10
1	A	709	GLN	N-CA-C	7.31	130.74	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5666	0	5620	356	0
1	B	5741	0	5699	120	0
2	A	5	0	0	1	0
2	B	5	0	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	201	0	0	22	0
4	B	168	0	0	21	0
All	All	11788	0	11319	474	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

The worst 5 of 474 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:24:LEU:CD2	1:A:663:HIS:ND1	1.81	1.43
1:A:24:LEU:HD23	1:A:663:HIS:ND1	1.14	1.40
1:A:96:THR:HA	1:A:99:ILE:CG2	1.56	1.35
1:A:688:ALA:C	1:A:690:LEU:H	1.04	1.35
1:A:96:THR:O	1:A:100:LYS:HD2	1.25	1.31

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	733/751 (98%)	626 (85%)	70 (10%)	37 (5%)	2	12
1	B	737/751 (98%)	673 (91%)	54 (7%)	10 (1%)	11	43
All	All	1470/1502 (98%)	1299 (88%)	124 (8%)	47 (3%)	4	22

5 of 47 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	114	GLU
1	A	117	ALA
1	A	647	ASP
1	A	654	THR
1	A	658	ASP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	601/628 (96%)	569 (95%)	32 (5%)	22	58
1	B	614/628 (98%)	587 (96%)	27 (4%)	28	65
All	All	1215/1256 (97%)	1156 (95%)	59 (5%)	25	61

5 of 59 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	682	GLU
1	B	100	LYS
1	B	609	GLU
1	A	708	VAL
1	A	725	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	646	ASN
1	B	342	ASN
1	A	677	GLN
1	A	160	HIS
1	B	296	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	PO4	B	801	-	4,4,4	0.97	0	6,6,6	0.64	0
2	PO4	A	801	-	4,4,4	0.87	0	6,6,6	0.64	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	801	PO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	735/751 (97%)	-0.13	39 (5%) 26 10	28, 50, 130, 141	0
1	B	739/751 (98%)	-0.61	0 100 100	22, 38, 64, 100	0
All	All	1474/1502 (98%)	-0.37	39 (2%) 56 27	22, 42, 125, 141	0

The worst 5 of 39 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	728	ALA	8.2
1	A	108	ALA	4.2
1	A	113	PRO	4.2
1	A	709	GLN	4.0
1	A	689	PRO	3.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	MG	B	802	1/1	0.94	0.33	38,38,38,38	0
3	MG	A	802	1/1	0.96	0.42	27,27,27,27	0
2	PO4	A	801	5/5	0.97	0.08	52,53,59,66	0
2	PO4	B	801	5/5	0.98	0.14	41,41,49,53	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.