



# Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 12:37 am BST

PDB ID : 5Z1Z  
Title : The apo-structure of D-lactate dehydrogenase from Escherichia coli  
Authors : Furukawa, N.; Miyanaga, A.; Nakajima, M.; Taguchi, H.  
Deposited on : 2017-12-28  
Resolution : 1.97 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	11647 (2.00-1.96)
Clashscore	141614	1014 (1.98-1.98)
Ramachandran outliers	138981	1006 (1.98-1.98)
Sidechain outliers	138945	1006 (1.98-1.98)
RSRZ outliers	127900	11410 (2.00-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	329	<div> <div>2%</div> <div>87%</div> <div>8%</div> <div>• •</div> </div>
1	B	329	<div> <div>4%</div> <div>88%</div> <div>7%</div> <div>• 5%</div> </div>
1	C	329	<div> <div>0%</div> <div>85%</div> <div>9%</div> <div>• •</div> </div>
1	D	329	<div> <div>6%</div> <div>85%</div> <div>8%</div> <div>• 5%</div> </div>

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 9901 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called D-isomer specific 2-hydroxyacid dehydrogenase NAD-binding.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	315	Total	C	N	O	S	0	0	0
			2446	1550	416	464	16			
1	B	314	Total	C	N	O	S	0	0	0
			2439	1545	415	463	16			
1	C	316	Total	C	N	O	S	0	0	0
			2454	1554	417	467	16			
1	D	311	Total	C	N	O	S	0	0	0
			2406	1524	406	460	16			

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	4	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	47	Total	O	0	0
			47	47		
4	B	41	Total	O	0	0
			41	41		
4	C	31	Total	O	0	0
			31	31		
4	D	29	Total	O	0	0
			29	29		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: D-isomer specific 2-hydroxyacid dehydrogenase NAD-binding




- Molecule 1: D-isomer specific 2-hydroxyacid dehydrogenase NAD-binding




- Molecule 1: D-isomer specific 2-hydroxyacid dehydrogenase NAD-binding



- Molecule 1: D-isomer specific 2-hydroxyacid dehydrogenase NAD-binding





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	139.60 Å   150.49 Å   62.76 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	48.24 – 1.97 48.20 – 1.97	Depositor EDS
% Data completeness (in resolution range)	96.3 (48.24-1.97) 96.3 (48.20-1.97)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.75 (at 1.97 Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, $R_{free}$	0.204   ,   0.227 0.212   ,   0.233	Depositor DCC
$R_{free}$ test set	4550 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	29.2	Xtriage
Anisotropy	0.047	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 44.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	9901	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.59% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.90	0/2487	0.85	0/3356
1	B	0.89	0/2480	0.89	0/3346
1	C	0.81	0/2495	0.83	0/3367
1	D	0.83	0/2446	0.85	0/3302
All	All	0.86	0/9908	0.86	0/13371

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2446	0	2447	23	0
1	B	2439	0	2438	22	0
1	C	2454	0	2451	26	0
1	D	2406	0	2403	32	0
2	A	1	0	0	0	0
3	A	7	0	10	3	0
4	A	47	0	0	0	0
4	B	41	0	0	0	0
4	C	31	0	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	D	29	0	0	2	0
All	All	9901	0	9749	89	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (89) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:260:TYR:HB2	1:D:263:GLU:OE1	1.62	0.99
1:A:260:TYR:HB2	1:A:263:GLU:CD	1.88	0.93
1:C:279:ASP:HB3	1:C:282:ARG:HD2	1.50	0.93
1:A:260:TYR:HB2	1:A:263:GLU:CG	1.99	0.92
1:B:283:ARG:NH1	1:B:287:CYS:SG	2.43	0.91
1:A:260:TYR:HB2	1:A:263:GLU:HG3	1.55	0.86
1:D:284:LEU:N	1:D:287:CYS:HG	1.72	0.86
1:D:210:GLU:OE1	1:D:210:GLU:N	2.12	0.81
1:D:260:TYR:HB2	1:D:263:GLU:CD	2.00	0.80
1:D:113:MET:SD	1:D:200:VAL:HG11	2.25	0.76
1:C:279:ASP:HB3	1:C:282:ARG:CD	2.17	0.74
1:A:324:CYS:HB3	1:A:327:GLU:HG3	1.69	0.74
1:D:209:PRO:HB2	1:D:210:GLU:OE1	1.93	0.68
1:D:261:GLU:HG2	1:D:295:HIS:HB3	1.76	0.67
1:D:284:LEU:N	1:D:287:CYS:SG	2.67	0.66
1:C:251:LYS:HA	1:C:251:LYS:HE2	1.78	0.66
1:D:181:ALA:HB1	1:D:186:VAL:HG11	1.77	0.65
1:C:1:MET:HG2	1:C:2:LYS:N	2.11	0.64
1:C:169:ARG:HH22	1:C:197:GLU:HG3	1.63	0.62
1:B:261:GLU:OE1	1:B:261:GLU:N	2.32	0.62
1:B:37:LYS:NZ	1:C:28:GLU:OE2	2.30	0.62
1:A:223:MET:O	1:A:251:LYS:CE	2.48	0.61
1:A:10:GLN:HB2	1:B:135:GLU:HG2	1.84	0.60
1:D:287:CYS:HB2	1:D:288:HIS:CD2	2.36	0.59
1:A:13:LYS:HE3	3:A:402:PEG:H41	1.84	0.58
1:D:53:ASN:OD1	1:D:262:ASN:HB2	2.03	0.58
1:A:283:ARG:HB2	1:C:287:CYS:SG	2.44	0.57
1:D:52:VAL:CG2	1:D:261:GLU:HB3	2.34	0.57
1:D:52:VAL:HG21	1:D:261:GLU:CB	2.35	0.57
1:A:223:MET:O	1:A:251:LYS:HE3	2.04	0.56
1:C:248:LYS:NZ	1:C:283:ARG:HE	2.03	0.56
1:B:286:ALA:HB2	1:D:286:ALA:CB	2.35	0.56

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:260:TYR:CB	1:A:263:GLU:HG3	2.31	0.56
1:C:52:VAL:HB	1:C:262:ASN:OD1	2.06	0.55
1:D:260:TYR:O	1:D:262:ASN:N	2.40	0.55
1:D:52:VAL:HG21	1:D:261:GLU:HB2	1.88	0.55
1:D:263:GLU:H	1:D:265:ASP:H	1.54	0.54
1:C:248:LYS:HZ1	1:C:283:ARG:HE	1.55	0.54
1:C:119:ARG:HG3	1:C:137:LEU:HA	1.88	0.54
1:B:284:LEU:H	1:B:284:LEU:HD12	1.73	0.53
1:C:249:ASN:O	1:C:250:GLN:HB2	2.09	0.53
1:B:150:ILE:HB	1:B:203:LEU:HD23	1.90	0.53
1:A:125:GLN:OE1	1:B:125:GLN:NE2	2.39	0.52
1:B:52:VAL:CG2	1:B:261:GLU:HB2	2.40	0.52
1:A:287:CYS:SG	1:C:283:ARG:HB2	2.50	0.51
1:B:119:ARG:HG3	1:B:137:LEU:HA	1.91	0.51
1:D:309:GLN:HG3	4:D:426:HOH:O	2.09	0.51
1:D:261:GLU:HG2	1:D:295:HIS:CB	2.41	0.51
1:D:260:TYR:CB	1:D:263:GLU:OE1	2.47	0.51
1:A:125:GLN:OE1	1:B:125:GLN:HG2	2.11	0.51
1:B:52:VAL:HG21	1:B:261:GLU:HB2	1.93	0.50
1:C:234:ARG:HD2	1:C:260:TYR:CE2	2.46	0.50
1:A:260:TYR:HB2	1:A:263:GLU:OE2	2.10	0.50
1:C:150:ILE:HB	1:C:203:LEU:HD23	1.94	0.50
1:C:234:ARG:NE	1:C:260:TYR:CD2	2.79	0.49
1:C:248:LYS:NZ	1:C:283:ARG:HH21	2.10	0.49
1:D:200:VAL:HG22	1:D:228:MET:HG2	1.93	0.49
1:B:176:TYR:CD1	1:B:176:TYR:N	2.81	0.49
1:C:264:ARG:O	1:C:265:ASP:HB2	2.13	0.49
1:B:286:ALA:CB	1:D:286:ALA:CB	2.90	0.49
1:B:286:ALA:HB2	1:D:286:ALA:HB1	1.95	0.49
1:D:309:GLN:CG	4:D:426:HOH:O	2.61	0.48
1:A:10:GLN:HG2	1:B:135:GLU:CD	2.34	0.48
1:A:84:LEU:HD22	1:A:329:VAL:HG23	1.97	0.47
1:D:113:MET:HG3	1:D:228:MET:SD	2.54	0.47
1:C:169:ARG:HH22	1:C:197:GLU:CG	2.26	0.47
1:D:150:ILE:HB	1:D:203:LEU:HD23	1.96	0.47
1:A:259:VAL:HG12	1:A:264:ARG:HG3	1.96	0.47
1:B:176:TYR:HD1	1:B:176:TYR:N	2.13	0.46
1:C:169:ARG:NH2	1:C:197:GLU:HG3	2.29	0.46
1:A:316:SER:O	1:A:320:LYS:HG3	2.15	0.45
1:B:32:PHE:HB3	1:C:32:PHE:HB3	1.99	0.44
1:B:248:LYS:HB2	1:B:248:LYS:HE2	1.86	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:135:GLU:OE1	1:B:10:GLN:HB2	2.18	0.44
1:A:283:ARG:HD2	1:C:283:ARG:HD2	2.00	0.43
1:C:234:ARG:CD	1:C:260:TYR:CE2	3.01	0.43
1:C:103:GLU:HG2	1:D:143:TYR:CG	2.54	0.43
1:D:124:TYR:O	1:D:128:ARG:CG	2.67	0.42
1:D:259:VAL:HG23	1:D:259:VAL:O	2.18	0.42
1:D:264:ARG:O	1:D:265:ASP:HB2	2.20	0.42
1:A:13:LYS:NZ	3:A:402:PEG:H12	2.34	0.42
1:D:262:ASN:OD1	1:D:265:ASP:HA	2.20	0.41
1:C:234:ARG:NE	1:C:260:TYR:CE2	2.87	0.41
1:B:259:VAL:HG23	1:B:259:VAL:O	2.20	0.41
1:A:320:LYS:HE2	1:A:322:GLU:OE1	2.21	0.41
1:B:328:LEU:O	1:B:329:VAL:HG13	2.20	0.41
1:A:10:GLN:OE1	3:A:402:PEG:O4	2.38	0.41
1:C:107:GLU:HA	1:C:110:ILE:HG22	2.03	0.40
1:D:52:VAL:CG2	1:D:261:GLU:CB	2.96	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	311/329 (94%)	306 (98%)	5 (2%)	0	100	100
1	B	310/329 (94%)	303 (98%)	7 (2%)	0	100	100
1	C	312/329 (95%)	306 (98%)	6 (2%)	0	100	100
1	D	307/329 (93%)	297 (97%)	10 (3%)	0	100	100
All	All	1240/1316 (94%)	1212 (98%)	28 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	262/276 (95%)	250 (95%)	12 (5%)	27	14
1	B	261/276 (95%)	252 (97%)	9 (3%)	37	25
1	C	263/276 (95%)	249 (95%)	14 (5%)	22	10
1	D	258/276 (94%)	248 (96%)	10 (4%)	32	19
All	All	1044/1104 (95%)	999 (96%)	45 (4%)	29	17

All (45) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	GLU
1	A	145	LYS
1	A	178	SER
1	A	183	GLU
1	A	203	LEU
1	A	213	HIS
1	A	228	MET
1	A	241	GLN
1	A	245	GLU
1	A	263	GLU
1	A	320	LYS
1	A	323	THR
1	B	145	LYS
1	B	176	TYR
1	B	178	SER
1	B	213	HIS
1	B	221	ASP
1	B	228	MET
1	B	264	ARG
1	B	283	ARG
1	B	284	LEU
1	C	80	ASN
1	C	129	ASP
1	C	145	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	178	SER
1	C	210	GLU
1	C	213	HIS
1	C	228	MET
1	C	234	ARG
1	C	241	GLN
1	C	248	LYS
1	C	251	LYS
1	C	261	GLU
1	C	262	ASN
1	C	279	ASP
1	D	80	ASN
1	D	128	ARG
1	D	145	LYS
1	D	178	SER
1	D	210	GLU
1	D	213	HIS
1	D	228	MET
1	D	241	GLN
1	D	265	ASP
1	D	287	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	PEG	A	402	-	6,6,6	0.35	0	5,5,5	0.61	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PEG	A	402	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	402	PEG	O2-C3-C4-O4
3	A	402	PEG	C1-C2-O2-C3

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	402	PEG	3	0

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	315/329 (95%)	0.26	8 (2%) 57 59	16, 32, 59, 76	0
1	B	314/329 (95%)	0.30	14 (4%) 33 35	15, 33, 65, 96	0
1	C	316/329 (96%)	0.33	2 (0%) 89 90	20, 38, 68, 79	0
1	D	311/329 (94%)	0.49	21 (6%) 17 19	21, 39, 67, 98	0
All	All	1256/1316 (95%)	0.35	45 (3%) 42 45	15, 35, 66, 98	0

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	287	CYS	6.1
1	B	286	ALA	5.6
1	B	262	ASN	5.1
1	D	286	ALA	5.1
1	D	262	ASN	4.6
1	B	284	LEU	4.3
1	D	321	GLY	4.1
1	D	323	THR	4.0
1	C	279	ASP	3.8
1	B	265	ASP	3.4
1	B	283	ARG	3.3
1	B	285	SER	3.3
1	D	173	PHE	3.3
1	A	176	TYR	3.2
1	A	324	CYS	3.2
1	D	241	GLN	3.2
1	D	288	HIS	3.2
1	D	221	ASP	3.2
1	A	321	GLY	3.2
1	D	261	GLU	3.0
1	B	263	GLU	2.9

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	287	CYS	2.9
1	D	176	TYR	2.8
1	B	261	GLU	2.7
1	D	218	ALA	2.6
1	B	260	TYR	2.6
1	D	210	GLU	2.6
1	A	323	THR	2.5
1	B	176	TYR	2.4
1	D	313	GLN	2.4
1	A	322	GLU	2.4
1	D	191	LEU	2.4
1	D	25	PHE	2.3
1	D	290	VAL	2.3
1	B	218	ALA	2.3
1	B	323	THR	2.3
1	D	240	SER	2.2
1	D	175	PRO	2.2
1	A	209	PRO	2.1
1	C	323	THR	2.1
1	D	320	LYS	2.1
1	D	284	LEU	2.1
1	A	329	VAL	2.1
1	A	208	THR	2.0
1	B	282	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	PEG	A	402	7/7	0.89	0.14	43,48,51,53	0
2	MG	A	401	1/1	0.97	0.05	39,39,39,39	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.