



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 15, 2020 – 05:26 am BST

PDB ID : 2Z2M  
Title : Cefditoren-Acylated Penicillin-Binding Protein 2X (PBP2X) from *Streptococcus pneumoniae*  
Authors : Yamada, M.; Watanabe, T.; Takeuchi, Y.  
Deposited on : 2007-05-23  
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	168	88% 7% 5%
1	D	168	57% 40%
2	B	385	85% 8% 6%
2	E	385	85% 8% 6%
3	C	125	86% 13%
3	F	125	93% 7%

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 9617 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Penicillin-binding protein 2X.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	159	Total	C	N	O	S	0	0	0
			1224	766	204	250	4			
1	D	100	Total	C	N	O	S	0	0	0
			743	459	126	155	3			

- Molecule 2 is a protein called Penicillin-binding protein 2X.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	363	Total	C	N	O	S	0	0	0
			2784	1752	459	556	17			
2	E	362	Total	C	N	O	S	0	0	0
			2778	1748	459	554	17			

- Molecule 3 is a protein called Penicillin-binding protein 2X.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	125	Total	C	N	O	S	0	0	0
			962	601	162	197	2			
3	F	125	Total	C	N	O	S	0	0	0
			962	601	162	197	2			

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is (2R)-2-[(1R)-1-{[(2Z)-2-(2-AMINO-1,3-THIAZOL-4-YL)-2-(METHOXYIMINO)ACETYL]AMINO}-2-OXOETHYL]-5-[(Z)-2-(4-METHYL-1,3-THIAZOL-5-YL)VINYL]-3,6-DIHYDRO-2H-1,3-THIAZINE-4-CARBOXYLIC ACID (three-letter code: CDS) (formula:  $C_{19}H_{20}N_6O_5S_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	N	O	S	
			33	19	6	5	3	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	E	1	Total	C	N	O	S	0	0
			33	19	6	5	3		

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	3	Total	O	0	0
			3	3		
6	B	30	Total	O	0	0
			30	30		
6	C	8	Total	O	0	0
			8	8		
6	D	13	Total	O	0	0
			13	13		
6	E	37	Total	O	0	0
			37	37		
6	F	2	Total	O	0	0
			2	2		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


#### • Molecule 1: Penicillin-binding protein 2X

Chain A: 





#### • Molecule 1: Penicillin-binding protein 2X

Chain D: 





#### • Molecule 2: Penicillin-binding protein 2X

Chain B: 




#### • Molecule 2: Penicillin-binding protein 2X

Chain E: 



- Molecule 3: Penicillin-binding protein 2X

Chain C:  86% 13% .



- Molecule 3: Penicillin-binding protein 2X

Chain F:  93% 7%



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	107.73Å 171.21Å 89.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.42 – 2.60	Depositor
% Data completeness (in resolution range)	89.8 (29.42-2.60)	Depositor
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.221 , 0.267	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	9617	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	46.0	wwPDB-VP



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, CDS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.35	0/1244	0.49	0/1676
1	D	0.36	0/754	0.51	0/1013
2	B	0.38	0/2837	0.51	0/3848
2	E	0.38	0/2831	0.52	0/3838
3	C	0.33	0/975	0.47	0/1320
3	F	0.34	0/975	0.47	0/1320
All	All	0.37	0/9616	0.50	0/13015

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1224	0	1191	7	0
1	D	743	0	708	1	0
2	B	2784	0	2717	18	0
2	E	2778	0	2712	17	0
3	C	962	0	975	11	0
3	F	962	0	975	5	0
4	B	5	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	33	0	18	0	0
5	E	33	0	18	0	0
6	A	3	0	0	0	0
6	B	30	0	0	0	0
6	C	8	0	0	0	0
6	D	13	0	0	0	0
6	E	37	0	0	0	0
6	F	2	0	0	0	0
All	All	9617	0	9314	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 52 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:327:ILE:HG22	2:E:331:SER:HB2	1.59	0.84
2:E:492:ARG:NH2	3:F:656:ASN:O	2.11	0.83
2:B:492:ARG:NH1	3:C:680:LEU:O	2.28	0.66
3:F:728:GLN:HG3	3:F:749:GLY:HA3	1.77	0.66
2:B:492:ARG:NH2	3:C:656:ASN:O	2.28	0.65

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	157/168 (94%)	151 (96%)	6 (4%)	0	100	100
1	D	94/168 (56%)	90 (96%)	4 (4%)	0	100	100
2	B	359/385 (93%)	347 (97%)	11 (3%)	1 (0%)	41	64

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	E	358/385 (93%)	344 (96%)	14 (4%)	0	100	100
3	C	123/125 (98%)	122 (99%)	1 (1%)	0	100	100
3	F	123/125 (98%)	118 (96%)	5 (4%)	0	100	100
All	All	1214/1356 (90%)	1172 (96%)	41 (3%)	1 (0%)	51	75

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	540	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	132/141 (94%)	130 (98%)	2 (2%)	65	83
1	D	80/141 (57%)	77 (96%)	3 (4%)	33	59
2	B	304/322 (94%)	297 (98%)	7 (2%)	50	75
2	E	303/322 (94%)	293 (97%)	10 (3%)	38	64
3	C	108/108 (100%)	105 (97%)	3 (3%)	43	69
3	F	108/108 (100%)	107 (99%)	1 (1%)	78	91
All	All	1035/1142 (91%)	1009 (98%)	26 (2%)	47	73

5 of 26 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	750	ASP
1	D	203	LEU
2	E	616	ASP
1	D	81	ASP
1	D	163	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	162	ASN
2	E	360	ASN
2	E	558	ASN
3	C	671	ASN
2	E	530	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
4	SO4	B	801	-	4,4,4	0.17	0	6,6,6	0.12	0
5	CDS	B	901	2	21,35,35	1.60	1 (4%)	16,48,48	3.05	9 (56%)
5	CDS	E	901	2	21,35,35	1.45	2 (9%)	16,48,48	2.91	7 (43%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	CDS	E	901	2	-	5/14/43/43	0/2/3/3
5	CDS	B	901	2	-	4/14/43/43	0/2/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	901	CDS	C10-C4	5.90	1.61	1.52
5	E	901	CDS	C10-C4	5.00	1.59	1.52
5	E	901	CDS	C29-S30	2.06	1.73	1.70

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	E	901	CDS	O23-C22-C24	5.47	126.66	120.35
5	B	901	CDS	C15-C14-C13	5.46	131.27	124.02
5	E	901	CDS	C15-C14-C13	4.98	130.62	124.02
5	B	901	CDS	O23-C22-C24	4.80	125.88	120.35
5	E	901	CDS	C6-C7-N21	4.74	120.31	109.98

There are no chirality outliers.

5 of 9 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	B	901	CDS	C14-C13-C3-C4
5	B	901	CDS	C14-C13-C3-C2
5	E	901	CDS	C24-N25-O26-C27
5	E	901	CDS	C14-C13-C3-C4
5	E	901	CDS	C14-C13-C3-C2


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates ⓘ

EDS was not executed - this section is therefore empty.

### 6.4 Ligands ⓘ

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers ⓘ

EDS was not executed - this section is therefore empty.