



wwPDB X-ray Structure Validation Summary Report ⓘ

May 11, 2022 – 04:08 pm BST

PDB ID : 7Z42
Title : Influenza B polymerase with Pol II pSer5 CTD peptide mimic bound in site 2B
Authors : Cusack, S.; Drncova, P.
Deposited on : 2022-03-03
Resolution : 2.42 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.28.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.28.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 2.42 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4647 (2.44-2.40)
Clashscore	141614	5161 (2.44-2.40)
Ramachandran outliers	138981	5073 (2.44-2.40)
Sidechain outliers	138945	5074 (2.44-2.40)
RSRZ outliers	127900	4543 (2.44-2.40)
RNA backbone	3102	1080 (2.80-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	751	
1	D	751	
2	B	772	
2	E	772	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	C	798	<div><div></div><div>16%</div><div>82%</div><div>8%</div><div>10%</div></div>
3	F	798	<div><div></div><div>19%</div><div>82%</div><div>8%</div><div>9%</div></div>
4	G	28	<div><div></div><div>11%</div><div>75%</div><div>25%</div></div>
4	I	28	<div><div></div><div>7%</div><div>64%</div><div>36%</div></div>
4	X	28	<div><div></div><div>11%</div><div>14%</div><div>86%</div></div>
4	Y	28	<div><div></div><div>4%</div><div>7%</div><div>89%</div></div>
5	H	13	<div><div></div><div>46%</div><div>46%</div><div>8%</div></div>
5	V	13	<div><div></div><div>46%</div><div>54%</div></div>

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 35949 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Polymerase acidic protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	721	Total	C	N	O	S	0	1	0
			5787	3675	970	1101	41			
1	D	715	Total	C	N	O	S	0	1	0
			5750	3652	962	1095	41			

There are 50 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-13	GLY	-	expression tag	UNP Q5V8Z9
A	-12	SER	-	expression tag	UNP Q5V8Z9
A	-11	HIS	-	expression tag	UNP Q5V8Z9
A	-10	HIS	-	expression tag	UNP Q5V8Z9
A	-9	HIS	-	expression tag	UNP Q5V8Z9
A	-8	HIS	-	expression tag	UNP Q5V8Z9
A	-7	HIS	-	expression tag	UNP Q5V8Z9
A	-6	HIS	-	expression tag	UNP Q5V8Z9
A	-5	HIS	-	expression tag	UNP Q5V8Z9
A	-4	HIS	-	expression tag	UNP Q5V8Z9
A	-3	GLY	-	expression tag	UNP Q5V8Z9
A	-2	SER	-	expression tag	UNP Q5V8Z9
A	-1	GLY	-	expression tag	UNP Q5V8Z9
A	0	SER	-	expression tag	UNP Q5V8Z9
A	727	GLY	-	expression tag	UNP Q5V8Z9
A	728	SER	-	expression tag	UNP Q5V8Z9
A	729	GLY	-	expression tag	UNP Q5V8Z9
A	730	SER	-	expression tag	UNP Q5V8Z9
A	731	GLY	-	expression tag	UNP Q5V8Z9
A	732	GLU	-	expression tag	UNP Q5V8Z9
A	733	ASN	-	expression tag	UNP Q5V8Z9
A	734	LEU	-	expression tag	UNP Q5V8Z9
A	735	TYR	-	expression tag	UNP Q5V8Z9
A	736	PHE	-	expression tag	UNP Q5V8Z9
A	737	GLN	-	expression tag	UNP Q5V8Z9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-13	GLY	-	expression tag	UNP Q5V8Z9
D	-12	SER	-	expression tag	UNP Q5V8Z9
D	-11	HIS	-	expression tag	UNP Q5V8Z9
D	-10	HIS	-	expression tag	UNP Q5V8Z9
D	-9	HIS	-	expression tag	UNP Q5V8Z9
D	-8	HIS	-	expression tag	UNP Q5V8Z9
D	-7	HIS	-	expression tag	UNP Q5V8Z9
D	-6	HIS	-	expression tag	UNP Q5V8Z9
D	-5	HIS	-	expression tag	UNP Q5V8Z9
D	-4	HIS	-	expression tag	UNP Q5V8Z9
D	-3	GLY	-	expression tag	UNP Q5V8Z9
D	-2	SER	-	expression tag	UNP Q5V8Z9
D	-1	GLY	-	expression tag	UNP Q5V8Z9
D	0	SER	-	expression tag	UNP Q5V8Z9
D	727	GLY	-	expression tag	UNP Q5V8Z9
D	728	SER	-	expression tag	UNP Q5V8Z9
D	729	GLY	-	expression tag	UNP Q5V8Z9
D	730	SER	-	expression tag	UNP Q5V8Z9
D	731	GLY	-	expression tag	UNP Q5V8Z9
D	732	GLU	-	expression tag	UNP Q5V8Z9
D	733	ASN	-	expression tag	UNP Q5V8Z9
D	734	LEU	-	expression tag	UNP Q5V8Z9
D	735	TYR	-	expression tag	UNP Q5V8Z9
D	736	PHE	-	expression tag	UNP Q5V8Z9
D	737	GLN	-	expression tag	UNP Q5V8Z9

- Molecule 2 is a protein called RNA-directed RNA polymerase catalytic subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	754	Total	C	N	O	S	0	0	0
			5911	3731	1024	1104	52			
2	E	743	Total	C	N	O	S	0	0	0
			5818	3671	1004	1091	52			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-8	GLY	-	expression tag	UNP Q5V8Y6
B	-7	SER	-	expression tag	UNP Q5V8Y6
B	-6	GLY	-	expression tag	UNP Q5V8Y6
B	-5	SER	-	expression tag	UNP Q5V8Y6
B	-4	GLY	-	expression tag	UNP Q5V8Y6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-3	SER	-	expression tag	UNP Q5V8Y6
B	-2	GLY	-	expression tag	UNP Q5V8Y6
B	-1	SER	-	expression tag	UNP Q5V8Y6
B	0	GLY	-	expression tag	UNP Q5V8Y6
B	753	GLY	-	expression tag	UNP Q5V8Y6
B	754	SER	-	expression tag	UNP Q5V8Y6
B	755	GLY	-	expression tag	UNP Q5V8Y6
B	756	SER	-	expression tag	UNP Q5V8Y6
B	757	GLY	-	expression tag	UNP Q5V8Y6
B	758	GLU	-	expression tag	UNP Q5V8Y6
B	759	ASN	-	expression tag	UNP Q5V8Y6
B	760	LEU	-	expression tag	UNP Q5V8Y6
B	761	TYR	-	expression tag	UNP Q5V8Y6
B	762	PHE	-	expression tag	UNP Q5V8Y6
B	763	GLN	-	expression tag	UNP Q5V8Y6
E	-8	GLY	-	expression tag	UNP Q5V8Y6
E	-7	SER	-	expression tag	UNP Q5V8Y6
E	-6	GLY	-	expression tag	UNP Q5V8Y6
E	-5	SER	-	expression tag	UNP Q5V8Y6
E	-4	GLY	-	expression tag	UNP Q5V8Y6
E	-3	SER	-	expression tag	UNP Q5V8Y6
E	-2	GLY	-	expression tag	UNP Q5V8Y6
E	-1	SER	-	expression tag	UNP Q5V8Y6
E	0	GLY	-	expression tag	UNP Q5V8Y6
E	753	GLY	-	expression tag	UNP Q5V8Y6
E	754	SER	-	expression tag	UNP Q5V8Y6
E	755	GLY	-	expression tag	UNP Q5V8Y6
E	756	SER	-	expression tag	UNP Q5V8Y6
E	757	GLY	-	expression tag	UNP Q5V8Y6
E	758	GLU	-	expression tag	UNP Q5V8Y6
E	759	ASN	-	expression tag	UNP Q5V8Y6
E	760	LEU	-	expression tag	UNP Q5V8Y6
E	761	TYR	-	expression tag	UNP Q5V8Y6
E	762	PHE	-	expression tag	UNP Q5V8Y6
E	763	GLN	-	expression tag	UNP Q5V8Y6

- Molecule 3 is a protein called Polymerase basic protein 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	718	Total	C	N	O	S	0	0	0
			5746	3654	1001	1051	40			
3	F	728	Total	C	N	O	S	0	1	0
			5838	3712	1023	1063	40			

There are 56 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-8	GLY	-	expression tag	UNP Q5V8X3
C	-7	SER	-	expression tag	UNP Q5V8X3
C	-6	GLY	-	expression tag	UNP Q5V8X3
C	-5	SER	-	expression tag	UNP Q5V8X3
C	-4	GLY	-	expression tag	UNP Q5V8X3
C	-3	SER	-	expression tag	UNP Q5V8X3
C	-2	GLY	-	expression tag	UNP Q5V8X3
C	-1	SER	-	expression tag	UNP Q5V8X3
C	0	GLY	-	expression tag	UNP Q5V8X3
C	771	GLY	-	expression tag	UNP Q5V8X3
C	772	TRP	-	expression tag	UNP Q5V8X3
C	773	SER	-	expression tag	UNP Q5V8X3
C	774	HIS	-	expression tag	UNP Q5V8X3
C	775	PRO	-	expression tag	UNP Q5V8X3
C	776	GLN	-	expression tag	UNP Q5V8X3
C	777	PHE	-	expression tag	UNP Q5V8X3
C	778	GLU	-	expression tag	UNP Q5V8X3
C	779	LYS	-	expression tag	UNP Q5V8X3
C	780	GLY	-	expression tag	UNP Q5V8X3
C	781	SER	-	expression tag	UNP Q5V8X3
C	782	GLY	-	expression tag	UNP Q5V8X3
C	783	SER	-	expression tag	UNP Q5V8X3
C	784	GLU	-	expression tag	UNP Q5V8X3
C	785	ASN	-	expression tag	UNP Q5V8X3
C	786	LEU	-	expression tag	UNP Q5V8X3
C	787	TYR	-	expression tag	UNP Q5V8X3
C	788	PHE	-	expression tag	UNP Q5V8X3
C	789	GLN	-	expression tag	UNP Q5V8X3
F	-8	GLY	-	expression tag	UNP Q5V8X3
F	-7	SER	-	expression tag	UNP Q5V8X3
F	-6	GLY	-	expression tag	UNP Q5V8X3
F	-5	SER	-	expression tag	UNP Q5V8X3
F	-4	GLY	-	expression tag	UNP Q5V8X3
F	-3	SER	-	expression tag	UNP Q5V8X3
F	-2	GLY	-	expression tag	UNP Q5V8X3
F	-1	SER	-	expression tag	UNP Q5V8X3
F	0	GLY	-	expression tag	UNP Q5V8X3
F	771	GLY	-	expression tag	UNP Q5V8X3
F	772	TRP	-	expression tag	UNP Q5V8X3
F	773	SER	-	expression tag	UNP Q5V8X3
F	774	HIS	-	expression tag	UNP Q5V8X3
F	775	PRO	-	expression tag	UNP Q5V8X3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	776	GLN	-	expression tag	UNP Q5V8X3
F	777	PHE	-	expression tag	UNP Q5V8X3
F	778	GLU	-	expression tag	UNP Q5V8X3
F	779	LYS	-	expression tag	UNP Q5V8X3
F	780	GLY	-	expression tag	UNP Q5V8X3
F	781	SER	-	expression tag	UNP Q5V8X3
F	782	GLY	-	expression tag	UNP Q5V8X3
F	783	SER	-	expression tag	UNP Q5V8X3
F	784	GLU	-	expression tag	UNP Q5V8X3
F	785	ASN	-	expression tag	UNP Q5V8X3
F	786	LEU	-	expression tag	UNP Q5V8X3
F	787	TYR	-	expression tag	UNP Q5V8X3
F	788	PHE	-	expression tag	UNP Q5V8X3
F	789	GLN	-	expression tag	UNP Q5V8X3

- Molecule 4 is a protein called DNA-directed RNA polymerase II subunit RPB1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	Y	3	Total	C	N	O	P	0	0	0
			24	12	3	8	1			
4	G	21	Total	C	N	O	P	0	0	0
			165	96	21	45	3			
4	X	4	Total	C	N	O	P	0	0	0
			30	15	4	10	1			
4	I	18	Total	C	N	O	P	0	0	0
			142	85	18	37	2			

- Molecule 5 is a RNA chain called RNA (5'-R(P*AP*GP*UP*AP*GP*UP*AP*AP*CP*AP*AP*GP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	H	13	Total	C	N	O	P	0	0	0
			266	117	52	84	13			
5	V	13	Total	C	N	O	P	0	0	0
			266	117	52	84	13			


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	43	Total	O	0	0
			43	43		


Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	30	Total 30	O 30	0	0
6	C	20	Total 20	O 20	0	0
6	D	39	Total 39	O 39	0	0
6	E	43	Total 43	O 43	0	0
6	F	16	Total 16	O 16	0	0
6	G	1	Total 1	O 1	0	0
6	H	6	Total 6	O 6	0	0
6	I	1	Total 1	O 1	0	0
6	V	7	Total 7	O 7	0	0




• Molecule 2: RNA-directed RNA polymerase catalytic subunit



• Molecule 3: Polymerase basic protein 2





- Molecule 5: RNA (5'-R(P*AP*GP*UP*AP*GP*UP*AP*AP*CP*AP*AP*GP*A)-3')



- Molecule 5: RNA (5'-R(P*AP*GP*UP*AP*GP*UP*AP*AP*CP*AP*AP*GP*A)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	130.93Å 202.27Å 135.73Å 90.00° 110.54° 90.00°	Depositor
Resolution (Å)	127.11 – 2.42 127.11 – 2.42	Depositor EDS
% Data completeness (in resolution range)	61.4 (127.11-2.42) 61.4 (127.11-2.42)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.38 (at 2.42Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.223 , 0.257 0.224 , 0.256	Depositor DCC
R_{free} test set	7573 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	44.1	Xtriage
Anisotropy	0.059	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.001 for l,-k,h	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	35949	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.67	0/5903	0.71	0/7959
1	D	0.67	0/5866	0.70	0/7910
2	B	0.67	0/6028	0.71	0/8127
2	E	0.67	0/5931	0.71	0/7995
3	C	0.67	0/5844	0.72	0/7852
3	F	0.67	0/5939	0.72	0/7982
4	G	0.60	0/140	0.63	0/189
4	I	0.62	0/127	0.63	0/173
4	X	0.71	0/20	0.59	0/27
4	Y	0.57	0/14	0.80	0/18
5	H	0.93	1/298 (0.3%)	1.26	3/462 (0.6%)
5	V	0.93	1/298 (0.3%)	1.25	2/462 (0.4%)
All	All	0.67	2/36408 (0.0%)	0.73	5/49156 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	V	1	A	OP3-P	-7.51	1.52	1.61
5	H	1	A	OP3-P	-7.43	1.52	1.61

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	V	5	G	P-O3'-C3'	-6.80	111.54	119.70
5	H	12	G	P-O3'-C3'	-6.47	111.93	119.70
5	H	5	G	P-O3'-C3'	-6.04	112.45	119.70
5	H	6	U	P-O3'-C3'	-5.63	112.95	119.70
5	V	6	U	P-O3'-C3'	-5.56	113.03	119.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5787	0	5761	33	0
1	D	5750	0	5717	28	0
2	B	5911	0	5932	30	0
2	E	5818	0	5824	34	0
3	C	5746	0	5909	25	0
3	F	5838	0	6007	33	0
4	G	165	0	130	0	0
4	I	142	0	115	0	0
4	X	30	0	21	0	0
4	Y	24	0	17	1	0
5	H	266	0	130	1	0
5	V	266	0	130	1	0
6	A	43	0	0	0	0
6	B	30	0	0	0	0
6	C	20	0	0	0	0
6	D	39	0	0	0	0
6	E	43	0	0	1	0
6	F	16	0	0	0	0
6	G	1	0	0	0	0
6	H	6	0	0	0	0
6	I	1	0	0	0	0
6	V	7	0	0	0	0
All	All	35949	0	35693	160	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 160 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:51:MET:O	1:A:62:THR:HA	1.92	0.70
2:B:282:LEU:HD22	2:B:440:LEU:HD13	1.76	0.66

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:F:717:LEU:HD21	3:F:728:ILE:HD11	1.77	0.66
2:E:282:LEU:HD22	2:E:440:LEU:HD13	1.78	0.66
3:F:273:ILE:HD11	3:F:516:ALA:CB	2.26	0.66

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	718/751 (96%)	691 (96%)	26 (4%)	1 (0%)	51	67
1	D	712/751 (95%)	688 (97%)	24 (3%)	0	100	100
2	B	752/772 (97%)	726 (96%)	26 (4%)	0	100	100
2	E	737/772 (96%)	714 (97%)	23 (3%)	0	100	100
3	C	712/798 (89%)	678 (95%)	34 (5%)	0	100	100
3	F	725/798 (91%)	693 (96%)	31 (4%)	1 (0%)	51	67
4	G	16/28 (57%)	16 (100%)	0	0	100	100
4	I	14/28 (50%)	13 (93%)	1 (7%)	0	100	100
4	X	2/28 (7%)	2 (100%)	0	0	100	100
4	Y	1/28 (4%)	1 (100%)	0	0	100	100
All	All	4389/4754 (92%)	4222 (96%)	165 (4%)	2 (0%)	100	100

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	65	GLU
3	F	89	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	641/664 (96%)	624 (97%)	17 (3%)	44	63
1	D	638/664 (96%)	613 (96%)	25 (4%)	32	49
2	B	646/657 (98%)	620 (96%)	26 (4%)	31	48
2	E	636/657 (97%)	614 (96%)	22 (4%)	36	53
3	C	628/694 (90%)	597 (95%)	31 (5%)	25	39
3	F	637/694 (92%)	609 (96%)	28 (4%)	28	43
4	G	18/24 (75%)	18 (100%)	0	100	100
4	I	16/24 (67%)	16 (100%)	0	100	100
4	X	3/24 (12%)	3 (100%)	0	100	100
4	Y	2/24 (8%)	2 (100%)	0	100	100
All	All	3865/4126 (94%)	3716 (96%)	149 (4%)	32	49

5 of 149 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	E	562	ARG
3	F	607	GLN
2	E	669	ARG
3	F	368	ARG
3	C	227	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	590	GLN
2	E	310	ASN
3	F	325	GLN
2	B	613	ASN
1	A	319	GLN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
5	H	11/13 (84%)	3 (27%)	0
5	V	11/13 (84%)	3 (27%)	0
All	All	22/26 (84%)	6 (27%)	0

5 of 6 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
5	H	4	A
5	H	7	A
5	H	11	A
5	V	4	A
5	V	7	A

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

7 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SEP	X	26	4	8,9,10	0.55	0	8,12,14	0.68	0
4	SEP	G	47	4	8,9,10	0.54	0	8,12,14	0.64	0
4	SEP	G	40	4	8,9,10	0.72	0	8,12,14	0.63	0
4	SEP	Y	26	4	8,9,10	0.57	0	8,12,14	0.64	0
4	SEP	I	40	4	8,9,10	0.51	0	8,12,14	0.64	0
4	SEP	G	33	4	8,9,10	0.56	0	8,12,14	0.64	0
4	SEP	I	33	4	8,9,10	0.56	0	8,12,14	0.62	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SEP	X	26	4	-	1/5/8/10	-
4	SEP	G	47	4	-	2/5/8/10	-
4	SEP	G	40	4	-	4/5/8/10	-
4	SEP	Y	26	4	-	1/5/8/10	-
4	SEP	I	40	4	-	0/5/8/10	-
4	SEP	G	33	4	-	2/5/8/10	-
4	SEP	I	33	4	-	2/5/8/10	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 12 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	G	40	SEP	CB-OG-P-O1P
4	G	40	SEP	CB-OG-P-O2P
4	G	40	SEP	CB-OG-P-O3P
4	Y	26	SEP	N-CA-CB-OG
4	X	26	SEP	N-CA-CB-OG

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	Y	26	SEP	1	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	721/751 (96%)	0.61	48 (6%) 17 15	21, 40, 126, 149	0
1	D	715/751 (95%)	0.45	19 (2%) 54 52	23, 39, 116, 139	0
2	B	754/772 (97%)	0.37	17 (2%) 60 57	23, 45, 76, 116	0
2	E	743/772 (96%)	0.40	14 (1%) 66 64	23, 42, 69, 110	0
3	C	718/798 (89%)	1.31	124 (17%) 1 1	25, 56, 185, 226	0
3	F	728/798 (91%)	1.50	149 (20%) 1 0	26, 54, 172, 193	0
4	G	18/28 (64%)	1.09	3 (16%) 1 1	46, 68, 131, 137	0
4	I	16/28 (57%)	0.87	2 (12%) 3 3	46, 63, 75, 89	0
4	X	3/28 (10%)	4.84	3 (100%) 0 0	94, 94, 96, 97	0
4	Y	2/28 (7%)	3.35	1 (50%) 0 0	98, 98, 98, 98	0
5	H	13/13 (100%)	-0.05	0 100 100	25, 29, 56, 69	0
5	V	13/13 (100%)	-0.02	0 100 100	25, 28, 40, 47	0
All	All	4444/4780 (92%)	0.77	380 (8%) 10 9	21, 46, 139, 226	0

The worst 5 of 380 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	F	425	ARG	28.0
3	F	423	LEU	24.7
3	C	460	SER	21.4
3	F	426	ALA	19.3
3	C	431	SER	18.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	SEP	G	47	10/11	0.64	0.34	124,135,140,141	0
4	SEP	Y	26	10/11	0.81	0.24	79,84,91,92	0
4	SEP	X	26	10/11	0.83	0.20	79,83,87,89	0
4	SEP	G	40	10/11	0.91	0.12	64,67,72,72	0
4	SEP	G	33	10/11	0.94	0.13	49,53,54,55	0
4	SEP	I	40	10/11	0.94	0.12	65,71,84,84	0
4	SEP	I	33	10/11	0.95	0.10	55,58,61,62	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.